



Contribution ID: 1046

Type: **Parallel Talk**

## Strong First-Order Electroweak Phase Transitions in the Standard Model with a Singlet Extension

Monday 4 May 2020 15:15 (15 minutes)

A common assumption about the early universe is that it underwent an electroweak phase transition (EWPT). Though the standard model (SM) is able to restore the electroweak symmetry through a smooth cross over PT, we require a strongly first-order PT to ensure electroweak baryogenesis, requiring us to look at new physics beyond the SM. The simplest case to extend the SM is to add a real singlet field, which allows first-order EWPTs (FOEPT) to occur.

Starting with the most general higgs+singlet lagrangian, we then fixed four of its coupling constants as functions of parameters whose range of values had more experimental motivation. Then by requiring a FOEPT and performing a Monte-Carlo scan over five free parameters, we were able to study the parameter space in this allowed region. Most notably, we observed the triple higgs coupling ( $\lambda_3$ ) take on values between 1.2 and 2.5. The possible values of  $\lambda_3$  could serve as motivation for future collider experiments to improve sensitivity in this range when looking at the cross sections of  $pp \rightarrow hh$  versus  $\lambda_3$ .

### Summary

**Primary authors:** Mr HOOPER, Anthony (University of Nebraska-Lincoln); Dr HUANG, Peisi (University of Nebraska-Lincoln)

**Presenter:** Mr HOOPER, Anthony (University of Nebraska-Lincoln)

**Session Classification:** BSM I

**Track Classification:** BSM