


LEARNING PHYSICS AT FUTURE LEPTON COLLIDERS WITH MACHINE

Lingfeng Li

Hong Kong University of Science and Technology



Pheno

May 4, 2020, Pittsburgh

Based on arXiv:2004.15013

in collaboration with Ying-Ying Li, Tao Liu and Sijun Xu

Precision Frontier of Next Decades

Led by future ee colliders (FCC-ee, CEPC, CLIC, ILC)
, measuring Higgs and EW precisely.

[A. Abada et al., (2019); H. Abramowicz et al., 1608.07538, F. An et al., 1810.09037,...]

Primary Higgs and electroweak processes


Jet Number	0	2	4	6
$e^-e^+ \rightarrow WW$	11%	44%	45%	0%
$e^-e^+ \rightarrow ZZ$	9%	42%	49%	0%
$e^-e^+ \rightarrow ZH$	3%	32%	55%	11%
$e^-e^+ \rightarrow H\nu\nu$	20%	69%	11%	0%
$e^-e^+ \rightarrow t\bar{t}$	0%	11%	44%	45%

Hadronic mode dominant

How would jet clustering affect the precisions?

Limitations of jet clustering

Hadrons from different Z clustered in a same jet (info distortion)



Detailed structures are gone after clustering (info loss)

Can we recover from these limitations?

First Way: Jet +Event-Level Obs.

- Jet substructure observables: extensively applied in boost kinematics
- Event shape: relatively intuition-based, e.g. thrust [[E. Farhi, 1977](#)]
- Fox-Wolfram moments [[G. C. Fox and S. Wolfram, 1978](#)] and their extensions: more systematic, but relatively less intuitive.

$$H_{AB;l} = \sum_{m=-l}^l H_{AB;l,m} = \frac{4\pi}{2l+1} \sum_{i,j} \frac{A_i B_j}{s} \sum_{m=-l}^l (Y_l^m(\Omega_i)^* Y_l^m(\Omega_j)) = \sum_{i,j} \frac{A_i B_j}{s} P_l(\cos \Omega_{ij})$$

- Pros: Simple framework. Physically intuitive.
- Cons: Less organized.

Another Way: Event-Level ML


Pursue analysis directly at event level

- Pro: Most information.
 - Lepton Collider: negligible pileups, colorless beam and fixed energy
- Con: Large complexity. -> ML as a solution.

Comparative studies to compare the two approaches using ML as a tool

- Jet Level: Fully Connected Network (FCN) :
Input: jet momenta (and FW moments $l \leq 50$ / track info).
- Event Level: Convolutional Neuron Network (CNN)
Based on ResNet-50 structure.
Input: 50×50 pixelized event-level image (and track info).

Cumulative Mollweide Projection





Back to back motion of qq

Halo size and structure: minimal included angle of quarks/ information missing at jet level

- Define a Cartesian coordinate system: z-axis being along beam line and x - y plane (equatorial plane) overlapping with its transverse plane
- Rotate the motion direction of the most energetic particle to be along x-axis.
- Project the particles to ``detector sphere''


“Dictionary” between Event Projection and CMB



Mollweide projection at e^-e^+ colliders	All-sky CMB map
Projection sphere	Celestial sphere
Equatorial plane	Galactic plane
Energy (p_T , timing, charge, d_0 , etc.) projection	Temperature (polarization) map
Event-level kinematics	Anisotropy
Fox-Wolfram moments	Power spectrum (TT , TB , BB , etc.)
Multi-spectra	Bispectrum, trispectrum, etc.
...

In such CMB-like information scheme, the event-level information is encoded as the FW moments at leading order and multi-spectra at higher orders.


Benchmark




$$e^-e^+ \rightarrow Zh \rightarrow \nu\nu + (bb, jj, gg, W_q W_q^*)$$

FW Moments of Energy Distribution

$$H_{EE;0} = \frac{(\sum_i E_i)^2}{s}$$



$$H_{EE;1} = \frac{|\sum_i \vec{p}_i|^2}{s}$$



- Analogue to CMB power spectrum
- Difference: suppressed sample (“cosmic”) variance, due to large size of data sample
- Similarity: physics at characteristic scales may result in “acoustic peaks”

Benchmark Performance

Jet

<

Jet + FW

~

Image







<

Jet + FW + track

~

Image + track

The performance gap between Jet+FW and Image may be explained by higher order correlation terms.



Application: Measurement of $\Gamma(h)$ @ 240 GeV, 5 ab⁻¹

The most important method for the Higgs factory mode:
Limitation mostly arise from BR(h→WW*) and $\sigma(\nu\nu h)$ rate measurements


$$\Gamma_h^* = \frac{\Gamma(h \rightarrow WW^*)}{\text{BR}(h \rightarrow WW^*)} \propto \frac{\sigma(\nu\nu h)}{\text{BR}(h \rightarrow WW^*)} = \frac{[\sigma(\nu\nu h_b)][\sigma(Zh)]^2}{[\sigma(Zh_b)][\sigma(Zh_W)]}$$

Γ_h (%)	CEPC ₂₄₀₍₂₅₀₎ [14, 65]	FCC ₂₄₀ [15]	FCC ₂₄₀₊₃₆₅ [15]	CLIC ₃₅₀ [66]	ILC ₂₅₀ [64, 67, 68]
Method A	5.1 (5.0)	4.5*	4.2*	-	20*
Method B	3.5 (3.2)	3.5*	1.7*	6.7	13
Method C	-	-	3.4*	-	-
Combined	2.8 (2.7)	2.7	1.3	6.7	11

*In our study, we also include h→ cc/gg/ττ decays to take the advantage of machine learning (~ 20% increase in net signal rate.)

Event level results @ 240 GeV, 5 ab⁻¹

	Jet	Jet+FW	Jet+FW+track	Image	Image+track
Precision (%)	J1	J2	J3	E1	E2
$\sigma(Z_\nu h_{W_{lq}})$	1.7 (1.6)	1.4 (1.6)	1.5 (1.6)	1.5 (1.4)	1.5 (1.4)
$\sigma(Z_\nu h_{W_{qq}})$	1.6 (1.6)	1.2 (1.2)	1.1 (1.1)	1.1 (1.1)	1.1 (1.1)
$\sigma(\nu\nu h_h)$	2.8 (2.7)	1.8 (1.7)	1.9 (1.8)	1.4 (1.4)	1.3 (1.3)
Γ_h	$3.2^{+0.9}_{-0.3}$ (3.1)	$2.3^{+0.7}_{-0.2}$ (2.2)	$2.3^{+0.7}_{-0.2}$ (2.3)	$1.9^{+0.5}_{-0.1}$ (1.9)	$1.9^{+0.4}_{-0.1}$ (1.9)



**2.3% with jet level inputs +
FW moments**

1.9% with event-level inputs

The precision achieved is robust against the rescaling of detector resolutions and different detector templates

Outlook

Can the Higgs decay width be measured at sub percent level @ 240+365 GeV or even @ 240 GeV, given the currently proposed detector baseline?

- Apply event-level ML to multiple channels
- Extra information: charge, pid, displacement, etc.
- Advanced ML techniques
-

We expect event-level analysis with ML to be broadly applied to other hadronic-event measurements at future e-e+ colliders. To what extent one can benefit from it?

- Higgs couplings to quarks/gluons
- CP properties of Higgs boson
- Flavor physics
-




Precision Frontier of Next Decades


The precision frontier of next decades in Higgs and electroweak physics is expected to be defined by a future e-e⁺ collider.

Measurements	CEPC ₂₅₀ [61]	FCC ₂₄₀ [62]	FCC ₃₆₅ [62]	CILC ₃₅₀ [63]	ILC ₂₅₀ [60, 64, 65]
$\sigma(Zh)$	0.5%	0.5%	0.9%	1.6%	2.6%
$\sigma(Zh)\text{BR}(h \rightarrow bb)$	0.3%	0.3%	0.5%	0.86%	1.2%
$\sigma(Zh)\text{BR}(h \rightarrow cc)$	3.1%	2.2%	3.5%	14%	8.3%
$\sigma(Zh)\text{BR}(h \rightarrow gg)$	1.2%	1.9%	6.5%	6.1%	7.0%
$\sigma(Zh)\text{BR}(h \rightarrow WW^*)$	0.9%	1.2%	2.6%	5.1%	6.4%
$\sigma(Zh)\text{BR}(h \rightarrow ZZ^*)$	4.9%	4.4%	12%	-	19%
$\sigma(h\nu\nu)\text{BR}(h \rightarrow bb)$	2.9%	3.1%	0.9%	1.9%	10.5%
$\sigma(h\nu\nu)\text{BR}(h \rightarrow cc)$	-	-	10%	26%	-
$\sigma(h\nu\nu)\text{BR}(h \rightarrow WW^*)$	-	-	3.0%	-	-
$\sigma(h\nu\nu)\text{BR}(h \rightarrow ZZ^*)$	-	-	10%	-	-


[F. An et al., 1810.09037; A. Abada et al., (2019); H. Abramowicz et al., 1608.07538]





Track Variables Defined at Event-Level









Benchmark Study (4j)



Benchmark Study (4j)




Benchmark Study (4j)






Signal	Backgrounds			
$Z_\nu h_{W_l q}$	$W_l W_q$	$Z_l Z_{q_5}$	$Z_\nu h_\tau$	
8.57×10^3	2.41×10^5	1.04×10^3	3.22×10^3	
$Z_\nu h_{W_{qq}}$	$Z_\nu Z_{q_5}$	$q_5 q_5 (\gamma)$	$\gamma\gamma \rightarrow q_5 q_5$	$W_q W_q / Z_{q_5} Z_{q_5}$
1.65×10^4	5.61×10^4	4.01×10^4	4.41×10^2	1.42×10^4
	$Z_\nu h_b$	$Z_\nu h_c$	$Z_\nu h_g$	$Z_\nu h_{Z_{q_5} q_5}$
	8.78×10^4	4.71×10^3	1.41×10^4	2.10×10^3

Signal	$\nu\nu h_b$	$\nu\nu h_c$	$\nu\nu h_g$	$\nu\nu h_\tau$
1.51×10^4	1.24×10^4	6.43×10^2	1.92×10^3	1.50×10^2
Higgs backgrounds	$Z_\nu h_b$	$Z_\nu h_c$	$Z_\nu h_g$	$Z_\nu h_\tau$
1.39×10^5	9.47×10^4	5.08×10^3	1.52×10^4	1.06×10^3
	$Z_\nu h_{V_{q_5 q_5}}$	$\nu\nu h_{V_{q_5 q_5}}$		
	2.01×10^4	2.51×10^3		
Non-Higgs backgrounds	$q_5 q_5 (\gamma) / \gamma\gamma \rightarrow q_5 q_5$	$W_q W_q$	$Z_{q_5} Z_{q_5}$	$Z_\nu Z_{q_5}$
1.40×10^5	$6.79 \times 10^4 / 2.81 \times 10^3$	1.26×10^4	6.61×10^2	5.61×10^4


Zh->vvWW* (2j1l)




Zh->vvWW* (4j)



Zh->vvWW* (2j1l)





Zh->vvWW* (4j)



Signal	$\nu\nu h_b$	$\nu\nu h_c$	$\nu\nu h_g$	$\nu\nu h_\tau$
1.51×10^4	1.24×10^4	6.43×10^2	1.92×10^3	1.50×10^2
Higgs backgrounds	$Z_\nu h_b$	$Z_\nu h_c$	$Z_\nu h_g$	$Z_\nu h_\tau$
1.39×10^5	9.47×10^4	5.08×10^3	1.52×10^4	1.06×10^3
	$Z_\nu h_{V_{q_5 q_5}}$	$\nu\nu h_{V_{q_5 q_5}}$		
	2.01×10^4	2.51×10^3		
Non-Higgs backgrounds	$q_5 q_5 (\gamma) / \gamma\gamma \rightarrow q_5 q_5$	$W_q W_q$	$Z_{q_5} Z_{q_5}$	$Z_\nu Z_{q_5}$
1.40×10^5	$6.79 \times 10^4 / 2.81 \times 10^3$	1.26×10^4	6.61×10^2	5.61×10^4

vvh (2j)



vvh (2j)

