

Testing SFOEWPT of 2HDM at future Colliders

Wei Su

Uni of Adelaide



THE UNIVERSITY
of ADELAIDE

2020.xxxxx (WS, M. White, A. Williams, M. Zhang)


1910.06269 (WS)

1909.09035 (WS, M. White, A. Williams, Y. Wu)


outline

- 🌸 Precision measurements at future collider
- 🌸 2HDM: Brief Introduction
- 🌸 Individual constraints: theory, EW, Higgs, flavour
- 🌸 Results and Conclusion

Precision Measurements





CMS-HIG-14-042
ATLAS-HIGG-2014-14




LHC Run-1:

$$m_h = 125.09$$



CEPC



LHC
HL-LHC
FCC

Precision: Higgs couplings

CEPC-CDR , FCC-ee, ILC Operating Scenarios

collider	CEPC	FCC-ee			ILC				
\sqrt{s}	240 GeV	240 GeV	365 GeV	250 GeV	350 GeV	500 GeV			
$\int \mathcal{L} dt$	5.6 ab ⁻¹	5 ab ⁻¹	1.5 ab ⁻¹	2 ab ⁻¹	200 fb ⁻¹	4 ab ⁻¹			
production	Zh	Zh	Zh	$\nu\bar{\nu}h$	Zh	Zh	$\nu\bar{\nu}h$	Zh	$\nu\bar{\nu}h$
$\Delta\sigma/\sigma$	0.5%	0.5%	0.9%	—	0.71%	2.0%	—	1.05	—
decay	$\Delta(\sigma \cdot BR)/(\sigma \cdot BR)$								
$h \rightarrow b\bar{b}$	0.27%	0.3%	0.5%	0.9%	0.46%	1.7%	2.0%	0.63%	0.23%
$h \rightarrow c\bar{c}$	3.3%	2.2%	6.5%	10%	2.9%	12.3%	21.2%	4.5%	2.2%
$h \rightarrow gg$	1.3%	1.9%	3.5%	4.5%	2.5%	9.4%	8.6%	3.8%	1.5%
$h \rightarrow WW^*$	1.0%	1.2%	2.6%	3.0%	1.6%	6.3%	6.4%	1.9%	0.85%
$h \rightarrow \tau^+\tau^-$	0.8%	0.9%	1.8%	8.0%	1.1%	4.5%	17.9%	1.5%	2.5%
$h \rightarrow ZZ^*$	5.1%	4.4%	12%	10%	6.4%	28.0%	22.4%	8.8%	3.0%
$h \rightarrow \gamma\gamma$	6.8%	9.0%	18%	22%	12.0%	43.6%	50.3%	12.0%	6.8%
$h \rightarrow \mu^+\mu^-$	17%	19%	40%	—	25.5%	97.3%	178.9%	30.0%	25.0%
$(\nu\bar{\nu})h \rightarrow b\bar{b}$	2.8%	3.1%	—	— ^{Wei Su}	3.7%	—	—	—	—

Precision: EW

	Current ($1.7 \times 10^7 Z's$)			CEPC ($10^{10} Z's$)			FCC-ee ($7 \times 10^{11} Z's$)			ILC ($10^9 Z's$)						
	σ	correlation			σ (10^{-2})	correlation			σ (10^{-2})	correlation			σ (10^{-2})	correlation		
		<i>S</i>	<i>T</i>	<i>U</i>		<i>S</i>	<i>T</i>	<i>U</i>		<i>S</i>	<i>T</i>	<i>U</i>		<i>S</i>	<i>T</i>	<i>U</i>
<i>S</i>	0.04 ± 0.11	1	0.92	-0.68	2.46	1	0.862	-0.373	0.67	1	0.812	0.001	3.53	1	0.988	-0.879
<i>T</i>	0.09 ± 0.14	-	1	-0.87	2.55	-	1	-0.735	0.53	-	1	-0.097	4.89	-	1	-0.909
<i>U</i>	-0.02 ± 0.11	-	-	1	2.08	-	-	1	2.40	-	-	1	3.76	-	-	1

CEPC-CDR , FCC-ee, ILC Operating Scenarios

2HDM: Brief Introduction

- Two Higgs Doublet Model

$$\Phi_i = \begin{pmatrix} \phi_i^+ \\ (v_i + \phi_i^0 + iG_i)/\sqrt{2} \end{pmatrix}$$

$$v_u^2 + v_d^2 = v^2 = (246\text{GeV})^2$$

$$\tan \beta = v_u/v_d$$

	ϕ_1	ϕ_2
Type I	u,d,l	
Type II	u	d,l
lepton-specific	u,d	l
flipped	u,l	d

$$\begin{pmatrix} H^0 \\ h^0 \end{pmatrix} = \begin{pmatrix} \cos \alpha & \sin \alpha \\ -\sin \alpha & \cos \alpha \end{pmatrix} \begin{pmatrix} \phi_1^0 \\ \phi_2^0 \end{pmatrix},$$

$$A = -G_1 \sin \beta + G_2 \cos \beta$$

$$H^\pm = -\phi_1^\pm \sin \beta + \phi_2^\pm \cos \beta$$

- Parameters (CP-conserving, Z_2 Symmetry)

$$m_{11}^2, m_{22}^2, \lambda_1, \lambda_2, \lambda_3, \lambda_4, \lambda_5$$



$$v, \tan \beta, \alpha, m_h, m_H, m_A, m_{H^\pm}$$

Soft Z_2 symmetry breaking: m_{12}^2

Wei Su

246 GeV

125. GeV


Constraints: theory

- Perturbativity
- Stability of the potential
- Unitarity of the scattering matrix

$$\cos(\beta - \alpha) = 0,$$
$$m_\Phi \equiv m_H = m_A = m_{H^\pm}$$



$$\lambda v^2 \equiv m_\Phi^2 - m_{12}^2 / s_\beta c_\beta$$

$$-125^2 \text{GeV}^2 < \lambda v^2 < 600^2 \text{GeV}^2$$



Constraints: EW+Higgs (indirect)


Type-II



[1912.01431](#) ,
N Chen, T Han, S Li, S Su, WS

Constraints: heavy Higgs (direct)


Type-II



[1910.06269](#), WS

Constraints: flavor

Type-II




2020.xxxx , gambit community

Results: Strong First Order Phase Transition

Allowed by LHC

SFOPT + LHC R-2

$m_H < 1 \text{ TeV}$





Results: Strong First Order Phase Transition

SFOPT + LHC R-2

SFOPT + CEPC

$m_H < 1 \text{ TeV}$

Preliminary





Results: Strong First Order Phase Transition

Preliminary

SFOPT + LHC R-2

Further study:
 κ_Z or $\cos(\beta - \alpha)$? SFOPT



Conclusion



🌸 Theory,

🌸 EW,

🌸 Higgs,


🌸 flavour

1. $m_H < 1$ TeV under SFOPT+ LHC Run-2
2. Type-II not allowed under SFOPT+ CEPC



Thanks!

SM-like Higgs



CMS-HIG-14-042
ATLAS-HIGG-2014-14

LHC Run-I:

$$m_h = 125.09 \pm 0.24 \text{ GeV}$$

LHC Run-II

$$\Delta = 160 \text{ MeV}$$

1. Higgs property
2. Higgs potential: EWPT

