


XVIII INTERNATIONAL WORKSHOP ON DEEP-INELASTIC SCATTERING AND RELATED SUBJECTS

Combined Measurement and QCD Analysis of the Inclusive ep Scattering Cross Sections at HERA

[Published in JHEP 1001:109,2010]

SHIRAZ HABIB

*on behalf of the **H1** and **ZEUS** Collaborations*



O
T
O
N
D
I
S
C
U
S
I
O
N
S
2
0
1
0

Combined Measurement and QCD Analysis of the Inclusive ep Scattering Cross Sections at HERA

[Published in JHEP 1001:109,2010]

SHIRAZ HABIB

*on behalf of the **H1** and **ZEUS** Collaborations*

Outline:

- HERA Collider and the H1 & ZEUS Detectors
- Combining the H1 & ZEUS Measurements
- HERAPDF1.0 QCD Fit – PDF Determination
- Summary


HERA Collider and the H1 & ZEUS Detectors

HERA : A 6.3 km $e^\pm p$ collider located in **Hamburg, Germany**.


2 Phases of HERA Running [I + II]



HERA I [1992 – 2000]



HERA II [2003 – 2007]




Analyses presented in this talk use HERA I data only






Colliding-Beam Experiments : **H1 & ZEUS**

Inclusive Processes:

$$\text{NC : } e p \rightarrow e' X$$




$$\text{CC : } e p \rightarrow \nu_e X$$



Colliding-Beam Experiments : **H1 & ZEUS**

Inclusive Measurement:


$$\sigma_{\text{NC}}(x, Q^2), \sigma_{\text{CC}}(x, Q^2)$$



$$x = Q^2 / (2p \cdot q)$$

**HERA provides unique opportunity
to study the proton.**

6 orders of magnitude in Q^2



6 orders of magnitude in x



Combining the H1 & ZEUS Measurements

Statistical Precision:

H1 & ZEUS collected *similar* amounts of data:

- 100 pb^{-1} of e^+p data
- 15 pb^{-1} of e^-p data

[HERA I]

A combined measurement should improve **statistical precision** : $\delta_{\text{stat}} \rightarrow 0.707 \delta_{\text{stat}}$

Systematic Precision:

H1 and ZEUS are *different* detectors and use *different* analysis techniques:

- Calorimetry
- Kinematic Reconstruction

The H1 and ZEUS cross sections have different sensitivities to similar sources of correlated systematic uncertainty \Rightarrow improve the **systematic precision**.

The combination method used to average our cross sections takes the uncorrelated errors as well as the systematic correlations into account.



COMBINING THE H1 & ZEUS MEASUREMENTS

Input: H1 & ZEUS published cross sections [Inclusive NC , CC $e^\pm p$].

Combination Method:

[1] ***Swim H1 and ZEUS measurements to common grid (x, Q^2) :***

$$\sigma_{\text{H1}} (x_{\text{H1}}, Q^2_{\text{H1}}) \rightarrow \sigma_{\text{H1}} (x, Q^2) \quad ; \quad \sigma_{\text{ZEUS}} (x_{\text{ZEUS}}, Q^2_{\text{ZEUS}}) \rightarrow \sigma_{\text{ZEUS}} (x, Q^2)$$



COMBINING THE H1 & ZEUS MEASUREMENTS

Input: H1 & ZEUS published cross sections [Inclusive NC , CC $e^\pm p$].

Combination Method:

[1] Swap H1 and ZEUS measurements to common grid (x, Q^2) :

$$\sigma_{\text{H1}} (x_{\text{H1}}, Q^2_{\text{H1}}) \rightarrow \sigma_{\text{H1}} (x, Q^2) \quad ; \quad \sigma_{\text{ZEUS}} (x_{\text{ZEUS}}, Q^2_{\text{ZEUS}}) \rightarrow \sigma_{\text{ZEUS}} (x, Q^2)$$

[2] For CC and NC [$y < 0.35$] : $\sigma_{820} \rightarrow \sigma_{920}$



COMBINING THE H1 & ZEUS MEASUREMENTS

Input: H1 & ZEUS published cross sections [Inclusive NC , CC $e^\pm p$].

Combination Method:

[1] Swap H1 and ZEUS measurements to common grid (x, Q^2) :

$$\sigma_{\text{H1}} (x_{\text{H1}}, Q^2_{\text{H1}}) \rightarrow \sigma_{\text{H1}} (x, Q^2) \quad ; \quad \sigma_{\text{ZEUS}} (x_{\text{ZEUS}}, Q^2_{\text{ZEUS}}) \rightarrow \sigma_{\text{ZEUS}} (x, Q^2)$$

[2] For CC and NC [$y < 0.35$] : $\sigma_{820} \rightarrow \sigma_{920}$

[3] Build a χ^2 function for each data-set, exp:

$$\chi^2_{\text{exp}} (\mathbf{m}, \mathbf{b}) = \sum_i \frac{\left[m^i - \sum_j \gamma_j^i m^i b_j - \mu^i \right]^2}{\delta_{i,\text{stat}}^2 \mu^i \left(m^i - \sum_j \gamma_j^i m^i b_j \right) + \left(\delta_{i,\text{uncor}} m^i \right)^2} + \sum_j b_j^2$$



Input: H1 & ZEUS published cross sections [Inclusive NC , CC $e^\pm p$].

Combination Method:

[1] Swap H1 and ZEUS measurements to common grid (x, Q^2) :

$$\sigma_{\text{H1}} (x_{\text{H1}}, Q^2_{\text{H1}}) \rightarrow \sigma_{\text{H1}} (x, Q^2) \quad ; \quad \sigma_{\text{ZEUS}} (x_{\text{ZEUS}}, Q^2_{\text{ZEUS}}) \rightarrow \sigma_{\text{ZEUS}} (x, Q^2)$$

[2] For CC and NC [$y < 0.35$] : $\sigma_{820} \rightarrow \sigma_{920}$

[3] Build a χ^2 function for each data-set, exp:

Combination at point i

[Estimate of 1 true cross section]

$$\chi^2_{\text{exp}} (\mathbf{m}, \mathbf{b}) = \sum_i \frac{[m^i - \sum_j \gamma_j^i m^i b_j - \mu^i]^2}{\delta_{i,\text{stat}}^2 \mu^i (m^i - \sum_j \gamma_j^i m^i b_j) + (\delta_{i,\text{uncor}} m^i)^2} + \sum_j b_j^2$$

Measurement at point i

Shift of the j^{th} source of correlated uncertainty

Sensitivity of the cross section to the j^{th} source of correlated uncertainty.

γ_j^i defined as the relative change of the measurement for a 1 sigma shift of the error source

$$\delta_{i,\text{stat}} / \delta_{i,\text{uncor}}$$

Relative stat. / syst. error on the measurement



[4] Build a total χ^2 for all data sets: $\chi_{\text{tot}}^2 = \sum_{\text{exp}} \chi_{\text{exp}}^2$

[5] Minimize χ_{tot}^2

Notes:



[1] Additive instead of multiplicative error treatment added as an extra “procedural uncertainty”. Typically less than 0.5%.

$$\chi_{\text{exp}}^2(\mathbf{m}, \mathbf{b}) = \sum_i \frac{\left[m^i - \sum_j \gamma_j^i m^i b_j - \mu^i \right]^2}{\delta_{i,\text{stat}}^2 \mu^i \left(m^i - \sum_j \gamma_j^i m^i b_j \right) + \left(\delta_{i,\text{uncor}} m^i \right)^2} + \sum_j b_j^2$$

[2] Only correlation assumed between H1 and ZEUS is due to normalization. Other correlations between the experiments which contribute significantly are due to background estimation [significant only at high y] and hadron energy scale. Added as procedural uncertainties. Can be a few %.



- 1402 measurements with 110 correlated sources of uncertainty combined to 741 cross sections.
- $\chi^2 / \text{dof} = 636.5 / 656$; No tension in Pulls ; $|b_j| < 2 \Rightarrow \text{H1 and ZEUS Agree!}$



Systematic Uncertainty:

- $\delta_{\text{H1 LAR}} \rightarrow 0.45 \delta_{\text{H1 LAR}}$
- $\delta_{\text{ZEUS BG}} \rightarrow 0.35 \delta_{\text{ZEUS BG}}$

Overall Precision:

- 2% for $3 < Q^2 < 500 \text{ GeV}^2$
- 1% for $2 < Q^2 < 100 \text{ GeV}^2$



HERAPDF1.0 QCD Fit – PDF Determination

Data Input:

The HERA Inclusive Combined Cross Sections [NC, CC, $e^\pm p$] is a **Consistent Data Set** which allows the extraction of valence, sea quark and gluon (scaling violation).

Model:

PDF evolution	:	$Q_0^2 = 1.9 \text{ GeV}^2$ use DGLAP @ NLO
Renormalization & Factorization scale	:	Q^2
m_c	:	1.4 GeV
m_b	:	4.75 GeV
$\alpha_s(M_z)$:	0.1176
Q^2_{\min} of Data	:	3.5 GeV 2
$f_s = \bar{s} / (\bar{s} + \bar{d})$ @ Q_0^2	:	0.31
Heavy Quark Coefficient Functions	:	GMVFNS Robert Thorne VFNS 2008



Parameterization $xf(x)$:

- Use General form $xf(x) = A x^B (1-x)^C$: $xg(x); xu_v(x); xd_v(x); x\bar{U}(x); x\bar{D}(x)$ and fit $\Rightarrow \chi^2_9$.
- Modify $xf(x)$: $xf(x) \rightarrow xf(x) (1 + \epsilon x^{1/2} + Dx + Ex^2)$ and fit $\Rightarrow \chi^2_{10}$. Find ϵ, D, E that gives best 10 parameter fit $\Rightarrow Eu_v$
- Repeat to find best 11 parameter fit. Find that χ^2 is saturated. Settle for **10 parameter fit**:

$$xg(x) = A_g x^{B_g} (1 - x)^{C_g},$$

$$xu_v(x) = A_{u_v} x^{B_{u_v}} (1 - x)^{C_{u_v}} (1 + E_{u_v} x^2),$$

$$xd_v(x) = A_{d_v} x^{B_{d_v}} (1 - x)^{C_{d_v}},$$

$$x\bar{U}(x) = A_{\bar{U}} x^{B_{\bar{U}}} (1 - x)^{C_{\bar{U}}},$$

$$x\bar{D}(x) = A_{\bar{D}} x^{B_{\bar{D}}} (1 - x)^{C_{\bar{D}}}.$$

Additional Constraints:

- Quark Number Sum Rules
- Momentum Sum Rule
- $B_{\bar{U}} = B_{\bar{D}}$ & $A_{\bar{U}} = A_{\bar{D}} (1-f_s)$
 $\bar{u} \rightarrow \bar{d}$ as $x \rightarrow 0$
- $B_{u_v} = B_{d_v}$



PDF Uncertainties : Experimental \oplus Model \oplus Parameterization

Experimental Uncertainties:

- Consistent NC, CC, $e^\pm p$ Data Sets \Rightarrow Use conventional tolerance : $\Delta\chi^2 = 1$
- 110 corr. syst. Uncertainties \Rightarrow Added in quadrature
- Procedural Uncertainties \Rightarrow Use offset method

Model Uncertainties : f_s , m_c , m_b , Q^2_{min}

Variation	Standard Value	Lower Limit	Upper Limit
f_s	0.31	0.23	0.38
m_c [GeV]	1.4	1.35	1.65
m_b [GeV]	4.75	4.3	5.0
Q^2_{min} [GeV 2]	3.5	2.5	5.0
Q_0^2 [GeV 2]	1.9	1.5	2.5

Parameterization Uncertainties :


- Vary Q_0^2 and allow for negative gluon at its lower limit
- Relax $B_{u_\nu} = B_{d_\nu}$ constraint
- Variations of 11 parameter fits

} Uncertainty = Envelope
of all Variations




Good $\chi^2 / \text{dof} = 574 / 582$

H1 and ZEUS



H1 and ZEUS




HERA and Fixed Target Data well described by HERAPDF1.0



Starting Scale $Q^2 = 1.9 \text{ GeV}^2$

H1 and ZEUS





For the sea and gluon experimental uncertainties are relatively small.



Evolve to $Q^2 = 10000 \text{ GeV}^2$

H1 and ZEUS





Few % uncertainty



W and Z Rapidity Distributions at LHC [7 TeV]

[Amanda Cooper-Sarkar]



- HERAPDF1.0 7TeV
- exp. uncert.
- model uncert.
- param uncert.

Uncertainty on Prediction 5% and better.



Summary & The Shape of Things to Come ...

- H1 and ZEUS have combined their HERA I data resulting in improved precision [1% in best measured region : NC $20 < Q^2 < 100 \text{ GeV}^2$]
- An NLO QCD Fit to the combined measurement gives the **HERAPDF1.0** PDFs with precision at the level of few % in low x region.
- New Data [Here at DIS 2010]:
 - HERA II 460/575 GeV NC (F_L) [*J. Grebenyuk, V. Radescu*]
 - HERA I/II F_2 charm [*R. Plačakyte / A. Cooper-Sarkar*]
- New Data:
 - HERA II NC, CC Inclusive
 - Jet Data

More to come ...


Thanks!




Backup



Pulls




Pulls on Sys




CC e+ Combination

H1 and ZEUS



ACOT

H1 and ZEUS



Alpha_s

H1 and ZEUS

