

Measurement of the J/ψ photoproduction at large momentum transfer at HERA

Alessia Bruni - INFN Bologna

on behalf of the ZEUS Collaboration

Deep Inelastic Scattering 2010

Florence, April 20, 2010

Measurement of the J/ψ photoproduction at large momentum transfer

Recent paper by ZEUS Collaboration

ZEUS Collab., DESY-09-137, accepted by JHEP

Results preliminary since a while, presented at DIS05 and ICHEP06:

<http://www-zeus.desy.de/physics/diff/pub/prelim/05-006/dszuba-dis05.pdf>

<http://www-zeus.desy.de/physics/phch/conf/ichep06/diff/6/ZEUS-prel-05-006.ps>

Similar results published by H1 Collaboration:


Diffractive Photoproduction of J/ψ Mesons with Large Momentum Transfer at HERA

A. Aktas et al., Phys Lett B568 (2003) 205-218

see also the talk of Boris Blok at this workshop:

DGLAP versus perturbative Pomeron in hard diffractive processes large momentum transfer at HERA and LHC

$\gamma p \rightarrow J/\psi Y$ at large $|t|$ - kinematics



Kinematic variables


- Q^2 , virtuality of the exchanged photon
- W , γp centre-of-mass energy
- t , 4-momentum transfer squared at the p-vertex
- M_Y mass of the p-dissociative state

- Usually the inelasticity variables z is introduced, fraction of the γ energy transferred to the J/ψ :

$$z = \frac{p_{J/\psi} \cdot p}{q \cdot p} \simeq 1 - \frac{M_Y^2 + |t|}{W^2} \simeq 1 \text{ for semi-exclusive processes}$$

At HERA (ep 27.5-820/920 GeV²), a wide range in W , t and M_Y can be accessed


$\gamma p \rightarrow J/\psi Y, J/\psi \rightarrow \mu^+ \mu^-$ experimental signature



- two tracks well reconstructed in the central tracking detector
- each track associated with a muon candidate in CAL and to a hit in the muon detector (trigger)
- some energy in the forward region, nothing else
- scattered electron not observed

J/ψ at large $|t|$, extraction of the signal


- HERA I data, 1996-2000, 112 pb^{-1} of luminosity
- $30 < W < 160 \text{ GeV}$
- $2 < |t| < 20 \text{ GeV}^2$
- $z > 0.95$, inelasticity cut
- 2817 events in $2.6 < M_{\mu\mu} < 3.5 \text{ GeV}$



Background from exclusive channel $ep \rightarrow e\mu^+\mu^-p$ suppressed
Background from $ep \rightarrow e\mu^+\mu^-Y$ subtracted ($\simeq 6 - 10\%$)

J/ψ at large $|t|$, control plots

ZEUS



Kinematic region


- $30 < W < 160$ GeV
- $|t| > 2$ GeV²
- $z > 0.95$ ($M_Y < 30$ GeV)

MC EPSOFT $\gamma p \rightarrow J/\psi Y$
MC GRAPE $\gamma p \rightarrow \mu^+ \mu^- Y$

Kinematic variables well described by the Monte Carlo

Measurement of the J/ψ photoproduction at large momentum transfer

$$\gamma p \rightarrow J/\psi Y$$



Hard inelastic diffractive process

- For hard scales pQCD can be applied
- Two scales: $M_{J/\psi}$ and t
- In perturbative QCD the VM production is mediated by an exchange of a gluon ladder in a colour singlet state


- $x = \frac{|t|}{M_Y^2 - M_p^2 + |t|} \simeq \frac{|t|}{(1-z)W^2}$

- large $|t|$ domain allows to investigate QCD dynamics vs x

Challenge is to describe all the variables simultaneously


QCD dynamics, DGLAP vs BFKL

QCD dynamics: $\alpha_S \ln \frac{x_0}{x} \ln \frac{Q^2}{Q_0^2}$



- DGLAP dynamics driven by terms $\ln \frac{M_{J/\psi}^2}{Q_0^2 - t}$
- BFKL evolution is driven by terms $\alpha_S^n \ln^n(W^2/|t|) \Rightarrow$ expected to dominate at larger $|t|$
- GLMN (Gotsman, Levin, Maor, Naftali 2002), DGLAP evolution tamed for $|t| \simeq M_{J/\psi}^2$
- FSZ (Frankfurt, Strikman, Zhalov 2008), increase of σ with t due to increase of $xG(x, Q^2)$ on Q^2
- EMP (Enberg, Motyka, Poludniowski 2003), BFKL evolution

J/ψ at large $|t|$, $|t|$ dependence





Fit $d\sigma/dt \simeq |t|^n$

- Data cannot be described by a single exponential fit $d\sigma/dt \simeq e^{bt}$ neither by a single power $d\sigma/dt \simeq |t|^n$
- All models predicts a power-law t -dependence, with n depending on t range

J/ψ at large $|t|$, $|t|$ dependence - model comparison

ZEUS




EMP (BFKL) below data

GLMN (DGLAP) validity range of $|t| < M_{J/\psi}^2$

FSZ describe data up to 12 GeV²

Models able to describe data but not for the full $t(x)$ range



J/ψ at large $|t|$, W dependence - model comparison



- Data rise with W for all t region
- BFKL (EMP) predictions too steep
- DGLAP (GLMN) approach fails to describe σ rise at low x
- FKS: increase of σ due to gluon distribution in the proton

$$x = \frac{|t|}{M_Y^2 - M_p^2 + |t|} \simeq \frac{|t|}{(1-z)W^2}$$



J/ψ at large $|t|$, W dependence - model comparison



ZEUS results are in good agreement with those of H1 in the common kinematic region


J/ψ at large $|t|$, W dependence as a function of t

Effective Pomeron trajectory $d\sigma/dt = F(t) \cdot W^{4(\alpha_P(t)-1)}$



- Pomeron intercept consistent with soft $\alpha_P(0) = 1.0808$
- Pomeron slope consistent with BFKL Pomeron
- H1 $\alpha_P(0) = 1.167 \pm 0.048(\text{stat}) \pm 0.024(\text{syst})$
- H1 $\alpha'_P = -0.0135 \pm 0.0074(\text{stat}) \pm 0.0051(\text{syst})$

Decay angular distribution - helicity frame



θ_h, ϕ_h angles of decay muons in the meson rest frame

Φ angle between scattering and production plane

Angular distribution are related to the spin of the γ^* and the meson


Angular distr. \rightarrow spin density matrix elements r_{ij}^{kl} , \rightarrow helicity amplitudes $T_{\lambda_{VM}\lambda_\gamma}$

s-channel helicity conservation: the outgoing VM retains the γ helicity

pQCD: during the interaction, the orbital angular momentum of $q\bar{q}$ can be modified due to the transfer of momentum of the gluons;
 \Rightarrow the helicity of the outgoing VM differs from the one of the γ , helicity flip between photon and meson is possible



Decay angular distributions in $|t|$ bins

ZEUS



- θ_h, ϕ_h angles of decay muons in the meson rest frame estimated in different t bins
- Spin density matrix elements are extracted from fit to the angular distributions

Helicity spin density matrix elements as a function of $|t|$



r_{1-1}^{04} is related to interference between non-flip and double-flip amplitude

r_{00}^{04} represents the probability that $J\psi$ has 0 helicity

$\text{Re}(r_{10}^{04})$ is proportional to the single flip amplitude

These spin density elements expected to be 0 in SCHC

Measurement of the J/ψ photoproduction at large momentum transfer

Summary

- $\gamma p \rightarrow J/\psi Y$ studied by ZEUS for t up to 20 GeV^2
- DGLAP models are able to describe data, but up to $|t| < 5 \text{ GeV}^2$
- BFKL models not yet able to describe data over all the kinematic range
- in general, no model gives a good description of all variables

NB: The discussion at the workshop has been added in next slide

Discussion at DIS2010

Comments raised by Boris Blok

- the model of Frankfurt, Strikman and Zhalov should be replaced by the model described in the recent papers *DGLAP versus perturbative Pomeron in large momentum transfer hard diffractive processes at HERA and LHC*

B. Blok, L. Frankfurt, M. Strikman, arXiv:1002.3048

and *The energy dependence of the hard exclusive diffractive processes in pQCD as the function of momentum transfer*

B. Blok, L. Frankfurt, M. Strikman, arXiv:1001.2469

This DGLAP model is able to describe data on the entire t range

- The α_P measured by ZEUS is an effective value and cannot be compared to the trajectory of the *soft* Pomeron or to the BFKL Pomeron