



THE OHIO STATE UNIVERSITY



**Full Run 2 Results
for Disappearing Tracks at CMS**


LLP7

**Brian Francis
for the CMS Collaboration**

**Accepted by PLB: [arXiv:2004.05153](https://arxiv.org/abs/2004.05153) ([CMS public page](#))
Featured: cms.cern/news/disappearing-act-cms**

A Striking Signature

- A long-lived charged particle decaying within the CMS tracker
- If the decay products are undetected, the track “disappears”
 - Neutral, weakly interacting
 - Too low momentum to be reconstructed
- Observation would be a striking sign of BSM physics
 - **Arises in many models**
 - Multiple handles to study — decay length, mass, dE/dx , potential recovery of decay products




Canonical Benchmark — AMSB


Simulation

- Consider anomaly-mediated supersymmetry breaking (AMSB)
- Small mass splitting between chargino ($\tilde{\chi}^\pm$) and neutralino ($\tilde{\chi}^0$)
 - Direct electroweak production:
$$pp \rightarrow \tilde{\chi}^\pm \tilde{\chi}^\mp, \tilde{\chi}^\pm \tilde{\chi}^0$$
- $\tilde{\chi}^\pm \rightarrow \pi^\pm \chi^0$ with lifetime $O(1)$ ns
- $\tilde{\chi}^0$ interacts only weakly, and π^\pm is too soft to be reconstructed

AMSB is one benchmark of many that would frequently produce disappearing tracks






Rare Backgrounds




- SM particles don't disappear, however there are rare backgrounds:
 - Charged hadrons can interact with detector material, e.g.
$$\pi^+ + n \rightarrow \pi^0 + p$$
 - Leptons can be mis-reconstructed
 - Spurious (“fake”) tracks — pattern recognition errors
- Approach to controlling these:
 - Explore and mitigate all possible ways to lose tracker hits
 - Estimate the remaining probability to fall into search region

Phase 1 CMS Pixel Upgrade



- CMS previously searched for disappearing tracks (@ 8 and 13 TeV) using a 3-layer pixel tracker
 - Limited sensitivity to short tracks (compared to ATLAS' JHEP 1806 (2018) 022)
- Since 2017 CMS now has a 4-layer pixel tracker
- New analysis bins results in the number of layers with measurements:
 - $n_{\text{layers}} = 4, 5, \geq 6$
 - **New sensitivity to shorter particle lifetimes**

Track Selection



- $p_t > 55$ GeV, isolated from other tracks/jets
- Require high track quality:
 - ≥ 4 pixel hits
 - No missing inner/middle hits
- Veto all tracks identified as leptons ($e/\mu/\tau_h$)
- Reject tracks in regions of lower lepton reconstruction efficiency
- “Disappearing” is defined as:
 - ≥ 3 missing outer hits — rejects most SM tracks
 - < 10 GeV energy deposited within $\Delta R < 0.5$
 - Rejects most electrons and charged hadrons
 - E.g. electrons with significant brem. energy causing a track reconstruction failure

Backgrounds


- Possible for leptons to enter search region
 - Dead/noisy channels, track fitting errors, nuclear interactions with the tracker material, etc
 - Estimate probability for several conditions:

$$N_{\text{est}}^{\ell} = \frac{N_{\text{ctrl}}^{\ell}}{\epsilon_{\text{trigger}}^{\ell}} P_{\text{veto}} P_{\text{off}} P_{\text{trig}} P_{\text{HEM}}$$

- “Spurious” tracks: not real charged trajectories, but pattern recognition errors
 - Tend to have missing hits and little associated calorimeter energy
 - Estimate probability of such tracks in $Z(\mu\mu)$ (plus track) events and extrapolate to search region

Lepton $P(\text{veto})$

- Measured using a tag-and-probe method for each flavor
- Calculate the probability for probes to disappear
- Also select a same-sign sample to subtract non-Z background events from T&P sample



$$P_{\text{veto}} = \frac{N_{\text{T\&P}}^{\text{veto}} - N_{\text{SS T\&P}}^{\text{veto}}}{N_{\text{T\&P}} - N_{\text{SS T\&P}}}$$



| Data-taking period | n_{lay} | P_{veto} | | |
|--------------------|------------------|--------------------------------------|---------------------------------------|--------------------------------------|
| | | Electrons | Muons | τ_h |
| 2017 | 4 | $(8.2 \pm 5.2) \times 10^{-4}$ | $(0.0_{-0.0}^{+3.9}) \times 10^{-3}$ | $(6.9_{-5.1}^{+8.3}) \times 10^{-2}$ |
| | 5 | $(2.2 \pm 0.9) \times 10^{-4}$ | $(3.2 \pm 1.3) \times 10^{-2}$ | $(6.5_{-2.7}^{+2.9}) \times 10^{-2}$ |
| | ≥ 6 | $(2.7 \pm 0.5) \times 10^{-5}$ | $(1.2 \pm 0.5) \times 10^{-6}$ | $(1.0 \pm 0.4) \times 10^{-3}$ |
| 2018 A | 4 | $(1.3 \pm 0.7) \times 10^{-3}$ | $(1.0 \pm 1.0) \times 10^{-1}$ | $(7.1_{-3.8}^{+5.5}) \times 10^{-2}$ |
| | 5 | $(0.9_{-0.9}^{+1.5}) \times 10^{-4}$ | $(7.4 \pm 4.2) \times 10^{-2}$ | $(4.4_{-4.4}^{+5.5}) \times 10^{-2}$ |
| | ≥ 6 | $(1.6 \pm 0.6) \times 10^{-5}$ | $(1.9 \pm 0.8) \times 10^{-6}$ | $(0.0_{-0.0}^{+7.3}) \times 10^{-4}$ |
| 2018 B | 4 | $(0.0_{-0.0}^{+1.1}) \times 10^{-4}$ | $(4.0_{-4.0}^{+15.0}) \times 10^{-2}$ | $(5.6_{-5.0}^{+6.5}) \times 10^{-2}$ |
| | 5 | $(1.4 \pm 1.1) \times 10^{-4}$ | $(5.8 \pm 3.8) \times 10^{-2}$ | $(5.1_{-3.7}^{+4.5}) \times 10^{-2}$ |
| | ≥ 6 | $(3.3 \pm 0.7) \times 10^{-5}$ | $(1.5 \pm 0.6) \times 10^{-6}$ | $(2.3 \pm 1.0) \times 10^{-3}$ |

Search Results



- Searched 101/fb 13 TeV data recorded in 2017-8

| Data-taking period | n_{lay} | Expected backgrounds | | | Observation |
|--------------------|------------------|-----------------------------|---------------------------------|------------------------------|-------------|
| | | Leptons | Spurious tracks | Total | |
| 2017 | 4 | $1.4 \pm 0.9 \pm 0.2$ | $10.9 \pm 0.7 \pm 4.7$ | $12.2 \pm 1.1 \pm 4.7$ | 17 |
| | 5 | $1.1 \pm 0.4 \pm 0.1$ | $1.0 \pm 0.2 \pm 0.6$ | $2.1 \pm 0.4 \pm 0.6$ | 4 |
| | ≥ 6 | $6.7 \pm 1.1 \pm 0.7$ | $0.04 \pm 0.04^{+0.08}_{-0.04}$ | $6.7 \pm 1.1 \pm 0.7$ | 6 |
| 2018 A | 4 | $1.1^{+1.0}_{-0.6} \pm 0.1$ | $6.2 \pm 0.5 \pm 3.5$ | $7.3^{+1.1}_{-0.8} \pm 3.5$ | 5 |
| | 5 | $0.2^{+0.6}_{-0.2} \pm 0.0$ | $0.5 \pm 0.1 \pm 0.3$ | $0.6^{+0.6}_{-0.2} \pm 0.3$ | 0 |
| | ≥ 6 | $1.8^{+0.6}_{-0.5} \pm 0.2$ | $0.04 \pm 0.04^{+0.06}_{-0.04}$ | $1.8^{+0.6}_{-0.5} \pm 0.2$ | 2 |
| 2018 B | 4 | $0.0^{+0.8}_{-0.0} \pm 0.0$ | $10.3 \pm 0.6 \pm 5.4$ | $10.3^{+1.0}_{-0.6} \pm 5.4$ | 11 |
| | 5 | $0.4^{+0.7}_{-0.3} \pm 0.1$ | $0.6 \pm 0.2 \pm 0.3$ | $1.0^{+0.7}_{-0.3} \pm 0.3$ | 2 |
| | ≥ 6 | $5.7^{+1.2}_{-1.1} \pm 0.6$ | $0.00^{+0.04}_{-0.00} \pm 0.00$ | $5.7^{+1.2}_{-1.1} \pm 0.6$ | 1 |

- Observation of 48 events is consistent with a total background estimate of $47.8^{+2.7}_{-2.3}$ (stat) ± 8.1 (syst) events
- Results are combined with those from 2015-6 to provide upper limits for the full 140/fb of the Run 2 data set




Exclusions



- In the context of AMSB, these results exclude charginos below:
 - Wino-like neutralino case — 884 (474) GeV for a lifetime of 3 (0.2) ns
 - Higgsino-like neutralino case — 750 (175) GeV for a lifetime of 3 (0.05) ns
- **New interpretation for 2017-8 data**

Summary



- CMS has updated its disappearing track search
 - Full Run 2 data set
 - Making use of the Phase 1 pixel upgrade to extend sensitivity to lower particle lifetimes
 - Adding a new higgsino-like neutralino interpretation
- No excess observed over estimated backgrounds
 - Currently obtains the world-best limits at 95% CL over a wide range of particle masses and lifetimes

Accepted by PLB: [arXiv:2004.05153](https://arxiv.org/abs/2004.05153) ([CMS public page](#))

Featured: cms.cern/news/disappearing-act-cms

Backup

Background Systematics

- Spurious tracks:
 - Test assumption that track occurrence probability is independent of physics content by comparing $Z \rightarrow \mu\mu$ to $Z \rightarrow ee$ estimates
 - Uncertainties in transfer factor (ζ) fit parameters
- Leptons:
 - Test assumption that unreconstructed leptons deposit no calorimeter energy by allowing them to instead deposit 10 GeV
 - Limited statistics in NLayers=4,5 muons/taus — compare to electrons, which have sufficient events

**Averages across
data-taking period,
NLayers categories**


| Background | Source | Uncertainty | | |
|-----------------|----------------------------|----------------------------|----------------------|-------------------------|
| | | $n_{\text{lay}} = 4$ | $n_{\text{lay}} = 5$ | $n_{\text{lay}} \geq 6$ |
| Spurious tracks | Control sample | $\pm 19\%$ | $\pm 29\%$ | $\pm 116\%$ |
| | ζ | $\pm 47\%$ | $\pm 47\%$ | $\pm 47\%$ |
| Electrons | Visible calorimeter energy | $\pm 14\%$ | $\pm 14\%$ | $\pm 13\%$ |
| Muons | P_{off} | +7% | +7% | — |
| | P_{trig} | +8% | +2% | — |
| | τ_h | Visible calorimeter energy | $\pm 19\%$ | $\pm 19\%$ |
| | P_{off} | +7% | +7% | — |
| | P_{trig} | +8% | +2% | — |

Signal Systematics


| Source | Uncertainty | | |
|--------------------------------------|----------------------|----------------------|-------------------------|
| | $n_{\text{lay}} = 4$ | $n_{\text{lay}} = 5$ | $n_{\text{lay}} \geq 6$ |
| Pileup | 3.0% | 3.3% | 2.8% |
| ISR | 13% | 13% | 13% |
| Trigger efficiency | 1.1% | 0.8% | 0.4% |
| Jet energy scale | 0.6% | 0.7% | 1.6% |
| Jet energy resolution | 0.5% | 0.5% | 1.3% |
| $p_{\text{T}}^{\text{miss}}$ | 0.3% | 0.3% | 0.4% |
| $E_{\text{calo}}^{\Delta R < 0.5}$ | 0.7% | 0.7% | 0.7% |
| Missing inner hits | 2.3% | 1.0% | 0.3% |
| Missing middle hits | 3.9% | 5.1% | 4.4% |
| Missing outer hits | — | — | 0.2% |
| Reconstructed lepton veto efficiency | 0.1% | 0.1% | — |
| Track reconstruction efficiency | 2.3% | 2.3% | 2.3% |
| Total | 14% | 15% | 14% |

**Averages across
data-taking conditions**

Previous Results in 2015-6 Data



- Previously published results from proton collision data from 2015-6
 - [JHEP 06 \(2018\) 022](https://arxiv.org/abs/1806.02222)



| Run period | Estimated number of background events | | | Observed events |
|------------|---------------------------------------|-----------------------|-----------------------|-----------------|
| | Leptons | Spurious tracks | Total | |
| 2015 | 0.1 ± 0.1 | $0_{-0}^{+0.1}$ | 0.1 ± 0.1 | 1 |
| 2016A | $2.0 \pm 0.4 \pm 0.1$ | $0.4 \pm 0.2 \pm 0.4$ | $2.4 \pm 0.5 \pm 0.4$ | 2 |
| 2016B | $3.1 \pm 0.6 \pm 0.2$ | $0.9 \pm 0.4 \pm 0.9$ | $4.0 \pm 0.7 \pm 0.9$ | 4 |
| Total | $5.2 \pm 0.8 \pm 0.3$ | $1.3 \pm 0.4 \pm 1.0$ | $6.5 \pm 0.9 \pm 1.0$ | 7 |