The 8th Asian Triangle Heavy-Ion Conference (ATHIC2021)



Contribution ID: 94

Type: not specified

Measurement of global polarization of Λ hyperons in Au+Au $\sqrt{sNN} = 7.2$ GeV fixed-target collisions at RHIC-STAR experiment

Monday 8 November 2021 10:51 (17 minutes)

Non-central heavy-ion collisions produce a large angular momentum that leads to vorticity of the created system.

Due to the spin-orbit coupling, spin directions of particles are aligned with the orbital angular momentum of the system.

Global polarization of Λ and $\overline{\Lambda}$ hyperons has been measured in Au+Au collisions from $\sqrt{s_{NN}}$ = 7.7 GeV to 200 GeV.[1,2]

The STAR fixed target program provides an opportunity to extend such measurements at even lower energies. Additionally, Λ global polarization is also influenced by magnetic field at the initial stage. It would be interesting to investigate such a effect towards lower beam energies.

In this talk, measurement of global polarization of Λ hyperons in Au+Au collisions at $\sqrt{s_{\text{NN}}}$ = 7.2 GeV with the fixed-target configuration is reported and compared with the results at other collision energies.

[1]L. Adamczyk et al.(STAR), Nature 548 62 (2017).

[2]J. Adam et al.(STAR), Phys. Rev. C 98 14910 (2018).

Primary author: Mr OKUBO (FOR THE STAR COLLABORATION), Kosuke (University of Tsukuba)

Presenter: Mr OKUBO (FOR THE STAR COLLABORATION), Kosuke (University of Tsukuba)

Session Classification: Contributed Session 5

Track Classification: Track group 2: Experiment