9th International Conference on New Frontiers in Physics (ICNFP 2020)



Contribution ID: 158

Type: Talk

Gluons, Heavy and Light Quarks in the Instanton Liquid Model

Saturday 5 September 2020 11:25 (25 minutes)

We are extending ILM to gluons, heavy quarks, and heavy-light quarks systems. In ILM $4N_c$ instanton collective coordinates = size $\rho\approx\bar{\rho}$, color orientation, position. $\rho\approx 0.3$ fm, inter-instanton distance $R\approx 1$ fm, packing parameter $\lambda=\rho^4/R^4\approx 0.01.$ ILM vacuum energy density $\approx-500\,{\rm MeV/fm}^3.$

Instanton vs hadron sizes.

 $r_{J/\psi} = 0.25$ fm, $r_{\Upsilon} = 0.14$ fm, $r_N \sim 0.3 - 0.5$ fm. Small quark core size hadrons are insensitive to the confinement, ILM safely applicable.

Light quarks in ILM. Dynamical quark mass M(q). $M(0) \approx 360 \text{ MeV} \sim \lambda^{1/2} \rho^{-1} \sim \text{strength of light quark-instanton interaction.}$ Successful reproducing of light hadrons physics with $O(m, 1/N_c, m/N_c)$ corrections.

Gluons in ILM. Dynamical gluon mass $M_g(q)$. $M_g(0) \approx M(0) \sim \lambda^{1/2} \rho^{-1} \sim \text{strength of gluon-instanton interaction.}$

Heavy quarks in ILM. ILM contribution to heavy quark mass $\Delta M(q)$. $\Delta M(0) \approx 70 \text{ MeV} \sim \lambda \rho^{-1} \sim \text{strength of heavy quark-instanton interaction.}$ Heavy quark-antiquark potential V(r) = ILM modified one gluon exchange $V_{ILM,g}(r)$ + direct instanton $V_{\text{dir}}(r)$ + confinement $V_{\text{conf}}(r)$ potentials. $V_{\text{cornell}}(r)$ = one gluon exchange $V_g(r)$ + confinement $V_{\text{conf}}(r)$ potentials. $V(r) \text{ vs } V_{\text{cornell}}(r) \Rightarrow +5 \div 10 \%$ correction for charmonium $(c\bar{c})$ ground state energy.

Heavy+light quarks in ILM. $(c\bar{c})' \rightarrow (c\bar{c})\pi\pi$. Light quark factor $F_{\pi Q} \approx 0.6F_{\pi}$. Heavy quark factor = dipole approximation $(1 + c r_{J/\psi}^2 / \rho^2 + ...)$. $c r_{J/\psi}^2 \approx -0.372 r_{J/\psi}^2 / \rho^2 \approx -0.26$. Standard approach = dipole approximation. Request for ILM reconsideration of heavy quarkonium light hadrons emission processes and light-heavy quarks meson states.

ILM is a framework for uniform and consistent description of light and heavy quark physics.

Is this abstract from experiment?

No

Internet talk

Yes

Name of experiment and experimental site

N/A

Is the speaker for that presentation defined?

Yes

Details

Mirzayusuf Musakhanov, Professor, National University of Uzbekistan, Uzbekistan, https://www.nuu.uz/eng

Primary author: Prof. MUSAKHANOV, Mirzayusuf (National University of Uzbekistan)Presenter: Prof. MUSAKHANOV, Mirzayusuf (National University of Uzbekistan)Session Classification: Workshop on QCD