



# Recent strangeness results from the RHIC beam energy scan

Xianglei Zhu

Tsinghua University

9/10/2020




*9th International Conference on New Frontiers in Physics*

*September 4 – 12, 2020*

*Kolymbari, Crete, Greece*

# QCD phase diagram

B. Müller, BEST Col. Meeting, 2016





- **RHIC Beam Energy Scan**

Cover the intermediate baryon density region

Look for **onset of de-confinement, phase boundary** and critical point

# STAR BES



- STAR:  
Collider experiment at RHIC
- full azimuthal coverage at mid-rapidity
- BES-I (completed)  
 $Au+Au \sqrt{s_{NN}} = 62.4 - 7.7$  GeV
- BES-II (on-going)  
 $Au+Au \sqrt{s_{NN}} = 19.6 - 7.7$  GeV
- Fixed-target (on-going)  
 $Au+Au \sqrt{s_{NN}} = 7.7 - 3.0$  GeV  
 $\mu_B$  up to 721 MeV

# Why strangeness?

Rafelski & Müller, 1982




- Strange quarks

- Not exist in colliding nuclei
- Current mass  $\sim 100 \text{ MeV} < T_c$
- Easily pair-produced in de-confined QGP medium



→ **Strangeness enhancement !**

- Hadrons with (multiple) strange quarks

- Small hadronic cross section
- Sensitive to the early stage dynamics of the medium
- Can be easily reconstructed and identified in experiment, up to high  $p_T$  !



# Particle yields, $dN/dy$ , at mid- $y$ (STAR BES-I)




STAR, PRC96, 044904, 2017

- Yield per participating pair increases towards central and higher energies in general
- Exceptions:
  - $p$  and  $\Lambda$  yields decrease towards higher energy
  - $\bar{p}$  and  $\bar{\Lambda}$  has weak centrality dependence

STAR, arXiv:1906.03732, to appear in PRC

# Particle yields in central collisions





STAR, PRC96, 044904, 2017  
 STAR, arXiv:1906.03732


- STAR BES-I data consistent with published data in general

- Rich structure in these excitation functions

- $p$  and  $\Lambda$  yields reach minimum at 39 GeV: interplay of baryon transport and pair production




# Average transverse mass





STAR, PRC96, 044904, 2017  
 STAR, arXiv:1906.03732

- A step-like structure can be seen in the energy dependence, first-order phase transition?
- $\Lambda$  and  $\bar{\Lambda}$  show split at lower energies might be due to baryon-antibaryon annihilations at high baryon density






# Directed flow (STAR BES-I)



STAR, PRL112, 162301, 2014  
 STAR, PRL120, 062301, 2018





- Sign change of proton  $dv_1/dy$ , softening of EOS, first-order phase transition
- Double sign change seen in net-proton, net- $\Lambda$ , not seen in net-kaon
- Need theory to explain




# $K^+/\pi^+$ ratio (NA61/SHINE)

P. Podlaski, SQM2019





- Plateau like structure visible in p+p
- Be+Be close to p+p
- Ar+Sc is higher than p+p but form of energy dependence is similar to p+p (no horn)

# Strange hadron to pion ratio (STAR BES-I)




## RHIC BES




- Particle ratios consistent with NA49, consistent with the picture of a **maximum net-baryon density around  $\sqrt{s_{NN}} \sim 8$  GeV at freeze-out**


# Anti-hadron to hadron ratio




STAR, PRC96, 044904, 2017  
 STAR, arXiv:1906.03732




- Centrality dependence of  $\bar{B}/B$  ratios:  
**peripheral > central**
- This effect is more prominent at lower energies.  
**baryon stopping and/or anti-baryon absorption**
- **Loss of low  $p_T$   $\bar{\Lambda}$  in central collisions**



# Anti-hadron to hadron ratio



STAR, PRC96, 044904, 2017  
 STAR, arXiv:1906.03732



- STAR BES data lie in a trend with NA49 data

- $\bar{B}/B$  ratios increase with number of strange quarks at low energies

$$\bar{\Omega}^+/\Omega^- > \bar{\Xi}^+/\Xi^- > \bar{\Lambda}/\Lambda > \bar{p}/p$$

# Anti-hyperon to hyperon ratio

$$n_i = \frac{g_i}{(2\pi^2)} \gamma_S^{|S_i|} m_i^2 T K_2(m_i/T) \exp(\mu_i/T)$$

$$\frac{\bar{\Lambda}}{\Lambda} = \exp\left(-\frac{2\mu_B}{T} + \frac{2\mu_S}{T}\right)$$

$$\ln\left(\frac{\bar{\Lambda}}{\Lambda}\right) = -\frac{2\mu_B}{T} + \frac{2\mu_S}{T}$$

$$\frac{\bar{\Xi}^+}{\Xi^-} = \exp\left(-\frac{2\mu_B}{T} + \frac{4\mu_S}{T}\right)$$



$$\ln\left(\frac{\bar{\Xi}^+}{\Xi^-}\right) = -\frac{2\mu_B}{T} + \frac{4\mu_S}{T}$$

$$\frac{\bar{\Omega}^+}{\Omega^-} = \exp\left(-\frac{2\mu_B}{T} + \frac{6\mu_S}{T}\right)$$



$$\ln\left(\frac{\bar{\Omega}^+}{\Omega^-}\right) = -\frac{2\mu_B}{T} + \frac{6\mu_S}{T}$$

- T is the temperature.
- $\mu_B$  is the baryon chemical potential.
- $\mu_S$  is the strangeness chemical potential.

(arXiv:nucl-th/9704046v1 by J.Cleymans & Phys. Rev. C 71(2005)054901)

# $\mu_S/T_{ch}$ and $\mu_B/T_{ch}$



STAR, arXiv:1906.03732



- Anti-hyperon to hyperon ratios are fit well with statistical thermal model
- Chemical freeze-out parameters,  $\mu_S/T_{ch}$  and  $\mu_B/T_{ch}$ , are extracted

# Chemical freeze-out parameters: $T_{ch}$ vs. $\mu_B$

STAR, Phys. Rev. C 96, 044904, 2017



✓ Particles used :  $\pi$ ,  $K$ ,  $p$ ,  $\Lambda$ ,  $\Xi$

✓ Ensemble used:

**Grand canonical (GCE)**

✓ Fit parameters:

$T_{ch}$ ,  $\mu_B$ ,  $\mu_S$  and  $\gamma_S$

Andronic: NPA 834 (2010) 237


Cleymans: PRC 73 (2006) 034905

Au+Au 200 GeV : Phys. Rev. C 83 (2011) 24901

Thermus, S. Wheaton & J. Cleymans, Comput. Phys. Commun. 180: 84-106, 2009.



# Nuclear modification factors $R_{CP}$




$$R_{CP}(p_T) = \frac{[d^2\sigma/(N_{\text{bin}}p_T dp_T dy)]_{\text{central}}}{[d^2\sigma/(N_{\text{bin}}p_T dp_T dy)]_{\text{peripheral}}}$$


- No  $K_S^0$  suppression in Au+Au 7.7 and 11.5 GeV
- Cronin effect and other effects (radial flow) compete with partonic energy loss
- Intermediate  $p_T$ , particle  $R_{CP}$  difference becomes smaller @ 7.7 and 11.5 GeV

STAR, arXiv:1906.03732

# Nuclear modification factors $R_{CP}$




- No suppression for lower energies
- Cronin effect and other effects (radial flow) compete with partonic energy loss



STAR, PRL121, 032301, 2018


# Baryon to meson ratio: $\bar{\Lambda}/K_S^0$



STAR, arXiv:1906.03732

$\sqrt{s_{NN}} < 19.6$  GeV, at intermediate  $p_T$ , the separation of central (0-5%) and peripheral (40-60%) collisions in  $\bar{\Lambda}/K_S^0$  becomes less significant

# $\Omega / \phi$ ratio




STAR, Phys. Rev. C 93, 021903 (R), 2016


➤ Intermediate  $p_T$   $\Omega/\phi$  ratios:

Indication of separation between  $\geq 19.6$  and 11.5 GeV

➤  $\Omega/\phi$  ratios: 40%-60% peripheral < 0-10% central for 19.6, 27 and 39 GeV




# The STAR fixed-target program



A 1 mm thick (4% inter. prob.) gold target




$V_y$  vs.  $V_x$  Distribution



1.3M events from half hour test run, top 30% central trigger, Au+Au  $\sqrt{s_{NN}}=4.5$  GeV


3.4M events from two hour test run, top 30% central trigger, Al+Au  $\sqrt{s_{NN}}=4.9$  GeV

# Hadron spectra and $dN/dy$ in Au+Au $\sqrt{s_{NN}}=4.5$ GeV




E895. Phys. Rev. C 68 (2003) 054905  
 E802. Phys. Rev. C 57 (1998) R466  
 E877. Phys. Rev. C 62 (2000) 024901

- Amplitude and width of rapidity densities are consistent with AGS experiments
- $m_T - m_0$  and  $y$  range will be extended by eTOF and iTPC upgrades




Y. Wu, QM2018  
 Top 5%



# Directed flow in Au+Au $\sqrt{s_{NN}}=4.5$ GeV

E895. Phys. Rev. Lett. 84 (2000) 005488  
 STAR. Phys. Rev. Lett. 112 (2014) 162301



*Y. Wu, QM2018*

Baryon  $v_1$  slope is consistent with E895 at 4.3 GeV



# Summary & outlook

- STAR BES-I have measured systematically the production of strangeness at intermediate baryon density
- Step/horn structures are now investigated at both BES and SPS
- Double sign change seen in directed flow of net-baryons, but not in net-kaons
- QGP signatures appear to turn off at lower collision energies, but need more statistics to confirm
- The ongoing STAR BES-II with detector upgrade (iTPC, eTOF, EPD) and larger luminosity allow precise measurement of the matter properties at intermediate baryon density ( $\mu_B$  up to 721 MeV)