

Supersymmetric theories and Graphene

Wednesday 29 July 2020 19:20 (30 minutes)

We discuss a 2+1 dimensional model holographically realized as the boundary theory of a four-dimensional gravity model in Anti de Sitter (AdS) spacetime. The result is achieved through suitable boundary conditions for the D=4 fields, and an effective model for massive spin-1/2 fields on a curved background is obtained.

The (unconventional) supersymmetry of the boundary model allows to introduce extra internal degrees of freedom, which can provide an application of the model to the description of the charge carriers properties of graphene-like 2D materials at the Dirac points K and K' . In particular, the two valleys correspond to the two independent sectors of the boundary model, connected by a parity transformation. The fermion masses entering the corresponding Dirac equations are related to the torsion parameters of the substrate in the three-dimensional model: the parity-even and odd components of the corresponding masses are identified with Semenoff and Haldane-type mass contributions, respectively.

The construction follows a top-down approach, in that the effective 2+1 dimensional theory for a condensed matter system at the boundary originates from a well-defined supersymmetric effective supergravity in the bulk.

Secondary track (number)

Primary author: GALLERATI, Antonio (Politecnico di Torino)

Presenter: GALLERATI, Antonio (Politecnico di Torino)

Session Classification: Formal Theory

Track Classification: 10. Formal Theory