

# Bekenstein bound from the Pauli principle

*Friday 31 July 2020 13:51 (3 minutes)*

Assuming that the degrees of freedom of a black hole are finite in number and of fermionic nature, we naturally obtain, within a second-quantized toy model of the evaporation, that the Bekenstein bound is a consequence of the Pauli exclusion principle for these fundamental degrees of freedom. We show that entanglement, Bekenstein and thermodynamic entropies of the black hole all stem from the same approach, based on the entropy operator whose structure is the one typical of Takahashi and Umezawa's Thermofield Dynamics. We then evaluate the von Neumann black hole–environment entropy and noticeably obtain a Page-like evolution. We finally show that this is a consequence of a duality between our model and a quantum dissipative-like fermionic system.

## I read the instructions

## Secondary track (number)

**Author:** Dr SMALDONE, Luca (Charles University Prague)

**Co-authors:** IORIO, Alfredo (Institute of Particle and Nuclear Physics, Faculty of Mathematics and Physics, Charles University ,Prague, Czech Republic); ACQUAVIVA, Giovanni (Charles University)

**Presenter:** Dr SMALDONE, Luca (Charles University Prague)

**Session Classification:** Beyond the Standard Model - Posters

**Track Classification:** 10. Formal Theory