

# A new way of understanding the role of each measurement in global SMEFT fit @ $e^+e^-$


(paper in preparation)

Keisuke Fujii (KEK), Michael Peskin (SLAC), [Junping Tian \(U. Tokyo\)](#)

ICHEP2020, July 31, 2020

based on earlier work: Barklow et al, arXiv:[1708.09079](#); [1708.08912](#)

# global SMEFT fit & roles of measurements



- When  $m_{\text{BSM}} \gg m_{\text{EW}}$ , all the SM measurements can fit into a SMEFT framework, providing coherent tests of BSM physics
- A global EFT fit involves many fitting parameters and many input measurements, making the accurate understanding of roles of each measurement increasingly difficult
- Roles of **EWPO / TGC / Beam polarizations / Top EW couplings** for Higgs coupling determination @  $e^+e^-$  get highly recognized

this talk:

- ☑ a brief recap on qualitative & quantitative understanding
- ☑ a new way of more transparent understanding

# global SMEFT fit @ future e+e-

[Barklow, Fujii, Jung, Peskin, JT, arXiv:1708.09079]

$$\begin{aligned} \Delta\mathcal{L} = & \frac{c_H}{2v^2} \partial^\mu (\Phi^\dagger \Phi) \partial_\mu (\Phi^\dagger \Phi) + \frac{c_T}{2v^2} (\Phi^\dagger \overleftrightarrow{D}^\mu \Phi) (\Phi^\dagger \overleftrightarrow{D}_\mu \Phi) - \frac{c_6 \lambda}{v^2} (\Phi^\dagger \Phi)^3 \\ & + \frac{g^2 c_{WW}}{m_W^2} \Phi^\dagger \Phi W_{\mu\nu}^a W^{a\mu\nu} + \frac{4gg' c_{WB}}{m_W^2} \Phi^\dagger t^a \Phi W_{\mu\nu}^a B^{\mu\nu} \\ & + \frac{g'^2 c_{BB}}{m_W^2} \Phi^\dagger \Phi B_{\mu\nu} B^{\mu\nu} + \frac{g^3 c_{3W}}{m_W^2} \epsilon_{abc} W_{\mu\nu}^a W^{b\nu\rho} W^{c\rho\mu} \\ & + i \frac{c_{HL}}{v^2} (\Phi^\dagger \overleftrightarrow{D}^\mu \Phi) (\bar{L} \gamma_\mu L) + 4i \frac{c'_{HL}}{v^2} (\Phi^\dagger t^a \overleftrightarrow{D}^\mu \Phi) (\bar{L} \gamma_\mu t^a L) \\ & + i \frac{c_{HE}}{v^2} (\Phi^\dagger \overleftrightarrow{D}^\mu \Phi) (\bar{e} \gamma_\mu e) . \end{aligned}$$

“Warsaw” basis,  
Grzadkowski et al,  
arXiv:1008.4884

$\Phi$ : higgs field  
W, B: SU(2), U(1) gauge  
L, e: left/right electron

- most of the non-trivial relations come from 7 operators

$$\begin{array}{ccccccc} c_H & & & & & & \\ & c_{HL} & c'_{HL} & c_{HE} & & & \\ & & & & c_{WW} & c_{WB} & c_{BB} \end{array}$$

## recap 1: absolute Higgs couplings (unique role of inclusive $\sigma_{Zh}$ )

$$\frac{c_H}{2v^2} \partial^\mu (\Phi^\dagger \Phi) \partial_\mu (\Phi^\dagger \Phi)$$

$$\frac{c_H}{2} \partial^\mu h \partial_\mu h \quad \longrightarrow \quad \text{renormalize SM Higgs field}$$

$$h \quad \longrightarrow \quad (1 - c_H/2)h$$

$\longrightarrow$  **shift all SM Higgs couplings by  $-c_H/2$**


- $c_H$  can not be determined by any BR or ratio of couplings
- $c_H$  has to rely on inclusive cross section of  $e^+e^- \rightarrow Zh$ , enabled by recoil mass technique at  $e^+e^-$

(precision of  $hZZ$ ,  $hWW \sim 1/2 \Delta c_H$ )


## recap 2: role of Electroweak Precision Observable (EWPO)

$$i \frac{c_{HL}}{v^2} (\Phi^\dagger \overleftrightarrow{D}^\mu \Phi) (\bar{L} \gamma_\mu L)$$


$$+ (c'_{HL}, c_{HE})$$



$e^+e^- \rightarrow Zhh$



$e^+e^- \rightarrow Zh$



Z-pole

- very useful EWPO at Z-pole:  $\mathbf{A}_{LR}, \mathbf{\Gamma}_{Z \rightarrow ee}$

- Z-e-e couplings can also get helped by  $\sigma_{zh}$ :  $\delta\sigma_{zh} \sim \mathbf{s}/m^2_z$

## recap 3: roles of Higgs measurements at (HL-)LHC

$$\boxed{\frac{g^2 c_{WW}}{m_W^2} \Phi^\dagger \Phi W_{\mu\nu}^a W^{a\mu\nu}} + (c_{BB}, c_{WB})$$

$$\delta\Gamma(h \rightarrow \gamma\gamma) = -c_H + 122(8c_{WW} - 16c_{WB} + 8c_{BB}) + \dots$$

$$\delta\Gamma(h \rightarrow \gamma Z) = -c_H + 122(8c_{WW} - 5.6c_{WB} - 2.4c_{BB}) + \dots$$


$$\delta\Gamma(h \rightarrow ZZ^*) = -c_H - 0.4(8c_{WW} + 3.7c_{WB} + 0.6c_{BB}) + \dots$$

- loop induced  $h \rightarrow \gamma\gamma/\gamma Z$  depend strongly on  $c_{WW}/c_{BB}/c_{WB}$
- very useful measurements: **BR(h → γγ/γZ)** at (HL-)LHC



$$R_{\gamma\gamma} = \frac{BR(h \rightarrow \gamma\gamma)}{BR(h \rightarrow ZZ^*)} \quad R_{\gamma Z} = \frac{BR(h \rightarrow \gamma Z)}{BR(h \rightarrow ZZ^*)}$$

use ratio of BR to keep model-independence

## recap 4: role of beam polarizations



$c_{WW}$



$c_{HL} + c'_{HL}$

$c_{HE}$

$P(e^-, e^+)$

$\sqrt{s}=250 \text{ GeV}$

$$(-0.8, +0.3) \quad \delta\sigma_L = -c_H + 10(8c_{WW}) + 27(c_{HL} + c'_{HL}) + \dots$$


$$(+0.8, -0.3) \quad \delta\sigma_R = -c_H + 1.6(8c_{WW}) + 2.1(c_{HL} + c'_{HL}) + \dots$$

$$(0, 0) \quad \delta\sigma_0 = -c_H + 6.6(8c_{WW}) + 16(c_{HL} + c'_{HL}) + \dots$$



- $\sigma_R$  has much weaker dependences on  $c_{WW}$  &  $c_{HL}+c_{HL}'$  (suppression of chiral new physics effects)  
-> results in better determination of  $c_H$

- redundant  $\sigma_L$  in turn improves  $c_{WW}$ ,  $c_{HL}+c_{HL}'$

# recap 4: role of beam polarizations



$c_{WW}$



$c_{HL} + c'_{HL}$

$c_{HE}$

with ILC inputs  $\sqrt{s}=250$  GeV

$\int L dt @ P(e^-, e^+)$	$\Delta c_H$	$\Delta(c_{HL} + c_{HL}')$	$\Delta(8c_{WW})$	$\Delta c_{HE}$
2 ab <sup>-1</sup> @ (0,0)	148	9.2	14	4.0
2 ab <sup>-1</sup> @ (+0.8,-0.3)	106	9.2	14	4.0
1 ab <sup>-1</sup> @ (+0.8,-0.3) 1 ab <sup>-1</sup> @ (-0.8,+0.3)	104	5.95	12	3.2
2 ab <sup>-1</sup> @ (-0.8,+0.3)	245	9.23	14	4.0
5 ab <sup>-1</sup> @ (0,0)	134	8.84	13	3.1


$\Delta$  in unit of  $10^{-4}$




## classical way of evaluating roles of measurements

- do global fit numerically: vary certain input measurements, and see how the precision of Higgs couplings vary
- hard to figure out synergies among multiple ( $\gg 2$ ) measurements

—> Is there a more transparent way?



[Bambade, et al, arXiv:1903.01629]



[de Blas, et al, arXiv:1907.04311]

## a new way for more transparent understanding

can we solve the global fit analytically?

we would like to express the uncertainty in a Higgs coupling ( $\Delta g$ )  
analytically in terms of the uncertainties in observables ( $\Delta O$ )

e.g. 
$$\Delta g_{hXX} = x_1 \Delta O_1 \oplus x_2 \Delta O_2 \oplus x_3 \Delta O_3 \oplus \dots$$

(all in physical quantities; should be EFT basis-independent)

as an intermediate step, we must get first the expression for  
the uncertainties in Wilson coefficients ( $\Delta c$ )

## basic notations (I)

$c_i$   $i = 1, 2, \dots, n$  Fitting parameters (Wilson coefficients)  
 $n$  = number of fitting parameters

$y_i$   $i = 1, 2, \dots, m$  Observables (deviation w.r.t. SM values)  
 $m$  = number of observables

$y_i = v_{ij} c_j$   $j = 1, 2, \dots, n$   $v_{ij}$  known from theory computation  
 $i = 1, 2, \dots, m$

in matrix form

$$y = Vc$$

$y$ : column vector of  $y_i$ ,  $m \times 1$   
 $c$ : column vector of  $c_j$ ,  $n \times 1$   
 $V$ : matrix of  $v_{ij}$ ,  $m \times n$

## basic notations (II)

in matrix form

$$\mathbf{y} = \mathbf{V}\mathbf{c}$$

$\mathbf{y}$ : column vector of  $y_i$ ,  $m \times 1$   
 $\mathbf{c}$ : column vector of  $c_j$ ,  $n \times 1$   
 $\mathbf{V}$ : matrix of  $v_{ij}$ ,  $m \times n$

measurements

$$\mathbf{E}_y$$

covariance matrix for all observables  $y_i$

which can be diagonalized

$$\mathbf{E}_y = \begin{pmatrix} \Delta_1^2 & 0 & \cdot & 0 \\ 0 & \Delta_2^2 & \cdot & 0 \\ \cdot & \cdot & \cdot & \cdot \\ 0 & 0 & \cdot & \Delta_m^2 \end{pmatrix}$$

where  $\Delta_i$  is measurement uncertainty for  $\mathbf{y}_i$

## global fit

minimizing

$$\begin{aligned}\chi^2 &= y^T \mathbf{E}_y^{-1} y \\ &= c^T \mathbf{V}^T \mathbf{E}_y^{-1} \mathbf{V} c = c^T \mathbf{D} c\end{aligned}$$

$$\mathbf{D} \equiv \mathbf{V}^T \mathbf{E}_y^{-1} \mathbf{V} \quad (i,j) \text{ element } d_{ij} = \sum_{k=1}^m \frac{v_{ki} v_{kj}}{\Delta_k^2}$$

$\mathbf{D}^{-1}$  is exactly the covariance matrix of fitting parameters

**global fit = how to obtain  $\mathbf{D}^{-1}$  = how to invert  $\mathbf{V}^T \mathbf{E}_y^{-1} \mathbf{V}$**

numerical solution is easy, e.g. Barklow et al, [1708.09079](#); [1708.08912](#)

## global fit: analytic solution

$$\mathbf{D} \equiv \mathbf{V}^T \mathbf{E}_y^{-1} \mathbf{V}$$

**global fit = how to obtain  $\mathbf{D}^{-1}$  = how to invert  $\mathbf{V}^T \mathbf{E}_y^{-1} \mathbf{V}$**

dimensions:  $\mathbf{V}$ :  $m \times n$   
 $\mathbf{E}_y$ :  $m \times m$   $\mathbf{D}$ :  $n \times n$   
 $\mathbf{V}^T$ :  $n \times m$

- **if  $m = n$** , namely there is no redundant measurement

solution is easy:

$$\mathbf{D}^{-1} = \mathbf{V}^{-1} \mathbf{E}_y (\mathbf{V}^T)^{-1}$$

$\mathbf{V}$  is invertible, otherwise global fit doesn't converge

## analytic solution: application of non-redundant case

for unpolarized  $e^+e^-$  at 250 GeV, there is almost no redundant measurement  
 (except W-fusion  $\nu\nu h$ , angular  $Zh$ ; either small contribution)

$$\delta g_{hZZ} = \frac{1}{2} \delta \sigma_{Zh} + 6.4 \delta \Gamma_l + 5.3 \delta g_{Z,eff} - 0.015 \delta R_{\gamma Z} - 2.4 \delta \kappa_{A,eff} + 8.9 \delta m_h + 0.098 \delta A_l + \dots$$

plug in measurement uncertainties for current EWPO + 2  $\text{ab}^{-1}$

$$= 41 \oplus 66 \oplus 30 \oplus 23 \oplus 14 \oplus 11 \oplus 8.7 \oplus \dots \times 10^{-4}$$

importance hierarchy

## analytic solution: application of non-redundant case

- **if  $m = n$** , namely there is no redundant measurement

$$\mathbf{D}^{-1} = \mathbf{V}^{-1} \mathbf{E}_y (\mathbf{V}^T)^{-1}$$

the expression in previous slide can be abstracted as

$$\Delta^2 c_i = \sum_{k=1}^n \frac{|\bar{V}_{ki}|^2}{|\mathbf{V}|^2} \Delta_k^2 = \frac{\sum_S C_{n-1}^n \frac{|V_S^i|^2}{\Delta_S^2}}{\sum_L C_n^n \frac{|V_L|^2}{\Delta_L^2}}$$



## global fit: analytic solution for general case ( $m > n$ )

$$\Delta^2 c_i = \frac{\sum_S C_{n-1}^m \frac{|V_S^i|^2}{\Delta_S^2}}{\sum_L C_n^m \frac{|V_L|^2}{\Delta_L^2}}$$

$$L = \{l_1, l_2, \dots, l_n\}$$

**n**-combination of  $\{1, 2, \dots, \mathbf{m}\}$

$$\Delta_L = \prod_{i=1}^n \Delta_{l_i}$$

$V_L$   $n \times n$  matrix formed by Rows **L** of  $V$

$$S = \{s_1, s_2, \dots, s_{n-1}\}$$

**(n-1)**-combination of  $\{1, 2, \dots, \mathbf{m}\}$

$$\Delta_S = \prod_{i=1}^{n-1} \Delta_{s_i}$$

$V_S^i$   $n-1 \times n-1$  matrix formed by Rows **S** of  $V$   
& **eliminating Column  $i$**


## summary

- highlighted a few important roles in global SMEFT fit, played by measurements  $\sigma_{Zh}$  at  $e^+e^-$  /  $\text{BR}(h \rightarrow \gamma\gamma / \gamma Z)$  at LHC / Z-pole / Beam polarizations
- developed a new way for more transparent understanding which is based on analytic expressions in terms of meas. uncertainties
- applied to non-redundant case: clear synergies among many meas.
- stay tuned for applications to redundant cases (of high interests: polarizations; multiple ECM; multiple Higgs prod. channels)


backup

# role of top-quark EW couplings


arXiv:2006.14631



$$\mathcal{O}_{Ht} = (\Phi^\dagger i \overleftrightarrow{D}_\mu \Phi) (\bar{t} \gamma^\mu t)$$



$$\mathcal{O}_{Hq}^{(3)} = (\Phi^\dagger i \overleftrightarrow{D}_\mu^a \Phi) (\bar{Q} \gamma^\mu \tau^a Q)$$



$$\mathcal{O}_{tB} = (\bar{Q} \sigma^{\mu\nu} t) \tilde{\Phi} B_{\mu\nu}$$


- NLO effects from top-quark operators are very important for Higgs couplings determination
- top-quark EW couplings measurements @ (HL-)LHC are invaluable for future e+e-

see talk by Martín Perelló @ top/EW session today

# roles of Triple Gauge Couplings (TGC) in $e^+e^- \rightarrow WW$

$$\boxed{\frac{4gg'c_{WB}}{m_W^2} \Phi^\dagger t^a \Phi W_{\mu\nu}^a B^{\mu\nu}} + (c_{WW}, c_{BB})$$

$e^+e^- \rightarrow WW$



$h \rightarrow ZZ$

$$\zeta_Z \frac{h}{2v} Z_{\mu\nu} Z^{\mu\nu}$$

$$\zeta_Z = c_w^2(8c_{WW}) + 2s_w^2(8c_{WB}) + s_w^4/c_w^2(8c_{BB})$$

- longitudinal modes of W/Z are from Higgs fields
- $c_{WB} / c_{HL}' / c_{3W}$  helped by meas. of TGCs in  $e^+e^- \rightarrow WW$

# Higgs couplings are related to themselves

$$\begin{aligned}
 \Delta\mathcal{L}_h = & \frac{1}{2}\partial_\mu h\partial^\mu h - \frac{1}{2}m_h^2 h^2 - (1 + \eta_h)\bar{\lambda}vh^3 + \frac{\theta_h}{v}h\partial_\mu h\partial^\mu h \\
 & + (1 + \eta_W)\frac{2m_W^2}{v}W_\mu^+W^{-\mu}h + (1 + \eta_{WW})\frac{m_W^2}{v^2}W_\mu^+W^{-\mu}h^2 \\
 & + (1 + \eta_Z)\frac{m_Z^2}{v}Z_\mu Z^\mu h + \frac{1}{2}(1 + \eta_{ZZ})\frac{m_Z^2}{v^2}Z_\mu Z^\mu h^2 \\
 & + \zeta_W\hat{W}_{\mu\nu}^+\hat{W}^{-\mu\nu}\left(\frac{h}{v} + \frac{1}{2}\frac{h^2}{v^2}\right) + \frac{1}{2}\zeta_Z\hat{Z}_{\mu\nu}\hat{Z}^{\mu\nu}\left(\frac{h}{v} + \frac{1}{2}\frac{h^2}{v^2}\right) \\
 & + \frac{1}{2}\zeta_A\hat{A}_{\mu\nu}\hat{A}^{\mu\nu}\left(\frac{h}{v} + \frac{1}{2}\frac{h^2}{v^2}\right) + \zeta_{AZ}\hat{A}_{\mu\nu}\hat{Z}^{\mu\nu}\left(\frac{h}{v} + \frac{1}{2}\frac{h^2}{v^2}\right).
 \end{aligned}$$

(SM structure: kappa like)

$$\eta_h = \delta\bar{\lambda} + \delta v - \frac{3}{2}c_H + c_6$$

$$\eta_W = 2\delta m_W - \delta v - \frac{1}{2}c_H$$

$$\eta_{WW} = 2\delta m_W - 2\delta v - c_H$$

$$\eta_Z = 2\delta m_Z - \delta v - \frac{1}{2}c_H - c_T$$

$$\eta_{ZZ} = 2\delta m_Z - 2\delta v - c_H - 5c_T$$

(Anomalous: new Lorentz structure)

$$\theta_h = c_H$$

$$\zeta_W = \delta Z_W = (8c_{WW})$$

$$\zeta_Z = \delta Z_Z = c_w^2(8c_{WW}) + 2s_w^2(8c_{WB}) + s_w^4/c_w^2(8c_{BB})$$

$$\zeta_A = \delta Z_A = s_w^2\left((8c_{WW}) - 2(8c_{WB}) + (8c_{BB})\right)$$

$$\zeta_{AZ} = \delta Z_{AZ} = s_w c_w \left( (8c_{WW}) - \left(1 - \frac{s_w^2}{c_w^2}\right)(8c_{WB}) - \frac{s_w^2}{c_w^2}(8c_{BB}) \right)$$

- hZZ/hWW/hγZ/hγγ highly related: SU(2)xU(1) gauge symmetries

# role of each measurement: more transparent understanding

for example: unpolarized e+e- at 250 GeV

$$\delta g_{hZZ} = \frac{1}{2} \delta \sigma_{Zh} + 6.4 \delta \Gamma_l + 5.3 \delta g_{Z,eff} - 0.015 \delta R_{\gamma Z} - 2.4 \delta \kappa_{A,eff} + 8.9 \delta m_h + 0.098 \delta A_l + \dots$$

$$\delta X = \frac{\Delta X}{X}$$

- $\sigma_{Zh}$  : cross section of e+e- -> Zh
- $A_l, \Gamma_l$  :  $A_{LR}$  and  $\Gamma(Z \rightarrow ll)$  at Z-pole
- $g_{Z,eff}, \kappa_{A,eff}$  : Triple Gauge Couplings
- $R_{\gamma Z}$  :  $BR(h \rightarrow \gamma Z) / BR(h \rightarrow ZZ^*)$
- $m_h$  : Higgs mass

# role of each measurement: more transparent understanding

for example: unpolarized  $e^+e^-$  at 250 GeV

plug in measurement precisions for current EWPOs + 2 ab<sup>-1</sup>

$$\begin{aligned}
 \delta g_{hbb} &= \frac{1}{2} \delta B_{bb} - \frac{1}{2} \delta B_{WW} + \frac{1}{2} \delta \sigma_{Zh} - 5.79 \delta \Gamma_l - 0.016 \delta \Gamma_{\gamma Z} + \dots \\
 &= 28 \oplus 91 \oplus 41 \oplus 59 \oplus 32 \oplus \dots \times 10^{-4}
 \end{aligned}$$

$\text{BR}(h \rightarrow bb)$        $\text{BR}(h \rightarrow WW)$        $\sigma_{Zh}$       EWPOs       $\text{BR}(h \rightarrow \gamma Z)$




# role of each measurement: more transparent understanding

for example: unpolarized e+e- at 250 GeV

plug in measurement precisions for current EWPOs + 2 ab-1

$$\delta g_{hcc} = \frac{1}{2} \delta B_{cc} - \frac{1}{2} \delta B_{WW} + \frac{1}{2} \delta \sigma_{Zh} - 5.79 \delta \Gamma_l - 0.016 \delta \Gamma_{\gamma Z} + \dots$$

$$= 160 \oplus 91 \oplus 41 \oplus 59 \oplus 32 \oplus \dots \times 10^{-4}$$



## global fit: analytic answer

correlation between  $c_i$  &  $c_j$

$$\rho_{ij} \equiv \frac{e_{ij}}{\sqrt{e_{ii}e_{jj}}} = \frac{(-1)^{i+j} \sum_S C_{n-1}^m \frac{|V_S^i V_S^j|}{\Delta_S^2}}{\sqrt{\sum_S C_{n-1}^m \frac{|V_S^i|^2}{\Delta_S^2} \sum_S C_{n-1}^m \frac{|V_S^j|^2}{\Delta_S^2}}}$$

## global fit: analytic solution

$$\mathbf{D} \equiv \mathbf{V}^T \mathbf{E}_y^{-1} \mathbf{V}$$

**global fit = how to obtain  $\mathbf{D}^{-1}$  = how to invert  $\mathbf{V}^T \mathbf{E}_y^{-1} \mathbf{V}$**

- what **if  $m > n$** , namely there are redundant measurements?

some hints from  $n=1,2,3$  cases,

e.g., if  $n=1$  (only one parameter  $c$ )

$$\frac{1}{\Delta_c^2} = \sum_{k=1}^m \frac{v_k^2}{\Delta_k^2} = \frac{1}{\Delta_{c1}^2} + \frac{1}{\Delta_{c2}^2} + \dots + \frac{1}{\Delta_{cm}^2}$$

## global fit: analytic answer

uncertainty of each fitting parameter: reorganize

$$\frac{1}{\Delta c_i^2} = \frac{\sum_L C_n^m \frac{|V_L|^2}{\Delta_L^2}}{\sum_S C_{n-1}^m \frac{|V_S^i|^2}{\Delta_S^2}} = \sum_L \frac{C_n^m}{\sum_S C_{n-1}^m \frac{|V_S^i|^2}{|V_L|^2} \frac{\Delta_L^2}{\Delta_S^2}}$$