

Z polarization as a probe of anomalous gauge-Higgs couplings

Thursday, 30 July 2020 11:30 (15 minutes)

We show how the Z boson polarization can be used to study the ZH production at future e^+e^- colliders and at the LHC. We calculate the 8 independent polarization parameters of the spin-1 Z boson which are sensitive to anomalous gauge-Higgs couplings, using Z spin density matrix. By constructing angular asymmetries from the Z boson decay leptons, which are related to the polarization observables, we constrain the anomalous ZZH couplings. Taking into account possible longitudinal beam polarizations at two different center of mass energies, we find that oppositely polarized beams at 500 GeV c.m. energy provides tighter bounds on the couplings than the same sign polarized and unpolarized beams. We find that most of the 1σ limits are of the order of a few times

10^{-3} for 14 TeV LHC with integrated luminosity of 1000 fb^{-1} and for 500 GeV e^+e^- colliders with oppositely polarized beams.

Secondary track (number)

Primary authors: Ms SARMAH, Priyanka (IIT Bombay); Prof. RAO, Kumar (IIT Bombay); Prof. RINDANI, Saurabh D. (Physical Research Laboratory, Ahmedabad)

Presenter: Ms SARMAH, Priyanka (IIT Bombay)

Session Classification: Beyond the Standard Model

Track Classification: 03. Beyond the Standard Model