



# First observation of production of three massive gauge bosons



Philip Chang on behalf of CMS collaboration

ICHEP  
July 29, 2020



Univ. of California  
San Diego



# Discovery of Higgs boson

July 4, 2012




Discovery advanced our knowledge of origin of mass in a major way

## Quarks




## Leptons

## Forces



- Is it the only Higgs boson? (or are there more?)
- Are multi-*bosons* interactions SM?
- Are there more states involved in electroweak symmetry breaking?
- Is the Higgs potential SM?


Many more to be studied on electroweak sector at the LHC



Triboson process has access to studying many multi-*boson* interactions


- ATLAS searched for WWW in 8 TeV:  $0.96\sigma$  ( $1.05\sigma$ ) arXiv:1610.05088
- CMS searched for WWW in 13 TeV  $36 \text{ fb}^{-1}$ :  $0.6\sigma$  ( $1.78\sigma$ ) arXiv:1905.04246
- ATLAS searched for VVV in 13 TeV  $80 \text{ fb}^{-1}$ :  $4.1\sigma$  ( $3.1\sigma$ ) arXiv:1903.10415

## VVV evidence




arXiv:1903.10415

## Axion-like-particle triboson signature limit



arXiv:1905.04246

## SMEFT Dim8 operator limit



arXiv:1610.05088

Both ATLAS and CMS have been searching for triboson processes and using them to test SM and constrain new physics




We are targeting all possible VVV productions both w/ and w/o Higgs:

- $pp \rightarrow WWW$
- $pp \rightarrow WWZ$
- $pp \rightarrow WZZ$
- $pp \rightarrow ZZZ$


And the combined production of all  $pp \rightarrow VVV$

Production cross section decreases with more Z's



Viable final states have O(fb) or less cross sections

Production cross section decreases with more Z's




Difficult final states  
w/ bkg rate of  
several orders of  
magnitude larger

Viable final states have O(fb) or less cross sections



Production cross section decreases with more Z's



Difficult final states w/ bkg rate of several orders of magnitude larger

$ZZZ \rightarrow 6L$   
( $L = e, \mu$ )

**11 attobarn**  
(~1.5 events produced at Run 2 of LHC)





Viable

Viable final states have O(fb) or less cross sections

# CMS detector measures leptons very well

$e/\mu$  among the **best** measured particles at CMS by combining tracker, calorimeter, and chambers measurements

(1-2% resolution for well measured ones)



Excellent lepton reconstruction and simulation at CMS

# Overview of the analysis


Target “fully” leptonic final states to go after first observation

One exception

|                 | Same-sign  | 3 leptons  | 4 leptons  | 5 leptons  | 6 leptons  |
|-----------------|--|--|--|--|--|
| Targeted signal | $W^\pm \rightarrow l^\pm \nu$<br>$W^\pm \rightarrow l^\pm \nu$<br>$W^\mp \rightarrow qq$                 | $W \rightarrow l\nu$<br>$W \rightarrow l\nu$<br>$W \rightarrow l\nu$       | $W \rightarrow l\nu$<br>$W \rightarrow l\nu$<br>$Z \rightarrow ll$ | $W \rightarrow l\nu$<br>$Z \rightarrow ll$<br>$Z \rightarrow ll$ | $Z \rightarrow ll$<br>$Z \rightarrow ll$<br>$Z \rightarrow ll$ |
| Dominant Bkgs.  | $WZ \rightarrow l^\pm \nu   l^\pm   \bar{l}^\mp$ (lost)<br>$t\bar{t} \rightarrow bb + l + X$<br>↳ fake l | $WZ \rightarrow l\nu ll$<br>$t\bar{t} \rightarrow bb + ll + X$<br>↳ fake l | $ZZ \rightarrow ll ll$<br>$ttZ \rightarrow ll ll + bbX$            | $ZZ \rightarrow ll ll$<br>+ fake lep                             | $ZZ \rightarrow ll ll$<br>+ 2 fake lep                         |

N.B.  $WZ \rightarrow 3l \sim 100k$ ,  $ZZ \rightarrow 4l \sim 10k$

Separate the analysis categories by N of leptons



Backgrounds are suppressed via disfavored decay topology of  $Z \rightarrow \tau\tau \rightarrow e\mu$

# Splitting signal regions by lepton flavors

|                    | Same-sign<br>2 leptons        | 3 leptons  | 4 leptons   | 5 leptons             | 6 leptons                              |
|--------------------|-------------------------------|--|---|-----------------------|--|
| Targeted<br>signal | $W^\pm \rightarrow l^\pm \nu$ | $W \rightarrow l \nu$  | $W \rightarrow l \nu$   | $W \rightarrow l \nu$ | $Z \rightarrow ll$                     |
|                    | $W^\pm \rightarrow l^\pm \nu$ | $W \rightarrow l \nu$  | $W \rightarrow l \nu$   | $Z \rightarrow ll$    | $Z \rightarrow ll$                     |
|                    | $W^\mp \rightarrow qq$        | $W \rightarrow l \nu$  | $Z \rightarrow ll$  | $Z \rightarrow ll$    | $Z \rightarrow ll$                     |
|                    | Split by<br>$ee/e\mu/\mu\mu$  | Split by<br># of<br>SFOS<br><br>e.g.<br>0: $e^\pm \mu^\mp e^\pm$<br>1: $e^\pm e^\mp \mu^\pm$<br>2: $e^\pm e^\mp e^\pm$ | tag $Z \rightarrow ll$<br>then split<br>$WW \rightarrow ee/\mu\mu$<br>v.<br>$WW \rightarrow e\mu$ |                       | Not enough<br>statistics<br>single bin |
|                    | 3 categories*                 | 3 categories   | 2 categories*   | 1 category            | 1 category                             |

\* marked ones will be further split


Each N lepton analysis is further split by flavors

# Splitting signal regions by lepton flavors



|                    | Same-sign<br>2 leptons  | 3 leptons   | 4 leptons   | 5 leptons  | 6 leptons  |
|--------------------|---|---|---|--|--|
| Targeted<br>signal | $W^\pm$<br>$W^\pm$<br>$W^\mp \rightarrow qq$  | $W$<br>$W$<br>$W \rightarrow l\nu$  | $V$<br>$V$<br>$Z \rightarrow ll$  | $W \rightarrow l\nu$<br>$Z \rightarrow ll$<br>$Z \rightarrow ll$ | $Z \rightarrow ll$<br>$Z \rightarrow ll$<br>$Z \rightarrow ll$ |
|                    | <p>Also have BDT-based analysis</p> <p>Split by <math>ee/e\mu/\mu\mu</math></p> <p>N.B. <math>\mu</math> is "cleaner" than e</p> <p>3 categories*</p> | <p>Also have BDT-based analysis</p> <p>Split by # of SFOS</p> <p>e.g.</p> <p>0: <math>e^\pm\mu^\mp e^\pm</math></p> <p>1: <math>e^\pm e^\mp \mu^\pm</math></p> <p>2: <math>e^\pm e^\mp e^\pm</math></p> <p>3 categories</p> | <p>Also have BDT-based analysis</p> <p>tag <math>Z \rightarrow ll</math> then split</p> <p><math>WW \rightarrow ee/\mu\mu</math></p> <p>v.</p> <p><math>WW \rightarrow e\mu</math></p> <p>2 categories*</p> | <p>Not enough statistics single bin</p> <p>1 category</p>        | <p>1 category</p>  |
|                    | * marked ones will be further split   |   |   |  |  |

Each N lepton analysis is further split by flavors

# Same-sign channel categorization




- $N_J \geq 2$ 
  - Two jets satisfy  $|m_{jj} - m_W| < 15 \text{ GeV}$  ( **$m_{jj}$ -in**)
  - Two jets satisfy  $|m_{jj} - m_W| \geq 15 \text{ GeV}$  ( **$m_{jj}$ -out**)
- $N_J = 1$ 
  - Only one jet exists (**1J**)



3 additional categories ( $m_{jj}$ -in,  $m_{jj}$ -out, 1J) each split by  $ee/e\mu/\mu\mu$   
 $\Rightarrow$  Total of 9 signal regions for same-sign analysis



- Separated by # of SFOS pairs:
  - 0 SFOS (low bkg.)
  - 1 SFOS
  - 2 SFOS
- 0SFOS is by far the cleanest
- One can further reduce WZ backgrounds in other SFOS channels
- e.g. For 1SFOS it is clear which lepton is from W:
 

|                         |                     |                             |                   |
|-------------------------|---------------------|-----------------------------|-------------------|
| $\frac{e^\pm e^\mp}{Z}$ | $\frac{\mu^\pm}{W}$ | $\frac{\mu^\pm \mu^\mp}{Z}$ | $\frac{e^\pm}{W}$ |
| Z                       | W                   | Z                           | W                 |
- (Similar strategy employed for 2SFOS)




By flavor, W lepton can be identified and kinematic endpoints can be used  
 $\Rightarrow$  Total of 3 signal regions for 3 lepton analysis






- Utilize  $m_{T2}$  variable: generalization of  $m_T$  for multiple missing particles
- $m_{T2}$  is sensitive to the end points of  $m_W$  from  $ZWW \rightarrow ll e \mu$
- $m_{T2}$  is sensitive to the end points of  $m_\tau$  from  $ZZ \rightarrow ll \tau \tau \rightarrow ll e \mu$




Exploit differences between  $Z \rightarrow ll \nu$ .  $WW \rightarrow ll \nu$

Trained two BDTs: WWZ v. ZZ and WWZ v. ttZ  
Below shows the 2D plane in BDT scores




5 bins are created from 2D planes



\*\*For  $Z \rightarrow ll + ee/\mu\mu$  event category, 2 bins are created (not shown)

Created multiple bins in BDTs to maximize sensitivity  
 $\Rightarrow$  Total of 7 signal regions for 4 lepton analysis


# 5 leptons



Require the 5 lepton events to contain two SFOS pair consistent with  $Z$  mass

The dominant background is  $ZZ \rightarrow 4l$  plus a fake lepton

The fake lepton has low transverse mass while the signal's  $W$  has transverse mass peaking at  $W$  mass




Simple cut-and-count  $\Rightarrow$  Total of 1 bin in 5 lepton

Select at least 6 leptons

Require  $\Sigma P_T \geq 250$  GeV

Less than 1 event expected

Very clean channel



Not enough stats, so search inclusively  $\Rightarrow$  Total of 1 bin in 6 lepton

# Putting it all together

|                         | Same-sign<br>2 leptons   | 3 leptons  | 4 leptons  | 5 leptons  | 6 leptons  |
|-------------------------|--|--|--|--|--|
| Signals                 | $W^\pm \rightarrow l^\pm \nu$<br>$W^\pm \rightarrow l^\pm \nu$<br>$W^\mp \rightarrow qq$ | $W \rightarrow l\nu$<br>$W \rightarrow l\nu$<br>$W \rightarrow l\nu$ | $W \rightarrow l\nu$<br>$W \rightarrow l\nu$<br>$Z \rightarrow ll$ | $W \rightarrow l\nu$<br>$Z \rightarrow ll$<br>$Z \rightarrow ll$ | $Z \rightarrow ll$<br>$Z \rightarrow ll$<br>$Z \rightarrow ll$ |
| Split Flavor            | 3  | 3  | 2  | 1  | 1  |
| Channel specific splits | mjj-in<br>mjj-out<br>1J  | -  | Split in kinematics or BDT   | -  | -  |
| Total                   | 9 bins   | 3 bins   | 7 bins   | 1 bin  | 1 bin  |

Total of 21 bins



|                   | Same-sign<br>2 leptons  | 3 leptons  | 4 leptons   | 5 leptons                            | 6 leptons                              |
|-------------------|---|--|---|--------------------------------------|--|
| Dominant<br>Bkgs. | $WZ \rightarrow l\bar{\nu}l\bar{\nu}$ (lost)<br>$t\bar{t} \rightarrow bb + l + X$<br>$\hookrightarrow$ fake $l$ | $WZ \rightarrow l\nu ll$<br>$t\bar{t} \rightarrow bb + ll + X$<br>$\hookrightarrow$ fake $l$ | $ZZ \rightarrow ll ll$<br>$ttZ \rightarrow ll ll + bbX$ | $ZZ \rightarrow ll ll$<br>+ fake lep | $ZZ \rightarrow ll ll$<br>+ 2 fake lep |


| Types of backgrounds      | Suppressed via                  | Bkg. estimation                         |
|---------------------------|---------------------------------|---|
| Fake leptons              | Isolation                       | Reliably extrapolate across isolation   |
| Backgrounds with $b$ jets | $b$ tagging                     | Reliably extrapolate across $b$ tagging |
| Lost leptons              | Removing events with 3rd lepton | Reliably extrapolate across N leptons   |
| Irreducible               | Smart flavor choices            | Reliably extrapolate across flavor      |

Reliably extrapolate across the method used to suppress background to estimate the size of residual backgrounds in signal region

# Fake lepton backgrounds



Pick one lepton events with phase space dominated by QCD (dijet) events

Well isolated  
"fake" lepton  
("tight")



Fake rate =

Less well  
isolated "fake"  
lepton ("Loose")



Fake rate is then applied to signal like region with "Loose"-ly identified leptons


"Side band" in isolation

Underlying effects ( $P_T$  of quarks) that govern fake rate are not measurable  
 $\Rightarrow$  **Source of systematics (~30%)**

Estimate fake lepton by measuring fake rate from QCD events


Devise control regions and extrapolate to signal region

$N_b$  in 3 lepton




Extrapolate across  $N_b$  tag ( $\sim 10\%$ )

4 lepton BDT score  
 $Z \rightarrow ll + e\mu + b$  jets




4 lepton  $m_{T2}$   
 $Z \rightarrow ll + ee/\mu\mu$



Extrapolate across flavor  
(uncertainty  $\sim 5\%$ )

Extrapolate from control region to estimate backgrounds





9 bins

3 bins



7 bins

1 1

More sensitive bins are generally to the right


# 4 lepton event

CMS experiment at the LHC, CERN  
Data recorded: 2016-Jul-23 08:13:27.898048 GMT  
Run 277168, Event No. 3219714497 LS 1799




# 5 lepton event

CMS experiment at the LHC, CERN  
Data recorded: 2016-Oct-09 21:24:05.010240 GMT  
Run 282735, Event No. 989682042 LS 491



| VVV mode   | Significance [ $\sigma$ ] |
|------------|---------------------------|
| <i>WWW</i> | <b>3.3</b> (3.1)          |
| <i>WWZ</i> | <b>3.4</b> (4.1)          |
| <i>WZZ</i> | 1.7 (0.7)                 |
| <i>ZZZ</i> | 0 (0.9)                   |
| Combined   | <b>5.7</b> (5.9)          |



$$\text{Signal strength } \mu = \frac{\text{Measured cross section}}{\text{Theoretical cross section}}$$



- We have **observed** production of three massive gauge boson for the **first** time!
- We also found **evidences** separately for the *WWW* and *WWZ* production.
- The cross sections are compatible with the standard model expectation.

First observation of VVV and evidences for *WWW* and *WWZ* productions

- First observation of  $VVV$  productions was made by CMS collaboration
- Also found evidences for  $WWW$  and  $WWZ$
- first hints for  $WZZ$  production and no hints for  $ZZZ$  yet
- The measured cross section is compatible with SM
- This establishes  $VVV$  process and opens a unique opportunity to test SM
- New physics can be also searched
- LHC will continue to probe electroweak interactions in various  $VVV$  channel

This paper is 1000th paper submitted by CMS!

CERN Courier




*“CMS is the first experiment in the history of high energy physics to reach this outstanding total of papers and with only a fraction of the data that the LHC anticipates to produce in its lifetime. The LHC accelerator at CERN will operate for another two decades.”*



# Backup

# Rare multi-boson processes




N.B.  $x\text{sec} \times \text{decay BR}$  can be even smaller

*Multi-boson processes*

$X = W, Z, H$

Electroweak multi-boson processes are rare and require LHC

# Rare multi-boson processes



Rarer events


N.B. xsec  $\times$  decay BR can be even smaller

*Multi-boson processes*


*X = W, Z, H*

Electroweak multi-boson processes are rare and require LHC







Triboson process has access to studying many multi-*boson* interactions



Triboson process has access to studying many multi-*boson* interactions




Triboson process has access to studying many multi-*boson* interactions



Triboson process has access to studying many multi-*boson* interactions

# Physics of VVV production ( $V = W, Z$ )



Triboson process has access to studying many multi-*boson* interactions

# Kinematic endpoints for 4 leptons





Events are separated into 2 categories by flavor:

- “ $e\mu$  channel”:  $(ee/\mu\mu)_{\text{on-Z-mass}} + e\mu$  (low bkg.)
- “ $ee/\mu\mu$  channel”:  $(ee/\mu\mu)_{\text{on-Z-mass}} + ee/\mu\mu$

$e\mu$  channel utilizes  $m_{T2}$  variable, which is a generalization of  $m_T$  for multiple missing particles.  $m_{T2}$  is sensitive to the end points of  $m_\tau$  from  $ZZ \rightarrow ll\tau\tau$


$ZZ$  bkg in  $ee/\mu\mu$  have low missing energy

Combine these and a few more kinematic variables to form total of 7 signal regions for 4 lepton analysis




Exploit differences between  $Z \rightarrow ll \nu$ .  $WW \rightarrow ll\nu$



# Reconstruct $W \rightarrow qq$ in $WWW \rightarrow l\bar{l}qq$



N.B. some signals are outside the window  
(See next slide)



dijet invariant mass for signal peaks around W mass



Difficult to match  $W \rightarrow qq$   
⇒ Select off-W-mass peak region

Difficult to reconstruct both jets  
⇒ Select 1 jet (1J) events

2 additional categories ( $m_{jj}$ -in,  $m_{jj}$ -out, 1J) each split by  $ee/e\mu/\mu\mu$   
⇒ Total of 9 signal regions for same-sign analysis

We cover wide range of possible jet final states to maximize sensitivity



# Kinematic endpoints for 4 leptons





Events are separated into 2 categories by flavor:

- “ $e\mu$  channel”:  $(ee/\mu\mu)_{\text{on-Z-mass}} + e\mu$  (low bkg.)
- “ $ee/\mu\mu$  channel”:  $(ee/\mu\mu)_{\text{on-Z-mass}} + ee/\mu\mu$

$e\mu$  channel utilizes  $m_{T2}$  variable, which is a generalization of  $m_T$  for multiple missing particles.  $m_{T2}$  is sensitive to the end points of  $m_\tau$  from  $ZZ \rightarrow ll\tau\tau$


$ZZ$  bkg in  $ee/\mu\mu$  have low missing energy

Combine these and a few more kinematic variables to form total of 7 signal regions for 4 lepton analysis



Exploit differences between  $Z \rightarrow ll \nu$ .  $WW \rightarrow ll\nu$

Boosted decision tree is widely used in many analyses at the LHC



[https://arogozhnikov.github.io/2016/07/05/gradient\\_boosting\\_playground.html](https://arogozhnikov.github.io/2016/07/05/gradient_boosting_playground.html)

Train dedicated boosted decision trees to maximize sensitivity

# Overview of BDT

|                | Same-sign   | 3 leptons  | 4 leptons  | 5 leptons  | 6 leptons  |
|----------------|---|--|--|--|--|
| Signals        | $W^\pm \rightarrow l^\pm \nu$<br>$W^\pm \rightarrow l^\pm \nu$<br>$W^\mp \rightarrow qq$                                      | $W \rightarrow l\nu$<br>$W \rightarrow l\nu$<br>$W \rightarrow l\nu$                         | $W \rightarrow l\nu$<br>$W \rightarrow l\nu$<br>$Z \rightarrow ll$ | $W \rightarrow l\nu$<br>$Z \rightarrow ll$<br>$Z \rightarrow ll$ | $Z \rightarrow ll$<br>$Z \rightarrow ll$<br>$Z \rightarrow ll$ |
| Dominant Bkgs. | $WZ \rightarrow l^\pm \nu l^\pm \bar{\nu}$ (lost $l^\pm$ )<br>$t\bar{t} \rightarrow bb + l + X$<br>$\hookrightarrow$ fake $l$ | $WZ \rightarrow l\nu ll$<br>$t\bar{t} \rightarrow bb + ll + X$<br>$\hookrightarrow$ fake $l$ | $ZZ \rightarrow ll ll$<br>$t\bar{t}Z \rightarrow ll ll + bbX$      | $ZZ \rightarrow ll ll$<br>$+ \text{fake lep}$                    | $ZZ \rightarrow ll ll$<br>$+ 2 \text{ fake lep}$               |

“Prompt” bkgs.



“Fake” bkgs.

$t\bar{t}Z$  bkg.

$ZZ$  bkg.

No BDT trained for 5/6 leptons (not enough stats)

Train different BDTs against different backgrounds





Maintained same categorizations but cut on BDT to maximize sensitivity

Total number of bins stayed same (9 for same-sign, 3 for 3 leptons)


Cut on each BDT scores to create a high sensitivity bin

## Quarks



## Leptons


## Forces



- Is it the only Higgs boson? (or are there more?)
- Are multi-*bosons* interactions SM?
- Are there more states involved in electroweak symmetry breaking?
- Is the Higgs potential SM?



Many more to be studied on electroweak sector at the LHC

## Quarks




“massive” bosons

## Forces




## Leptons




- Is it the only Higgs boson? (or are there more?)
- Are multi-*bosons* interactions SM?
- Are there more states involved in electroweak symmetry breaking?
- Is the Higgs potential SM?

Many more to be studied on electroweak sector at the LHC



- Is it the only Higgs boson? (or are there more?)
- Are multi-*bosons* interactions SM?
- Are there more states involved in electroweak symmetry breaking?
- Is the Higgs potential SM?

Many more to be studied on electroweak sector at the LHC



- Is it the only Higgs boson? (or are there more?)
- Are multi-*bosons* interactions SM?
- Are there more states involved in electroweak symmetry breaking?
- Is the Higgs potential SM?

1978 (way) before top/W/Z/Higgs discovery  
Chanowitz, Furman, Hinchliffe

$F, W^\pm, Z$  and  $H$  become “sthenons” in the sense of Appelquist and Bjorken [4]: they couple strongly to one another<sup>#1</sup> but weakly to non-sthenons (i.e., the light particles in the theory).

Many more to be studied on electroweak sector at the LHC




# We need LHC's large and energetic pp collision data

because rare

because "heavy"

Chang  
UCSD



← Multiply by 1000 to get the number of events produced for a picobarn process

During Run 2, CMS recorded 150 fb<sup>-1</sup> of which 137 fb<sup>-1</sup> have been validated as good quality data useable for physics analysis

LHC's large data enables us to study rare EW multi-X processes


Identifying leptons is not enough

We need to further classify the origin


$$\text{Isolation} = \frac{\sum \text{"stuff" in cone } P_T}{P_{T,\text{Lepton}}}$$

N.B. electrons and muons have different effects (muons are cleaner)

non-isolated lepton




isolated lepton



Use isolation to discriminate against leptons from heavy flavor decay


Dubbed "fake lepton"



The more leptons produced the lower the rate (i.e. lower bkg.)

Useful to organize physics analyses by N leptons

dilepton invariant mass of  $Z \rightarrow \ell\ell$  decay



If one selects  $|m_{\ell\ell} - m_Z| > 10 \text{ GeV}$  of  $ee/\mu\mu$  final state  $Z$  is reduced by **an order** of magnitude

If one selects  $e\mu$  final state,  $Z$  is reduced by **2 orders** of magnitude ( $e, \mu$  from  $\tau$  are soft)

$\Rightarrow ZZ$  suppressed in 4 leptons:  $ee/\mu\mu + e\mu$   
 $WZ$  suppressed in  $e^\pm\mu^\mp e^\pm$



↑  
 0 "SFOS"  
 (Zero same-flavor opposite sign pair)

i.e. "ee" or "μμ"

$Z$  decays predominantly to  $ee/\mu\mu \Rightarrow$  select away from  $Z \rightarrow ee/\mu\mu$



# Rejecting events with $b$ jets

EW processes generally do not come with  $b$  jets  $\Rightarrow$  Require # of  $b = 0$



Signals do not have  $b$  jets

EW processes generally do not come with b jets  $\Rightarrow$  Require # of b = 0



Signals do not have *b* jets



## same-sign selection

| Variable                         | $m_{jj}$ -in and $m_{jj}$ -out                                | 1j         |
|----------------------------------|---|------------|
| Trigger                          | Signal triggers, tab. 3.2                                     |            |
| Signal leptons                   | Exactly 2 tight SS leptons with $p_T > 25$ GeV                |            |
| Additional leptons               | No additional very loose lepton                               |            |
| Isolated tracks                  | No additional isolated tracks                                 |            |
| Jets                             | $\geq 2$ jets   | 1 jet      |
| b-tagging                        | no b-tagged jets and soft b-tag objects                       |            |
| $m_{\ell\ell}$                   | $> 20$ GeV  |            |
| $m_{\ell\ell}$                   | $ m_{\ell\ell} - m_Z  > 20$ GeV if $e^\pm e^\pm$              |            |
| $p_T^{\text{miss}}$              | $> 45$ GeV  |            |
| $m_{JJ}$ (leading jets)          | $< 500$ GeV   | —          |
| $\Delta\eta_{JJ}$ (leading jets) | $< 2.5$   | —          |
| $m_{jj}$ (closest $\Delta R$ )   | $65 < m_{jj} < 95$ GeV or<br>$ m_{jj} - 80$ GeV $\geq 15$ GeV | —          |
| $\Delta R_{\ell j}^{\text{min}}$ | —   | $< 1.5$    |
| $m_T^{\text{max}}$               | $> 90$ GeV if not $\mu^\pm \mu^\pm$                           | $> 90$ GeV |

## Three leptons selection

| Variable   | 0 SFOS  | 1 and 2 SFOS |
|--|---|--------------|
| Trigger  | Signal triggers, tab. 3.2   |              |
| Signal leptons   | 3 tight leptons with charge sum = $\pm 1e$<br>$p_T > 25/25/25$ GeV $p_T > 25/20/20$ GeV |              |
| Additional leptons   | No additional very loose lepton   |              |
| $m_{\text{SFOS}}$  | $m_{\text{SFOS}} > 20$ GeV and $ m_{\text{SFOS}} - m_Z  > 20$ GeV                       |              |
| $m_{\ell\ell\ell}$   | $ m_{\ell\ell\ell} - m_Z  > 10$ GeV   |              |
| SF lepton mass   | $> 20$ GeV  | —            |
| Dielectron mass  | $ m_{ee} - m_Z  > 20$ GeV   | —            |
| Jets   | $\leq 1$ jet  | 0 jets       |
| b-tagging  | No b-tagged jets and soft b-tag objects   |              |
| $\Delta\phi(\vec{p}_T(\ell\ell\ell), \vec{p}_T^{\text{miss}})$ | —   | $> 2.5$      |
| $p_T(\ell\ell\ell)$  | —   | $> 50$ GeV   |
| $m_T^{\text{3rd}}$ (1 SFOS) or $m_T^{\text{max}}$ (2 SFOS)     | —   | $> 90$ GeV   |

## Four leptons selection

| Variable                              | $e\mu$ category                                      | $ee/\mu\mu$ category   |
|---------------------------------------|--|--|
| Preselection                          |  | Selections in Table 20   |
| W candidate lepton flavors            | $e\mu$   | $ee/\mu\mu$  |
| $m_{\ell\ell}$                        | Separated into 4 bins in (0, 40, 60, 100, $\infty$ ) | $ m_{\ell\ell} - m_Z  > 10$ GeV  |
| $m_{T2}$                              | $m_{T2} > 25$ GeV (for $m_{\ell\ell} > 100$ GeV)     | ...  |
| $p_{T,4\ell}$ and $p_T^{\text{miss}}$ | ...  | No $p_{T,4\ell}$ cuts and $p_T^{\text{miss}} > 120$ GeV (Bin A)<br>$p_{T,4\ell} > 70$ GeV and $70 < p_T^{\text{miss}} < 120$ GeV (Bin B)<br>$40 < p_{T,4\ell} < 70$ GeV and $70 < p_T^{\text{miss}} < 120$ GeV (Bin C) |

5/6L will be explained later

This is the full selections but I will not go in details for every single one


Lepton finding efficiency is well modeled by MC

(factors:  $P_T$ ,  $\eta$ , lepton ID)

Construct a control region with 3 leptons and extrapolate across 3 lepton  $\rightarrow$  2 leptons

Experimental systematics assigned




Control region data statistics dominates uncertainty (20%)



Estimate lost lepton background by extrapolating across # of leptons




Now that we have established VVV production we can use it to test SM and also search new physics (cf. Four fermion interaction with Fermi constant)



Establishment of VVV production opens up a new physics program

# Fully leptonic v. Semi leptonic channel





┌ If BSM exists, effects are same ─┐

- Physics of  $V \rightarrow ff$  is well understood
- We have now established  $pp \rightarrow VVV$  production in “fully” leptonic decay
- Therefore, there ought to be  $pp \rightarrow VVV \rightarrow$  semi-leptonic  
 $\Rightarrow$  If new physics alters  $pp \rightarrow VVV$ , it will alter fully / semi leptonic the same



$VVV \rightarrow$  semi-leptonic ought to have same physics as  $VVV \rightarrow$  fully leptonic

# Fully leptonic v. Semi leptonic channel



Clean channel for discovery but probing tail is **difficult**

Bkg is larger but distinct high  $P_T$  feature can **discriminate** bkg.



Signal  
Bkg.

Small

Large

Signal  
Bkg.


We can probe VVV  $\rightarrow$  semi-leptonic for new physics

# More multi-massive-X processes for future


listing a few multi-massive-X processes with **same-sign**

$$pp \rightarrow W^\pm W^\pm H$$


Same-sign  
is special



$$pp \rightarrow tW^\pm W^\pm j$$




High  $P_T$  top  
( $> 500$  GeV)




Many more

WWH, tWWj, ttWW, ttZH

$$pp \rightarrow ttWW$$



$$pp \rightarrow ttZH$$




arXiv:1812.09299 Henning, Lombardo, Riemann, Riva  
arXiv:1511.03674 Dror, Farina, Salvioni, Serra  
arXiv:1904.05637 Maltoni, Mantani, Mimasu

There are many more multi-massive-X production to be explored at LHC



## Compact Muon Solenoid LHC, CERN



Visit us: [CMS Public Website](#), [CMS Physics](#) ; Contact us: [CMS Publications Committee](#)


### CMS Publications

|      |                            |   |                    |              |
|------|----------------------------|---|--------------------|--------------|
| 1000 | <a href="#">SMP-19-014</a> | <b>Observation of the production of three massive gauge bosons at <math>\sqrt{s} = 13</math> TeV</b>        | Submitted to PRL   | 19 June 2020 |
| 999  | <a href="#">HIN-19-001</a> | <b>Evidence for top quark production in nucleus-nucleus collisions</b>                                      | Submitted to NP    | 19 June 2020 |
| 998  | <a href="#">TRG-17-001</a> | <b>Performance of the CMS Level-1 trigger in proton-proton collisions at <math>\sqrt{s} = 13</math> TeV</b> | Submitted to JINST | 18 June 2020 |



April 2020

CMS Preliminary



| Quantities  | WWW   | WWZ   | WZZ  | ZZZ  |
|---|-------|-------|------|------|
| $\sigma_{pp \rightarrow VVV \text{ non-VH}} \text{ (fb)}$   | 216.0 | 165.1 | 55.7 | 14.0 |
| $\sigma_{VH \rightarrow VVV} \text{ (fb)}$  | 293.4 | 188.9 | 36.0 | 23.1 |
| $\sigma_{\text{total}} \text{ (fb)}$  | 509.4 | 354.0 | 91.6 | 37.1 |
| $\mathcal{B}_{VVV \rightarrow SS} \text{ (%)}$  | 7.16  | -     | -    | -    |
| $\mathcal{B}_{VVV \rightarrow 3\ell} \text{ (%)}$   | 3.46  | 4.82  | 6.37 | -    |
| $\mathcal{B}_{VVV \rightarrow 4\ell} \text{ (%)}$   | -     | 1.16  | 0.81 | 3.22 |
| $\mathcal{B}_{VVV \rightarrow 5\ell} \text{ (%)}$   | -     | -     | 0.39 | -    |
| $\mathcal{B}_{VVV \rightarrow 6\ell} \text{ (%)}$   | -     | -     | -    | 0.13 |
| $\sigma_{\text{total}} \times \mathcal{B}_{VVV \rightarrow SS} \text{ (fb)}$                                    | 36.4  | -     | -    | -    |
| $\sigma_{\text{total}} \times \mathcal{B}_{VVV \rightarrow 3\ell} \text{ (fb)}$                                 | 17.6  | 17.1  | 5.83 | -    |
| $\sigma_{\text{total}} \times \mathcal{B}_{VVV \rightarrow 4\ell} \text{ (fb)}$                                 | -     | 4.12  | 0.74 | 1.19 |
| $\sigma_{\text{total}} \times \mathcal{B}_{VVV \rightarrow 5\ell} \text{ (fb)}$                                 | -     | -     | 0.36 | -    |
| $\sigma_{\text{total}} \times \mathcal{B}_{VVV \rightarrow 6\ell} \text{ (fb)}$                                 | -     | -     | -    | 0.05 |
| $\sigma_{\text{total}} \times \mathcal{B}_{VVV \rightarrow SS} \times 137 \text{ fb}^{-1} (N_{\text{evts}})$    | 4987  | -     | -    | -    |
| $\sigma_{\text{total}} \times \mathcal{B}_{VVV \rightarrow 3\ell} \times 137 \text{ fb}^{-1} (N_{\text{evts}})$ | 2411  | 2343  | 799  | -    |
| $\sigma_{\text{total}} \times \mathcal{B}_{VVV \rightarrow 4\ell} \times 137 \text{ fb}^{-1} (N_{\text{evts}})$ | -     | 564   | 101  | 163  |
| $\sigma_{\text{total}} \times \mathcal{B}_{VVV \rightarrow 5\ell} \times 137 \text{ fb}^{-1} (N_{\text{evts}})$ | -     | -     | 49.3 | -    |
| $\sigma_{\text{total}} \times \mathcal{B}_{VVV \rightarrow 6\ell} \times 137 \text{ fb}^{-1} (N_{\text{evts}})$ | -     | -     | -    | 6.85 |



| Features                         | Selections  |         |                                     |
|----------------------------------|---|---------|-------------------------------------|
|                                  | SS + $\geq 2j$  | SS + 1j | 3 $\ell$                            |
| Triggers                         | Select events passing dilepton triggers   |         |                                     |
| Number of leptons                | Select events with 2 (3) leptons passing SS-ID (3 $\ell$ -ID) for SS (3 $\ell$ ) final states |         |                                     |
| Number of leptons                | Select events with 2 (3) leptons passing veto-ID for SS (3 $\ell$ ) final states              |         |                                     |
| Isolated tracks                  | No additional isolated tracks   |         | —                                   |
| b-tagging                        | no b-tagged jets and soft b-tag objects   |         |                                     |
| Jets                             | $\geq 2$ jets   | 1 jet   | $\leq 1$ jet                        |
| $m_{JJ}$ (leading jets)          | $< 500$ GeV   |         | —                                   |
| $\Delta\eta_{JJ}$ (leading jets) | $< 2.5$   |         | —                                   |
| $m_{\ell\ell}$                   | $> 20$ GeV  |         | —                                   |
| $m_{\ell\ell}$                   | $ m_{\ell\ell} - m_Z  > 20$ GeV if $e^\pm e^\pm$  |         | —                                   |
| $m_{\text{SFOS}}$                | —   | —       | $m_{\text{SFOS}} > 20$ GeV          |
| $m_{\text{SFOS}}$                | —   | —       | $ m_{\text{SFOS}} - m_Z  > 20$ GeV  |
| $m_{\ell\ell\ell}$               | —   | —       | $ m_{\ell\ell\ell} - m_Z  > 10$ GeV |





| Variable                         | $m_{jj}$ -in and $m_{jj}$ -out                                       | 1j         |
|----------------------------------|--|------------|
| Trigger                          | Signal triggers, tab. 3.2  |            |
| Signal leptons                   | Exactly 2 tight SS leptons with $p_T > 25$ GeV                       |            |
| Additional leptons               | No additional very loose lepton                                      |            |
| Isolated tracks                  | No additional isolated tracks  |            |
| Jets                             | $\geq 2$ jets  | 1 jet      |
| b-tagging                        | no b-tagged jets and soft b-tag objects                              |            |
| $m_{\ell\ell}$                   | $> 20$ GeV   |            |
| $m_{\ell\ell}$                   | $ m_{\ell\ell} - m_Z  > 20$ GeV if $e^\pm e^\pm$                     |            |
| $p_T^{\text{miss}}$              | $> 45$ GeV   |            |
| $m_{JJ}$ (leading jets)          | $< 500$ GeV  | —          |
| $\Delta\eta_{JJ}$ (leading jets) | $< 2.5$  | —          |
| $m_{jj}$ (closest $\Delta R$ )   | $65 < m_{jj} < 95$ GeV or<br>$ m_{jj} - 80 \text{ GeV}  \geq 15$ GeV | —          |
| $\Delta R_{\ell j}^{\text{min}}$ | —  | $< 1.5$    |
| $m_T^{\text{max}}$               | $> 90$ GeV if not $\mu^\pm \mu^\pm$                                  | $> 90$ GeV |



| Variable   | 0 SFOS  | 1 and 2 SFOS         |
|--|---|----------------------|
| Trigger  | Signal triggers, tab. 3.2   |                      |
| Signal leptons   | 3 tight leptons with charge sum = $\pm 1e$                        |                      |
|  | $p_T > 25/25/25$ GeV  | $p_T > 25/20/20$ GeV |
| Additional leptons   | No additional very loose lepton                                   |                      |
| $m_{\text{SFOS}}$  | $m_{\text{SFOS}} > 20$ GeV and $ m_{\text{SFOS}} - m_Z  > 20$ GeV |                      |
| $m_{\ell\ell\ell}$   | $ m_{\ell\ell\ell} - m_Z  > 10$ GeV                               |                      |
| SF lepton mass   | $> 20$ GeV  | —                    |
| Dielectron mass  | $ m_{ee} - m_Z  > 20$ GeV   | —                    |
| Jets   | $\leq 1$ jet  | 0 jets               |
| b-tagging  | No b-tagged jets and soft b-tag objects                           |                      |
| $\Delta\phi(\vec{p}_T(\ell\ell\ell), \vec{p}_T^{\text{miss}})$ | —   | $> 2.5$              |
| $p_T(\ell\ell\ell)$  | —   | $> 50$ GeV           |
| $m_T^{\text{3rd}}$ (1 SFOS) or $m_T^{\text{max}}$ (2 SFOS)     | —   | $> 90$ GeV           |




| Features            | Selections   |
|---------------------|--|
| Number of leptons   | Select events with 4 leptons passing common veto-ID  |
| Triggers            | Select events passing dilepton triggers  |
| Z lepton            | Find opposite charge lepton pairs, passing ZID, closest to $m_Z$<br>Require Z leptons to have $p_T > 25, 15$ GeV |
| W lepton            | Require that leftover leptons are opposite charge and pass WID<br>Require W leptons to have $p_T > 25, 15$ GeV   |
| Low mass resonances | Require any opposite charge pair invariant mass to be greater than 12 GeV  |
| b-tagged jets       | no b-tagged jet  |
| Z mass window       | Require invariant mass of the Z leptons to be within 10 GeV of Z boson mass                                      |




| Variable                              | $e\mu$ category   | $ee/\mu\mu$ category   |
|---------------------------------------|---|--|
| Preselection                          |   | Selections in Table 20   |
| W candidate lepton flavors            | $e\mu$  | $ee/\mu\mu$  |
| $m_{\ell\ell}$                        | Separated into 4 bins in (0, 40, 60, 100, $\infty$ )              | $ m_{\ell\ell} - m_Z  > 10 \text{ GeV}$  |
| $m_{T2}$                              | $m_{T2} > 25 \text{ GeV}$ (for $m_{\ell\ell} > 100 \text{ GeV}$ ) | ...  |
| $p_{T,4\ell}$ and $p_T^{\text{miss}}$ | ...   | No $p_{T,4\ell}$ cuts and $p_T^{\text{miss}} > 120 \text{ GeV}$ (Bin A)<br>$p_{T,4\ell} > 70 \text{ GeV}$ and $70 < p_T^{\text{miss}} < 120 \text{ GeV}$ (Bin B)<br>$40 < p_{T,4\ell} < 70 \text{ GeV}$ and $70 < p_T^{\text{miss}} < 120 \text{ GeV}$ (Bin C) |

$$m_{T2} = \min_{\vec{p}_T^{\nu(1)} + \vec{p}_T^{\nu(2)} = \vec{p}_T^{\text{miss}}} \left[ \max \left( m_T^{(1)}(\vec{p}_T^{\nu(1)}, \vec{p}_T^e), m_T^{(2)}(\vec{p}_T^{\nu(2)}, \vec{p}_T^\mu) \right) \right]$$





For  $WW \rightarrow l\nu l\nu$  sub-system of  $WWZ$ , endpoint is at  $m_W$

For  $Z \rightarrow \tau\tau \rightarrow ll\nu\nu\nu\nu$  sub-system of  $ZZ$ , endpoint is at  $m_\tau$





|                       | ZZ BDT range        | $t\bar{t}Z$ BDT range |
|-----------------------|---------------------|-----------------------|
| $e\mu$ BDT bin 1      | $(-\infty, -0.908)$ | $(-\infty, \infty)$   |
| $e\mu$ BDT bin 2      | $(-0.908, \infty)$  | $(-\infty, 0.015)$    |
| $e\mu$ BDT bin 3      | $(-0.908, 0.733)$   | $(0.015, \infty)$     |
| $e\mu$ BDT bin 4      | $(0.733, \infty)$   | $(0.015, 3.523)$      |
| $e\mu$ BDT bin 5      | $(0.733, \infty)$   | $(3.523, \infty)$     |
| $ee/\mu\mu$ BDT bin A | $(0, 3)$            | -                     |
| $ee/\mu\mu$ BDT bin B | $(3, \infty)$       | -                     |
















| Process | Higgs boson contributions as signal |           | Higgs boson contributions as background |           |
|---------|-------------------------------------|-----------|---|-----------|
|         | sequential-cut                      | BDT-based | sequential-cut                          | BDT-based |
| WWW     | 2.5 (2.9)                           | 3.3 (3.1) | 1.0 (1.8)                               | 1.6 (1.9) |
| WWZ     | 3.5 (3.6)                           | 3.4 (4.1) | 0.9 (2.2)                               | 1.3 (2.2) |
| WZZ     | 1.6 (0.7)                           | 1.7 (0.7) | 1.7 (0.8)                               | 1.7 (0.8) |
| ZZZ     | 0.0 (0.9)                           | 0.0 (0.9) | 0.0 (0.9)                               | 0.0 (0.9) |
| VVV     | 5.0 (5.4)                           | 5.7 (5.9) | 2.3 (3.5)                               | 2.9 (3.5) |



| Process | Higgs boson contributions as signal |                             | Higgs boson contributions as background |                             |
|---------|-------------------------------------|-----------------------------|---|-----------------------------|
|         | sequential-cut                      | BDT-based                   | sequential-cut                          | BDT-based                   |
| WZZ     | 5.2 ( $3.7^{+2.2}_{-1.3}$ )         | 6.1 ( $3.8^{+2.2}_{-1.3}$ ) | 5.8 ( $3.7^{+2.3}_{-1.3}$ )             | 5.8 ( $3.7^{+2.3}_{-1.3}$ ) |
| ZZZ     | 5.4 ( $6.0^{+4.6}_{-2.6}$ )         | 5.4 ( $6.2^{+4.9}_{-2.7}$ ) | 5.6 ( $6.3^{+5.3}_{-2.8}$ )             | 5.7 ( $6.3^{+5.3}_{-2.8}$ ) |



| Signal region                         | SS $m_{jj}$ -in |                 |                   | SS $m_{jj}$ -out |                 |                   | SS 1j         |                 |                   | $3\ell$  |          |          |
|---------------------------------------|-----------------|-----------------|-------------------|------------------|-----------------|-------------------|---------------|-----------------|-------------------|----------|----------|----------|
|                                       | $e^\pm e^\pm$   | $e^\pm \mu^\pm$ | $\mu^\pm \mu^\pm$ | $e^\pm e^\pm$    | $e^\pm \mu^\pm$ | $\mu^\pm \mu^\pm$ | $e^\pm e^\pm$ | $e^\pm \mu^\pm$ | $\mu^\pm \mu^\pm$ | 0SFOS    | 1SFOS    | 2SFOS    |
| Lost/three $\ell$                     | 1.4±0.9         | 5.5±1.6         | 7.0±1.7           | 10.7±2.6         | 9.7±3.6         | 31.4±3.8          | 2.5±1.1       | 41.0±6.1        | 5.8±1.6           | 3.5±0.7  | 25.6±4.2 | 36.1±3.1 |
| Irreducible                           | 1.0±0.1         | 0.6±0.1         | 2.9±0.2           | 4.7±0.4          | 1.9±0.2         | 15.5±1.2          | 0.4±0.0       | 4.6±0.2         | 0.5±0.1           | 1.3±0.1  | 1.2±0.1  | 0.3±0.0  |
| Nonprompt $\ell$                      | 0.6±0.6         | 3.6±2.4         | 4.2±1.5           | 0.8±1.0          | 2.8±1.5         | 9.1±4.5           | 2.5±5.2       | 2.9±1.4         | 0.2±0.1           | 1.8±0.5  | 7.5±2.3  | 1.8±1.1  |
| Charge flips                          | <0.1            | <0.1            | <0.1              | 4.5±2.5          | <0.1            | <0.1              | <0.1          | 0.1±0.1         | <0.1              | <0.1     | 0.8±1.2  | 0.3±0.1  |
| $\gamma \rightarrow$ nonprompt $\ell$ | 0.1±0.2         | 0.1±0.4         | <0.1              | 1.4±0.5          | 1.1±0.4         | 0.7±0.4           | 0.6±1.2       | 4.8±8.0         | <0.1              | <0.1     | 1.0±0.4  | 0.1±1.5  |
| Background sum                        | 3.1±1.1         | 9.8±2.9         | 14.2±2.3          | 22.1±3.8         | 15.6±4.0        | 56.8±6.0          | 6.0±5.4       | 53.5±10.1       | 6.4±1.6           | 6.6±0.9  | 36.2±5.0 | 38.7±3.6 |
| WWW onshell                           | 0.9±0.4         | 2.3±0.9         | 4.6±1.7           | 0.9±0.4          | 1.0±0.6         | 3.3±1.3           | 0.3±0.2       | 1.2±0.4         | 0.4±0.2           | 6.7±2.4  | 4.3±1.6  | 1.8±0.7  |
| WH $\rightarrow$ WWW                  | 0.4±0.3         | 1.3±0.9         | 1.2±0.5           | 0.5±0.3          | 1.3±1.3         | 2.7±1.2           | 1.1±0.8       | 6.5±3.1         | 2.2±1.1           | 3.4±1.6  | 5.0±2.1  | 0.6±0.6  |
| WWW total                             | 1.3±0.5         | 3.7±1.3         | 5.8±1.7           | 1.5±0.5          | 2.3±1.4         | 6.0±1.7           | 1.4±0.8       | 7.7±3.1         | 2.5±1.1           | 10.1±2.9 | 9.3±2.6  | 2.4±0.9  |
| WWZ onshell                           | <0.1            | <0.1            | <0.1              | <0.1             | <0.1            | <0.1              | <0.1          | <0.1            | <0.1              | 0.2±0.1  | <0.1     | <0.1     |
| ZH $\rightarrow$ WWZ                  | <0.1            | <0.1            | <0.1              | <0.1             | <0.1            | <0.1              | <0.1          | <0.1            | <0.1              | 0.1±0.1  | 0.1±0.1  | <0.1     |
| WWZ total                             | <0.1            | <0.1            | <0.1              | <0.1             | <0.1            | <0.1              | <0.1          | <0.1            | <0.1              | 0.3±0.1  | 0.1±0.1  | <0.1     |
| WZZ onshell                           | <0.1            | <0.1            | <0.1              | <0.1             | <0.1            | <0.1              | <0.1          | <0.1            | <0.1              | <0.1     | <0.1     | <0.1     |
| WH $\rightarrow$ WZZ                  | <0.1            | <0.1            | <0.1              | <0.1             | <0.1            | <0.1              | <0.1          | <0.1            | <0.1              | <0.1     | <0.1     | <0.1     |
| WZZ total                             | <0.1            | <0.1            | <0.1              | <0.1             | <0.1            | <0.1              | <0.1          | <0.1            | <0.1              | <0.1     | <0.1     | <0.1     |
| ZZZ onshell                           | <0.1            | <0.1            | <0.1              | <0.1             | <0.1            | <0.1              | <0.1          | <0.1            | <0.1              | <0.1     | <0.1     | <0.1     |
| ZH $\rightarrow$ ZZZ                  | <0.1            | <0.1            | <0.1              | <0.1             | <0.1            | <0.1              | <0.1          | <0.1            | <0.1              | <0.1     | <0.1     | <0.1     |
| ZZZ total                             | <0.1            | <0.1            | <0.1              | <0.1             | <0.1            | <0.1              | <0.1          | <0.1            | <0.1              | <0.1     | <0.1     | <0.1     |
| VVV onshell                           | 0.9±0.4         | 2.3±0.9         | 4.6±1.7           | 0.9±0.4          | 1.0±0.6         | 3.3±1.3           | 0.3±0.2       | 1.2±0.4         | 0.4±0.2           | 6.9±2.4  | 4.3±1.6  | 1.8±0.7  |
| VH $\rightarrow$ VVV                  | 0.4±0.3         | 1.3±0.9         | 1.2±0.5           | 0.5±0.3          | 1.3±1.3         | 2.7±1.2           | 1.1±0.8       | 6.5±3.1         | 2.2±1.1           | 3.6±1.6  | 5.1±2.1  | 0.6±0.6  |
| VVV total                             | 1.3±0.5         | 3.7±1.3         | 5.8±1.7           | 1.5±0.5          | 2.3±1.4         | 6.0±1.7           | 1.4±0.8       | 7.7±3.1         | 2.5±1.1           | 10.4±2.9 | 9.3±2.6  | 2.4±0.9  |
| Total                                 | 4.4±1.2         | 13.5±3.2        | 20.0±2.9          | 23.6±3.8         | 17.8±4.2        | 62.7±6.3          | 7.4±5.5       | 61.2±10.6       | 9.0±2.0           | 17.0±3.0 | 45.5±5.6 | 41.1±3.7 |
| Observed                              | 3               | 14              | 15                | 22               | 22              | 67                | 13            | 69              | 8                 | 17       | 42       | 39       |

| Signal region  | $4\ell e\mu$ |         |         |         |         | $4\ell ee/\mu\mu$ |          | $5\ell$   | $6\ell$   |
|----------------|--------------|---------|---------|---------|---------|-------------------|----------|-----------|-----------|
|                | bin 1        | bin 2   | bin 3   | bin 4   | bin 5   | bin A             | bin B    |           |           |
| ZZ             | 15.9±1.0     | 1.6±0.1 | 0.6±0.1 | 0.6±0.1 | 0.2±0.0 | 76.4±4.3          | 2.9±0.3  | 0.30±0.09 | 0.01±0.01 |
| t $\bar{t}$ Z  | 0.2±0.1      | 0.1±0.1 | 2.8±0.5 | 1.4±0.2 | 0.1±0.1 | 1.5±0.3           | 2.3±0.3  | <0.01     | <0.01     |
| tWZ            | 0.1±0.1      | 0.1±0.1 | 0.6±0.1 | 0.7±0.1 | 0.1±0.1 | 0.5±0.1           | 0.7±0.1  | <0.01     | <0.01     |
| WZ             | 0.5±0.2      | 0.2±0.2 | 0.5±0.2 | 0.3±0.3 | 0.1±0.1 | 1.0±0.4           | 0.2±0.1  | <0.01     | <0.01     |
| Other          | 1.1±0.4      | 0.5±0.5 | 0.5±0.2 | 0.6±0.2 | <0.1    | 2.7±0.6           | 0.5±0.2  | <0.01     | <0.01     |
| Background sum | 17.8±1.1     | 2.5±0.5 | 5.0±0.6 | 3.6±0.4 | 0.5±0.1 | 82.2±4.3          | 6.6±0.5  | 0.30±0.09 | 0.01±0.01 |
| WWW onshell    | <0.1         | <0.1    | <0.1    | <0.1    | <0.1    | <0.1              | <0.1     | <0.01     | <0.01     |
| WH → WWW       | <0.1         | <0.1    | <0.1    | <0.1    | <0.1    | <0.1              | <0.1     | <0.01     | <0.01     |
| WWW total      | <0.1         | <0.1    | <0.1    | <0.1    | <0.1    | <0.1              | <0.1     | <0.01     | <0.01     |
| WWZ onshell    | 0.3±0.1      | 0.4±0.2 | 1.4±0.7 | 3.6±1.5 | 1.0±0.5 | 2.7±1.2           | 3.2±1.4  | <0.01     | <0.01     |
| ZH → WWZ       | 1.1±0.5      | 1.1±0.5 | 0.5±0.2 | 1.3±0.5 | 1.8±0.8 | 2.9±1.2           | 1.5±0.6  | <0.01     | <0.01     |
| WWZ total      | 1.3±0.5      | 1.5±0.5 | 1.9±0.8 | 4.9±1.6 | 2.9±0.9 | 5.6±1.7           | 4.7±1.5  | <0.01     | <0.01     |
| WZZ onshell    | 0.2±0.2      | 0.1±0.1 | 0.2±0.2 | 0.4±0.4 | 0.1±0.1 | 0.5±0.4           | 0.2±0.2  | 2.62±1.82 | 0.03±0.05 |
| WH → WZZ       | 0.2±0.3      | 0.2±0.3 | <0.1    | 0.5±0.5 | <0.1    | <0.1              | <0.1     | <0.01     | <0.01     |
| WZZ total      | 0.4±0.3      | 0.3±0.3 | 0.2±0.2 | 0.9±0.7 | 0.1±0.1 | 0.5±0.4           | 0.2±0.2  | 2.62±1.82 | 0.03±0.05 |
| ZZZ onshell    | <0.1         | <0.1    | <0.1    | <0.1    | <0.1    | <0.1              | <0.1     | <0.01     | <0.01     |
| ZH → ZZZ       | <0.1         | <0.1    | <0.1    | <0.1    | <0.1    | <0.1              | <0.1     | <0.01     | <0.01     |
| ZZZ total      | <0.1         | <0.1    | <0.1    | <0.1    | <0.1    | <0.1              | <0.1     | <0.01     | <0.01     |
| VVV onshell    | 0.5±0.2      | 0.4±0.2 | 1.6±0.8 | 4.0±1.5 | 1.1±0.5 | 3.2±1.3           | 3.4±1.4  | 2.62±1.82 | 0.03±0.05 |
| VH → VVV       | 1.2±0.5      | 1.3±0.6 | 0.5±0.2 | 1.7±0.8 | 1.8±0.8 | 2.9±1.2           | 1.5±0.6  | <0.01     | <0.01     |
| VVV total      | 1.7±0.6      | 1.7±0.6 | 2.1±0.8 | 5.8±1.7 | 3.0±0.9 | 6.1±1.8           | 4.8±1.5  | 2.62±1.82 | 0.03±0.05 |
| Total          | 19.5±1.2     | 4.2±0.8 | 7.1±1.0 | 9.4±1.8 | 3.5±0.9 | 88.2±4.7          | 11.4±1.6 | 2.92±1.82 | 0.04±0.05 |
| Observed       | 22           | 9       | 7       | 8       | 3       | 80                | 11       | 3         | 0         |





| Signal region                         | SS $m_{jj}$ -in |                 |                   | SS $m_{jj}$ -out |                 |                   | SS 1j         |                 |                   | $3\ell$  |          |          |
|---------------------------------------|-----------------|-----------------|-------------------|------------------|-----------------|-------------------|---------------|-----------------|-------------------|----------|----------|----------|
|                                       | $e^\pm e^\pm$   | $e^\pm \mu^\pm$ | $\mu^\pm \mu^\pm$ | $e^\pm e^\pm$    | $e^\pm \mu^\pm$ | $\mu^\pm \mu^\pm$ | $e^\pm e^\pm$ | $e^\pm \mu^\pm$ | $\mu^\pm \mu^\pm$ | 0 SFOS   | 1 SFOS   | 2 SFOS   |
| Lost/three $\ell$                     | 1.8±0.4         | 10.9±2.0        | 8.7±1.0           | 8.8±1.7          | 46.0±6.2        | 44.8±4.4          | 8.4±1.3       | 43.5±4.4        | 34.5±2.7          | 4.6±0.8  | 15.1±1.5 | 58.3±2.4 |
| Irreducible                           | 2.1±0.4         | 13.0±3.6        | 8.4±1.4           | 9.8±1.4          | 41.1±4.5        | 42.8±4.7          | 2.6±0.6       | 22.8±8.6        | 13.2±1.9          | 2.5±0.9  | 2.2±1.2  | 2.5±0.8  |
| Nonprompt $\ell$                      | 1.3±0.9         | 5.8±2.4         | 6.8±2.2           | 2.3±1.3          | 12.0±6.1        | 11.2±3.8          | 1.8±2.9       | 2.4±1.3         | 2.8±1.1           | 3.0±0.9  | 5.7±1.6  | 5.9±1.6  |
| Charge flips                          | <0.1            | 1.2±2.0         | <0.1              | 2.6±1.6          | 1.0±0.5         | <0.1              | 6.9±4.7       | 0.2±0.1         | <0.1              | <0.1     | 1.1±1.3  | 0.7±0.2  |
| $\gamma \rightarrow$ nonprompt $\ell$ | 1.4±0.4         | 2.3±0.9         | 0.1±0.8           | 8.6±3.1          | 19.2±5.1        | 2.3±0.9           | 3.8±1.1       | 19.7±6.0        | 13.8±7.0          | <0.1     | 0.6±0.7  | 0.2±0.3  |
| Background sum                        | 6.7±1.2         | 33.3±5.2        | 24.0±2.9          | 32.1±4.3         | 119±11          | 101±8             | 23.6±5.8      | 88.7±11.4       | 64.4±7.8          | 10.1±1.5 | 24.7±2.9 | 67.6±3.1 |
| WWW onshell                           | 1.0±0.5         | 3.3±1.5         | 3.5±1.6           | 0.9±0.5          | 3.9±1.8         | 4.1±1.9           | 0.5±0.3       | 1.8±0.8         | 1.7±0.9           | 5.9±2.6  | 3.8±1.7  | 2.5±1.2  |
| WH $\rightarrow$ WWW                  | 0.2±0.3         | 1.9±1.5         | 0.6±0.4           | 0.4±0.4          | 1.3±0.8         | 1.7±1.0           | 0.8±0.5       | 4.5±2.7         | 3.3±2.0           | 3.0±1.7  | 2.7±1.5  | 1.3±0.8  |
| WWW total                             | 1.2±0.6         | 5.1±2.2         | 4.1±1.6           | 1.3±0.6          | 5.3±2.0         | 5.7±2.1           | 1.4±0.6       | 6.3±2.8         | 5.0±2.2           | 8.8±3.1  | 6.6±2.3  | 3.8±1.4  |
| WWZ onshell                           | 0.1±0.1         | 0.3±0.2         | 0.2±0.1           | <0.1             | <0.1            | 0.1±0.1           | 0.1±0.1       | <0.1            | <0.1              | 0.3±0.2  | 0.2±0.2  | 0.2±0.1  |
| ZH $\rightarrow$ WWZ                  | 0.1±0.1         | <0.1            | <0.1              | <0.1             | <0.1            | 0.3±0.3           | <0.1          | <0.1            | 0.4±0.4           | 0.2±0.1  | <0.1     | <0.1     |
| WWZ total                             | 0.1±0.2         | 0.3±0.2         | 0.2±0.1           | <0.1             | <0.1            | 0.4±0.3           | 0.1±0.1       | <0.1            | 0.4±0.4           | 0.4±0.2  | 0.2±0.2  | 0.2±0.1  |
| WZZ onshell                           | <0.1            | <0.1            | <0.1              | <0.1             | <0.1            | <0.1              | <0.1          | <0.1            | <0.1              | <0.1     | <0.1     | <0.1     |
| WH $\rightarrow$ WZZ                  | <0.1            | <0.1            | <0.1              | <0.1             | <0.1            | <0.1              | <0.1          | <0.1            | <0.1              | <0.1     | <0.1     | <0.1     |
| WZZ total                             | <0.1            | <0.1            | <0.1              | <0.1             | <0.1            | <0.1              | <0.1          | <0.1            | <0.1              | <0.1     | <0.1     | <0.1     |
| ZZZ onshell                           | <0.1            | <0.1            | <0.1              | <0.1             | <0.1            | <0.1              | <0.1          | <0.1            | <0.1              | <0.1     | <0.1     | <0.1     |
| ZH $\rightarrow$ ZZZ                  | <0.1            | <0.1            | <0.1              | <0.1             | <0.1            | <0.1              | <0.1          | <0.1            | <0.1              | <0.1     | <0.1     | <0.1     |
| ZZZ total                             | <0.1            | <0.1            | <0.1              | <0.1             | <0.1            | <0.1              | <0.1          | <0.1            | <0.1              | <0.1     | <0.1     | <0.1     |
| VVV onshell                           | 1.0±0.5         | 3.5±1.5         | 3.7±1.6           | 0.9±0.5          | 3.9±1.8         | 4.2±1.9           | 0.6±0.3       | 1.8±0.8         | 1.7±0.9           | 6.1±2.6  | 4.0±1.8  | 2.7±1.2  |
| VH $\rightarrow$ VVV                  | 0.3±0.3         | 1.9±1.5         | 0.6±0.4           | 0.4±0.4          | 1.3±0.8         | 2.0±1.0           | 0.8±0.5       | 4.5±2.7         | 3.7±2.0           | 3.1±1.7  | 2.7±1.5  | 1.3±0.8  |
| VVV total                             | 1.3±0.6         | 5.4±2.2         | 4.2±1.6           | 1.3±0.6          | 5.3±2.0         | 6.1±2.1           | 1.4±0.6       | 6.3±2.8         | 5.4±2.2           | 9.3±3.1  | 6.8±2.3  | 3.9±1.4  |
| Total                                 | 8.0±1.3         | 38.7±5.6        | 28.2±3.4          | 33.5±4.4         | 125±11          | 107±8             | 25.0±5.8      | 95.0±11.8       | 69.8±8.1          | 19.4±3.4 | 31.4±3.7 | 71.5±3.4 |
| Observed                              | 5               | 46              | 20                | 31               | 112             | 118               | 29            | 101             | 69                | 20       | 32       | 69       |





| Signal region  | $4\ell e\mu$ |         |         |         | $4\ell ee/\mu\mu$ |          |         | $5\ell$   | $6\ell$   |
|----------------|--------------|---------|---------|---------|-------------------|----------|---------|-----------|-----------|
|                | bin 4        | bin 3   | bin 2   | bin 1   | bin A             | bin B    | bin C   |           |           |
| ZZ             | 0.3±0.0      | 0.7±0.0 | 0.7±0.0 | 0.4±0.0 | 1.8±0.2           | 6.0±0.6  | 5.0±0.5 | 0.30±0.08 | 0.01±0.01 |
| t $\bar{t}$ Z  | 0.2±0.0      | 0.3±0.1 | 0.8±0.1 | 2.3±0.4 | 1.4±0.2           | 1.1±0.2  | 0.2±0.0 | <0.01     | <0.01     |
| tWZ            | 0.1±0.1      | 0.1±0.1 | 0.3±0.0 | 0.8±0.1 | 0.5±0.1           | 0.3±0.1  | 0.1±0.1 | <0.01     | <0.01     |
| WZ             | 0.2±0.1      | 0.1±0.1 | 0.1±0.2 | 0.6±0.2 | <0.1              | 0.2±0.1  | 0.1±0.1 | <0.01     | <0.01     |
| Other          | <0.1         | 0.2±0.1 | 0.6±0.3 | 0.2±0.1 | <0.1              | 1.4±0.5  | 0.1±0.1 | <0.01     | <0.01     |
| Background sum | 0.8±0.1      | 1.4±0.1 | 2.5±0.3 | 4.3±0.4 | 3.7±1.9           | 9.1±0.8  | 5.5±0.5 | 0.30±0.08 | 0.01±0.01 |
| WWW onshell    | <0.1         | <0.1    | <0.1    | <0.1    | <0.1              | <0.1     | <0.1    | <0.01     | <0.01     |
| WH → WWW       | <0.1         | <0.1    | <0.1    | <0.1    | <0.1              | <0.1     | <0.1    | <0.01     | <0.01     |
| WWW total      | <0.1         | <0.1    | <0.1    | <0.1    | <0.1              | <0.1     | <0.1    | <0.01     | <0.01     |
| WWZ onshell    | 0.5±0.2      | 0.5±0.2 | 1.1±0.4 | 4.0±1.6 | 2.1±0.9           | 1.2±0.4  | 0.6±0.2 | <0.01     | <0.01     |
| ZH → WWZ       | 2.3±0.9      | 1.1±0.4 | 0.3±0.1 | 0.1±0.1 | 0.8±0.3           | 0.9±0.4  | 0.5±0.2 | <0.01     | <0.01     |
| WWZ total      | 2.8±0.9      | 1.6±0.5 | 1.4±0.4 | 4.1±1.6 | 2.9±1.0           | 2.1±0.6  | 1.1±0.3 | <0.01     | <0.01     |
| WZZ onshell    | <0.1         | 0.1±0.1 | 0.1±0.1 | 0.4±0.3 | 0.2±0.2           | 0.1±0.1  | 0.1±0.1 | 2.17±1.46 | 0.03±0.04 |
| WH → WZZ       | <0.1         | 0.4±0.3 | 0.1±0.2 | <0.1    | <0.1              | <0.1     | <0.1    | <0.01     | <0.01     |
| WZZ total      | <0.1         | 0.4±0.4 | 0.2±0.2 | 0.4±0.3 | 0.2±0.2           | 0.1±0.1  | 0.1±0.1 | 2.17±1.46 | 0.03±0.04 |
| ZZZ onshell    | <0.1         | <0.1    | <0.1    | <0.1    | <0.1              | <0.1     | <0.1    | <0.01     | <0.01     |
| ZH → ZZZ       | <0.1         | <0.1    | <0.1    | <0.1    | <0.1              | <0.1     | <0.1    | <0.01     | <0.01     |
| ZZZ total      | <0.1         | <0.1    | <0.1    | <0.1    | <0.1              | <0.1     | <0.1    | <0.01     | <0.01     |
| VVV onshell    | 0.5±0.2      | 0.6±0.2 | 1.2±0.4 | 4.4±1.6 | 2.3±0.9           | 1.3±0.5  | 0.7±0.2 | 2.17±1.46 | 0.03±0.04 |
| VH → VVV       | 2.3±0.9      | 1.5±0.5 | 0.4±0.3 | 0.1±0.1 | 0.8±0.3           | 0.9±0.4  | 0.5±0.2 | <0.01     | <0.01     |
| VVV total      | 2.8±0.9      | 2.1±0.6 | 1.6±0.5 | 4.5±1.6 | 3.1±1.0           | 2.2±0.6  | 1.2±0.3 | 2.17±1.46 | 0.03±0.04 |
| Total          | 3.6±0.9      | 3.5±0.6 | 4.1±0.6 | 8.8±1.7 | 6.8±2.1           | 11.3±1.0 | 6.6±0.6 | 2.47±1.46 | 0.04±0.04 |
| Observed       | 7            | 1       | 5       | 7       | 6                 | 8        | 7       | 3         | 0         |


# History lesson



...after analysis of Run I data, ... **2**  $m_W$  shifted a full s.d. ... the  $m_{Higgs}$  must be **3** much lower than anyone had anticipated. ... Surprises happen.


– D. Amidei, R. Brock Fermi news 1/17/2003

History tells us with more data we get smarter; also surprises happen




ment with the results obtained from simulation. The  $\sigma(p_T)/p_T$  averaged over  $\phi$  and  $\eta$  varies in  $p_T$  from  $(1.8 \pm 0.3(\text{stat.}))\%$  at  $p_T = 30 \text{ GeV}/c$  to  $(2.3 \pm 0.3(\text{stat.}))\%$  at  $p_T = 50 \text{ GeV}/c$ , again in good agreement with the expectations from simulation.

<https://arxiv.org/pdf/1206.4071.pdf>



(a)



(b)



arXiv.org > physics > arXiv:1502.02701

Search...

Help | Advanced


Physics > Instrumentation and Detectors

*[Submitted on 9 Feb 2015 (v1), last revised 1 Jul 2015 (this version, v2)]*

## Performance of electron reconstruction and selection with the CMS detector in proton–proton collisions at $\sqrt{s} = 8$ TeV

[CMS Collaboration](#)

The performance and strategies used in electron reconstruction and selection at CMS are presented based on data corresponding to an integrated luminosity of 19.7 inverse femtobarns, collected in proton–proton collisions at  $\sqrt{s} = 8$  TeV at the CERN LHC. The paper focuses on prompt isolated electrons with transverse momenta ranging from about 5 to a few 100 GeV. A detailed description is given of the algorithms used to cluster energy in the electromagnetic calorimeter and to reconstruct electron trajectories in the tracker. The electron momentum is estimated by combining the energy measurement in the calorimeter with the momentum measurement in the tracker. Benchmark selection criteria are presented, and their performances assessed using Z, Upsilon, and J/psi decays into electron–positron pairs. The spectra of the observables relevant to electron reconstruction and selection as well as their global efficiencies are well reproduced by Monte Carlo simulations. The momentum scale is calibrated with an uncertainty smaller than 0.3%. The momentum resolution for electrons produced in Z boson decays ranges from 1.7 to 4.5%, depending on electron pseudorapidity and energy loss through bremsstrahlung in the detector material.





<https://twiki.cern.ch/twiki/pub/CMSPublic/BTV13TeV2017FIRST2018/PT30GeV.pdf>

