

# Paleo-Detectors for Galactic Supernova Neutrinos

*Friday, 31 July 2020 08:30 (15 minutes)*

Paleo-detectors are a proposed experimental technique in which one would search for traces of recoiling nuclei in ancient minerals. Natural minerals on Earth are as old as  $\mathcal{O}(1)$  Gyr and, in many minerals, the damage tracks left by recoiling nuclei are also preserved for time scales long compared to 1 Gyr once created. Thus, even reading out relatively small target samples of order 100 g, paleo-detectors would allow one to search for very rare events thanks to the large exposure,  $\varepsilon \sim 100 \text{ g Gyr} = 10^5 \text{ t yr}$ . Here, we explore the potential of paleo-detectors to measure nuclear recoils induced by neutrinos from galactic core collapse supernovae. We find that they would not only allow for a direct measurement of the average core collapse supernova rate in the Milky Way, but would also contain information about the time-dependence of the local supernova rate over the past  $\sim 1$  Gyr. Since the supernova rate is thought to be directly proportional to the star formation rate, such a measurement would provide a determination of the local star formation history. We investigate the sensitivity of paleo-detectors to both a smooth time evolution and an enhancement of the core collapse supernova rate on relatively short time scales, as would be expected for a starburst period in the local group.

## Secondary track (number)

**Primary author:** Dr STENGEL, Patrick (Stockholm University)

**Presenter:** Dr STENGEL, Patrick (Stockholm University)

**Session Classification:** Neutrino Physics

**Track Classification:** 02. Neutrino Physics