

A combined explanation of the B -decay anomalies with a single vector leptoquark

Friday, 31 July 2020 13:48 (3 minutes)

In order to simultaneously account for both $R_{D^{(*)}}$ and $R_{K^{(*)}}$ anomalies in B -decays, we consider an extension of the Standard Model by a single vector leptoquark field, and study how one can achieve the required lepton flavour non-universality, starting from a priori universal gauge couplings. While the unitary quark-lepton mixing induced by $SU(2)_L$ breaking is insufficient, we find that effectively nonunitary mixings hold the key to simultaneously address the $R_{K^{(*)}}$ and $R_{D^{(*)}}$ anomalies. As an intermediate step towards various UV-complete models, we show that the mixings of charged leptons with additional vector-like heavy leptons successfully provide a nonunitary framework to explain $R_{K^{(*)}}$ and $R_{D^{(*)}}$. These realisations have a strong impact for electroweak precision observables and for flavour violating ones: isosinglet heavy lepton realisations are already excluded due to excessive contributions to lepton flavour violating Z -decays. Furthermore, in the near future, the expected progress in the sensitivity of charged lepton flavour violation experiments should allow to fully probe this class of vector leptoquark models.

Secondary track (number)

05.

Primary author: KRIEWALD, Jonathan (LPC Clermont)

Co-authors: HATI, Chandan (LPC Clermont); TEIXEIRA, Ana M. (LPC Clermont); ORLOFF, Jean (Université Blaise Pascal - Clermont-Ferrand)

Presenter: KRIEWALD, Jonathan (LPC Clermont)

Session Classification: Beyond the Standard Model - Posters

Track Classification: 03. Beyond the Standard Model