

A search for dark matter using sub-PeV gamma-rays observed by Tibet AS _{γ}


Tarak Nath Maity

Based on

TNM, A K Saha, A Dubey, R Laha

2105.05680

CHEP, IISc Bangalore, India





भारतीय विज्ञान संस्थान

The XXVIII International Conference on Supersymmetry and Unification of Fundamental Interactions

(SUSY 2021)

Tibet AS+MD



- ✓ Area: ~1.5 Eiffel tower
- ✓ Hybridize with muon detector.
- ✓ 2.4 m underground
- ✓ Muon with energy greater than 1 GeV

Livetime: 719 days from February 2014 to May 2017



©Tibet AS_γ Collaboration

Muon detector: gamma and cosmic ray (CR) discrimination


What is it observing?

- ✓ Are not deflected by interstellar magnetic fields.
- ✓ Observation of ~ 100 TeV gamma ray predict the Galactic origin of the PeV cosmic ray.

CR Source ?




How? Photon and Proton Shower




© Google

Occasional γ -p interaction gives rises shower similar to hadronic shower

How? Photon and Proton Shower





200TeV Gamma-ray induced AS



200TeV Proton induced AS

- ✓ After muon cut, $\sim 99.9\%$ CR rejection & $\sim 90\%$ gamma efficiency @100 TeV


Result: Observed Flux



- ✓ First detection of sub-PeV diffuse gamma rays.
- ✓ Space dependent and space independent cosmic ray model seems to fit well with data, proposed in 1804.10116
- ✓ Several recent proposals e.g., see 2104.09491, 2104.03729, 2104.05609

What are we doing?

Observed Flux: Whether this observation could be used for detection of dark matter?



Decaying DM: gamma-ray spectrum


$$\frac{dN_\gamma}{dE_\gamma}$$

PPPC

1012.4515

HDMSpectra

2007.15001



© C Weniger

Decaying DM + Background < Data

DM Flux

$$\frac{d^2 \phi_\gamma}{dE_\gamma d\Omega}(E_\gamma) = \frac{1}{\Delta\Omega} \int_{\Delta\Omega} d\Omega \frac{1}{4\pi m_\chi \tau_\chi} \frac{dN_\gamma}{dE_\gamma}(E_\gamma)$$

$$\int_0^{s_{\max}} \rho_\chi(s, b, l) e^{-\tau_{\gamma\gamma}(E_\gamma, s, b, l)} ds$$

NFW \leftarrow $\rho_\chi(s, b, l)$ \rightarrow Attenuation $e^{-\tau_{\gamma\gamma}(E_\gamma, s, b, l)}$

Background

Different cosmic ray models

- Space dependent CR, 1804.10116
- Space independent CR, 1804.10116
- Hybrid gamma-model, 2104.09491

Data


TABLE S2. Galactic diffuse gamma-ray fluxes measured by the Tibet AS+MD array.

Energy bin (TeV)	Representative E (TeV)	Flux ($25^\circ < l < 100^\circ, b < 5^\circ$) ($\text{TeV}^{-1} \text{cm}^{-2} \text{s}^{-1} \text{sr}^{-1}$)	Flux ($50^\circ < l < 200^\circ, b < 5^\circ$) ($\text{TeV}^{-1} \text{cm}^{-2} \text{s}^{-1} \text{sr}^{-1}$)
100 – 158	121	$(3.16 \pm 0.64) \times 10^{-15}$	$(1.69 \pm 0.41) \times 10^{-15}$
158 – 398	220	$(3.88 \pm 1.00) \times 10^{-16}$	$(2.27 \pm 0.60) \times 10^{-16}$
398 – 1000	534	$(6.86^{+3.30}_{-2.40}) \times 10^{-17}$	$(2.99^{+1.40}_{-1.02}) \times 10^{-17}$

Amenomori et al 2104.05181 PRL


Decaying DM: Limits

✓ We have done a χ^2 analysis to set the limits.



*TNM, Saha, Dubey,
Laha 2105.05680*

Decaying DM Limits: DM profile



@ Our limits are robust.

Conclusion


✓ Recently, Tibet AS γ collaboration has discovered the first sub-PeV diffuse gamma-rays from the MW Galactic disk.

✓ Data broadly agrees with prior theoretical expectations

✓ We study the impact of this discovery on PeV scale decaying DM


✓ We find that data provides strongest bound on most of the final states

➔ Near future data of these high-energy gamma-rays can be used to discover heavy decaying DM.



email: tarak.maity.physics@gmail.com



Photon Proton Shower: Tibet AS $_{\gamma}$




	$N_{\mu} < 75$	$N_{\mu} < 100$	$N_{\mu} < 200$	$N_{\mu} < 300$
Percentage of γ -ray signals retained	10%	20%	60%	83%
Level of cosmic-ray background				
Solid line fit	10^{-5}	1.5×10^{-5}	4×10^{-5}	10^{-4}
Dashed line fit	$< 10^{-7}$	10^{-7}	6.6×10^{-7}	4×10^{-6}

Gaisser et al PRD 91'

- ✓ After muon cut, $\sim 99.9\%$ CR rejection & $\sim 90\%$ gamma efficiency @100 TeV



Tibet AS_γ



Decaying DM: Attenuation


Pair production: $\gamma + \gamma_b \rightarrow e^+ e^-$

γ_b $\left\{ \begin{array}{l} \text{CMB} \\ \text{Starlight} \\ \text{Infrared} \end{array} \right.$


Attenuation $\sim e^{(-L/\lambda)}$

Mean free path $\lambda = 1/n_b \sigma_{\gamma\gamma}$

CMB





SL+IR



@ A 100 TeV photon must originate from our galaxy.

Decaying DM: Limits



- ✓ For some of the channels (e.g., $t\bar{t}$) our bounds are stronger than previous limits.
- ✓ For some of the channels (e.g., $\nu\bar{\nu}$) our bounds are not as strong as previous limits.