

Enhanced Long-Lived Dark Photon signal @ LHC

Mingxuan Du

Nanjing University



In collaboration with Zuowei Liu, Van Que Tran,
Rundong Fang

arXiv: 1912.00422 2108.XXXXX


SUSY21 2021.08.25

Outline

- Motivation
- Our long-lived dark photon model (LLDP)
- Experimental constraints
- Enhanced LLDP signal @ timing detectors
- Enhanced LLDP signal @ LHCb
- Far detectors
- Conclusion

Motivation

Long-lived particles (LLP) present in a lot of BSM models




distance travelled by long-lived dark photon (LLDP)

$$d = \gamma v \tau \simeq 1 \text{ meter} \left[\frac{10^{-6}}{g} \right]^2 \left[\frac{E_{A'}}{100 \text{ GeV}} \right] \left[\frac{\text{GeV}}{M_{A'}} \right]^2$$

Suppressed ? Boosted Collider

To enhance LLDP signals@LHC

Make the LLDP production process different with its decay process



If $\epsilon_2 \gg \epsilon_1 \sim 10^{-6}$, A' is LLDP and its signal can be enhanced

Stueckelberg w/ 2 U(1) extension

- SM extended by a hidden sector (HS) with **two** U(1) gauge bosons X and C

$$-4\mathcal{L}_F = X_{\mu\nu}^2 + 2(\partial_\mu\sigma_1 + m_1\epsilon_1 B_\mu + m_1 X_\mu)^2$$

$$-4\mathcal{L}_W = C_{\mu\nu}^2 + 2(\partial_\mu\sigma_2 + m_2\epsilon_2 B_\mu + m_2 C_\mu)^2$$

SM $U(1)_Y$

Hidden $U(1)_F$

Hidden $U(1)_W$

- Both 2 extra gauge bosons obtain mass via **Stueckelberg** mechanism

E. C. G. Stueckelberg 1938

D. Feldman, Z. Liu, P. Nath, B.D. Nelson 2009

Kors & Nath, hep-ph/0402047

V. I. Ogievetskii & I. V. Polubarinov 1962

- Dirac fermion ψ $(g_F X_\mu + g_W C_\mu)\bar{\psi}\gamma^\mu\psi$

Mass matrix of neutral gauge bosons


4 by 4 mass square matrix in $V = (C, X, B, A^3)$

$$m^2 = \begin{pmatrix} m_2^2 & 0 & m_2^2 \epsilon_2 & 0 \\ 0 & m_1^2 & m_1^2 \epsilon_1 & 0 \\ m_2^2 \epsilon_2 & m_1^2 \epsilon_1 & m_1^2 \epsilon_1^2 + m_2^2 \epsilon_2^2 + \frac{g'^2 v^2}{4} & -\frac{g' g v^2}{4} \\ 0 & 0 & -\frac{g' g v^2}{4} & \frac{g^2 v^2}{4} \end{pmatrix}$$


Mass eigenstates $E = (Z', A', Z, A)$ via $E = VO$
where $O^T m^2 O$ is diagonal

ψ couples w/ A due to the neutral gauge bosons mixing; ψ is **millicharged**


CMB constraint



$\sim 1\%$ of DM
 can be charged




Experimental constraints



Timing Detector



CMS timing detector (phase 2)



- between the tracker and calorimeter
- $\delta t = 30\text{ps}$
- $1.17\text{m} \sim \text{O}(\text{ns})$ away from the beam axis

<https://cds.cern.ch/record/2296612/files/LHCC-P-009.pdf>

LLDP @ LHC





$$m_{Z'} \sim O(1)\text{TeV and } \epsilon_2 \sim 10^{-2}$$

$$m_{A'} \sim O(1)\text{GeV and } \epsilon_1 \sim 10^{-7}$$

GeV LLDP w/ $\tau \sim 1\text{m}$

Time delay for LLP



lepton: $p_T > 3 \text{ GeV}$

time delay: $\Delta t > 1.2 \text{ ns}$


DP: $0.2 \text{ m} < L_T < 1.17 \text{ m} \ \& \ z < 3.04 \text{ m}$

ISR jet: $p_T > 30 \text{ GeV} \ \& \ |\eta| < 2.5$


$$\Delta t = \frac{l_X}{\beta_X} + \frac{l_a}{\beta_a} - \frac{l_{SM}}{\beta_{SM}} \quad \beta_a \simeq \beta_{SM} \simeq 1$$

Liu, Liu, Wang, 1805.05957

Lepton p_T distribution




Time delay distribution




- SV: same vertex, due to time resolution: 30ps
- PU: pile-up, due to the spread of proton bunch

Timing detector sensitivity on LLDP




LHCb sensitivity on LLDP



Far forward detectors


- For the **light** LLDP $O(0.1 \sim 1)$ GeV with **longer lifetime** $O(10 \sim 100)$ m, the timing detectors do not probe them effectively
- **Far detectors** with distance $O(10 \sim 100)$ m from the IP can probe the LLDP with **longer lifetime** $O(10 \sim 100)$ m
- At hadron colliders, the **light** final particles are produced more **forward**
- **Far forward** detectors at LHC are ideal places to search the **light** LLDP with **longer lifetime** $O(10 \sim 100)$ m
- Negligible BG

Forward-Aperture CMS ExTension (FACET)




G. Landsberg, "Searches for new physics with FACET: Forward-Aperture CMS ExTension".
<https://indico.cern.ch/event/994582/>


ForwArd Search ExpeRiment (FASER)




LLDP production @ far forward detector



Meson Decay





**Proton
Bremsstrahlung**





Hidden Radiation

The HR mode enhances
the LLDP signal in the far
forward detectors

Lifetime frontier detectors



Lifetime frontier detectors



Conclusion

- “Ordinary” long-lived dark photon has a small LHC signal due to the extremely weak coupling
- We construct a BSM model in which the LLDP signal is greatly **enhanced**
- We analyzed various experimental constraints on the LLDP model
- Sensitivity to the LLDPs in the **precision timing** detector is computed
- Sensitivity to the LLDPs in the **forward** detector is computed... (work in progress)

Conclusion

- “Ordinary” long-lived dark photon has a small LHC signal due to the extremely weak coupling
- We construct a BSM model in which the LLDP signal is greatly **enhanced**
- We analyzed various experimental constraints on the LLDP model
- Sensitivity to the LLDPs in the **precision timing** detector is computed
- Sensitivity to the LLDPs in the **forward** detector is computed... (work in progress)


Thank you!



Backups

Mingxuan Du - LLDP NJU

CMB constraint



~ 1 % of DM
can be charged

