



Contribution ID: 436

Type: **not specified**

## **Search for charginos and neutralinos in final states with two boosted hadronically decaying bosons and missing transverse momentum with the ATLAS experiment**

*Tuesday, 24 August 2021 22:50 (20 minutes)*

Searches for electroweakinos light enough to be produced at the LHC are well motivated by consideration on dark matter, naturalness and the recently observed muon  $g-2$  anomaly. This talk presents a search for electroweakinos in fully-hadronic final states to exploit the advantage of the large branching ratio, and the efficient background rejection by identifying the high- $p_T$  bosons using large-radius jets and jet substructure information. Compared to the more traditionally considered lepton final states, it allows to reach sensitivity to higher electroweakinos masses and to set strong limits on a variety of simplified models. For example, for the case of the wino-bino simplified model a wino mass up to 1060 GeV is excluded. Additionally, more concrete complete models and parameter scans are also considered.

**Primary author:** OKAZAKI, Yuta (Kyoto U)

**Presenters:** OKAZAKI, Yuta (Kyoto U); OKAZAKI, Yuta (Kyoto University (JP))

**Session Classification:** Supersymmetry: Models, Phenomenology and Experimental Results

**Track Classification:** Supersymmetry: Models, Phenomenology and Experimental Results