



Contribution ID: 454

Type: **not specified**

Performance and calibration for the identification of boosted Higgs bosons decaying into beauty quark pairs in ATLAS

Thursday, 26 August 2021 22:35 (20 minutes)

The physics programme at ATLAS involves a variety of Standard Model and Beyond Standard Model resonances decaying to two b quarks, including the Higgs Boson. In order to identify these resonances at high momentum, ATLAS has developed the boosted $X \rightarrow bb$ tagger, a new NN-based tagging algorithm which combines the flavour information of up to three sub-jets associated to the large-R jet capturing the decays of these particles. This talk presents the Monte Carlo performance for the boosted $X \rightarrow bb$ tagger and the corresponding calibration strategy using the full Run-2 dataset gathered by ATLAS and comparing to simulation. Foreseen results include the signal tagging efficiencies derived using $Z (->bb)+jets$ and $Z(->bb)+gamma$ events, and background mistag rates measured using $t\bar{t}$ and $g \rightarrow bb$ splitting in multi-jet events.

Primary author: LI, Changqiao (Shanghai Jiao Tong University (CN))

Presenter: LI, Changqiao (Shanghai Jiao Tong University (CN))

Session Classification: Electroweak, Top quark, and Higgs Physics

Track Classification: Electroweak, Top quark, and Higgs Physics