



FZU

Institute of Physics
of the Czech
Academy of Sciences



ATLAS
EXPERIMENT

Searches for rare top production processes

Jiri Hejbal (Institute of Physics in Prague)
on behalf of the ATLAS Collaboration

Introduction

Run 2 data allow to probe the rarest processes with the lowest cross sections

- ▶ Stringent tests of the Standard Model
- ▶ Tiny anomalies may appear from new physics and can be explored in EFT

Flavour Changing Neutral Currents (FCNC)

- ▶ FCNC processes are forbidden at tree level and highly suppressed at higher order in the Standard Model (SM)

FCNC couplings can be described by an EFT:

$$\mathcal{L}_{\text{eff}} = \mathcal{L}_{\text{SM}} + \frac{1}{\Lambda_{\text{NP}}^2} \sum_k C_k O_k$$

$\Lambda_{\text{NP}} \dots$ scale of new physics



$O_k \dots$ dimension-6 operator

We present result of searches for FCNC and rare SM processes involving top quarks

- ▶ tqg
- ▶ tqZ
- ▶ $t\bar{t}\bar{t}$
- ▶ single top s-channel
- ▶ $tq\gamma$
- ▶ tqH
- ▶ $t\gamma$
- ▶ $t\bar{t}W$ charge asymmetry

For more top quark related results see [Mario's talk](#)

- ▶ Probes single top quark production via FCNC
- ▶ Reconstruct top in $t \rightarrow e/\mu vb$ final states, where $t \rightarrow \tau vb$ may also contribute
 - ▶ =1 lepton, ≥ 1 b-jet, $E_T^{\text{miss}} > 30$ GeV, $m_T(W) > 50$ GeV
 - ▶ Nr. of b-jet to define validation region, in signal region =1 b-jet




- ▶ The analysis targets separate contributions from cgt and ugt
 - ▶ Two Neural Network were used to construct two discriminants D_1, D_2

Upper limits on the production:

$$\begin{aligned} \sigma(ugt) \times \mathcal{B}(t \rightarrow Wb) \times \mathcal{B}(W \rightarrow \ell\nu) &< 3.0 \text{ pb} && 2.4 \text{ pb exp.} \\ \sigma(cgt) \times \mathcal{B}(t \rightarrow Wb) \times \mathcal{B}(W \rightarrow \ell\nu) &< 4.7 \text{ pb} && 2.5 \text{ pb exp.} \\ \mathcal{B}(W \rightarrow \ell\nu) &= 0.325 \end{aligned}$$

- ▶ Leading systematics:
 - ugt : related to W+jets process
 - cgt : modelling of the parton shower


- ▶ Target both production and decay of FCNC $t\gamma q$ vertices
- ▶ Background estimation
 - ▶ $e \rightarrow \gamma$: estimate a fake factor to correct simulation
 - ▶ $h \rightarrow \gamma$: transfer factor from control region
- ▶ Two neural network targeting $t\bar{u}\gamma$ and $t\bar{c}\gamma$ signal separately



▶ Upper limits of BR

Effective coupling	Coefficient limits		Coupling	BRs [10^{-5}]	
	Expected	Observed		Expected	Observed
$ C_{uW}^{(13)*} + C_{uB}^{(13)*} $	$0.104^{+0.020}_{-0.016}$	0.103	$t \rightarrow u\gamma$ LH	$0.88^{+0.37}_{-0.25}$	0.85
$ C_{uW}^{(31)} + C_{uB}^{(31)} $	$0.122^{+0.023}_{-0.018}$	0.123	$t \rightarrow u\gamma$ RH	$1.20^{+0.50}_{-0.33}$	1.22
$ C_{uW}^{(23)*} + C_{uB}^{(23)*} $	$0.205^{+0.037}_{-0.031}$	0.227	$t \rightarrow c\gamma$ LH	$3.40^{+1.35}_{-0.95}$	4.16
$ C_{uW}^{(32)} + C_{uB}^{(32)} $	$0.214^{+0.039}_{-0.032}$	0.235	$t \rightarrow c\gamma$ RH	$3.70^{+1.47}_{-1.03}$	4.46

- ▶ Major systematic: **statistical uncertainty**
- ▶ **Factor of 3.3 – 5.4 improvement wrt ATLAS 13 TeV 81 fb⁻¹ results**
 - ▶ More signal region, more optimised analysis and higher luminosity




ATLAS Simulation

$\sqrt{s} = 13$ TeV

- other prompt γ
- $h \rightarrow \gamma$ fakes
- $Z\gamma$ +jets
- $t\bar{t}\gamma$
- $e \rightarrow \gamma$ fakes
- $W\gamma$ +jets

SR







▶ Target both production and decay of FCNC tqZ vertices:

▶ $Z \rightarrow ll$, semi-leptonic top decay \Rightarrow tri-leptons

▶ Analysis regions

▶ Orthogonality cut applied on reconstructed top mass

▶ ≥ 2 jets, 1 b-jet (SR1) targeting decay mode or ≥ 1 jet, 1 b-jet (SR2) targeting production mode






Observable	Vertex	Coupling	Observed	Expected
SR1+CRs				
$\mathcal{B}(t \rightarrow Zq) [10^{-5}]$	tZu	LH	9.7	$8.6^{+3.6}_{-2.4}$
$\mathcal{B}(t \rightarrow Zq) [10^{-5}]$	tZu	RH	9.5	$8.2^{+3.4}_{-2.3}$
SR2+CRs				
$\mathcal{B}(t \rightarrow Zq) [10^{-5}]$	tZu	LH	7.8	$6.1^{+2.7}_{-1.7}$
$\mathcal{B}(t \rightarrow Zq) [10^{-5}]$	tZu	RH	9.0	$6.6^{+2.9}_{-1.8}$
SRs+CRs				
$\mathcal{B}(t \rightarrow Zq) [10^{-5}]$	tZu	LH	6.2	$4.9^{+2.1}_{-1.4}$
$\mathcal{B}(t \rightarrow Zq) [10^{-5}]$	tZu	RH	6.6	$5.1^{+2.1}_{-1.4}$
$\mathcal{B}(t \rightarrow Zq) [10^{-5}]$	tZc	LH	13	11^{+5}_{-3}
$\mathcal{B}(t \rightarrow Zq) [10^{-5}]$	tZc	RH	12	10^{+4}_{-3}

▶ Upper limits on branching ratios were improved with respect to the previous results by factor 2 - 5

▶ Dominant systematic: **statistical uncertainty**

- Explored both production and decay of FCNC tqH vertices
- Top quark: leptonic or hadronic decay
- $H \rightarrow \tau\tau$: $\tau_{had} \tau_{had}$ OR $\tau_{lep} \tau_{had}$ (depending on τ -lepton decay)



Analysis regions

- Employ seven signal regions in a combination of top and di-tau decay, and additional jets
- BDT is trained in each of the SR to separate signal from SM background

Background estimation

- Fake τ : estimate a transfer factor in CR
- Others: Monte-Carlo simulation



$\mathcal{B}(t \rightarrow cH) < 9.9 \times 10^{-4} (5.0^{+2.2}_{-1.4} \times 10^{-4}),$ assuming $\mathcal{B}(t \rightarrow uH) = 0$

▶ Upper limits of BR:

$\mathcal{B}(t \rightarrow uH) < 7.2 \times 10^{-4} (3.6^{+1.7}_{-1.0} \times 10^{-4}),$ assuming $\mathcal{B}(t \rightarrow cH) = 0$

▶ Limits translate to tqH Wilson coefficients: $C_{c\phi} < 1.38 (0.97)$ and $C_{u\phi} < 1.18 (0.83)$

▶ 2D contours:




▶ Major systematic: statistical uncertainty

▶ A slight excess of data is observed above background with a significance of 2.3σ



▶ A factor of 5 improvement wrt ATLAS 13 TeV 36 fb^{-1} results

$t\bar{t}\bar{t}$ production



JHEP 11 (2021) 118



- ▶ **Measurements done in the all of the leptonic final states**
 - ▶ SS dilepton and multi-lepton channel (**2LSS/ML**) -> [Eur. Phys. J. C 80 \(2020\)](#)
 - ▶ single-lepton and OS dilepton channel (**1L/2LOS**) -> [this talk](#)
- ▶ Never observed by ATLAS or CMS yet
- ▶ Sensitive to the magnitude and CP properties of the Yukawa coupling of top-quark to Higgs boson



- ▶ BDT that is used to separate the signal from the background
- ▶ **Targeting events with high jet and b-jet multiplicities**
 - ▶ 4-top final state features 10 (8) jets in 1L (2LOS) and 4 b-jets at truth level
- ▶ **Pre-selected events are orthogonal to 2LSS/3L**
 - ▶ 1L channel: One lepton ($>28 \text{ GeV}$) and at least 7 jets and at least 2 b-tagged jets
 - ▶ 2LOS channel: Two leptons ($>28, 10 \text{ GeV}$) with OS charge and at least 5 jets and at least 2 b-tagged jets
- ▶ **$t\bar{t} + \text{jets}$ background is estimated using corrected MC simulations**





► Measured cross section for 1L/2LOS : $\sigma_{t\bar{t}t\bar{t}} = 26_{-15}^{+17} \text{ fb}$

- With an observed (expected) significance of 1.9 (1.0) σ
- Uncertainties dominated by 4-top and $t\bar{t}$ +HF modelling uncertainties

► Combined cross section with 2LSS/3L analysis : 24_{-6}^{+7} fb

- With an observed (expected) significance of 4.7 (2.6) σ
- To be compared with the 4.3 σ observed significance from 2LSS/3L analysis

- ▶ First observation of t -channel single top quark production in association with a photon
- ▶ $tq\gamma$ (prod) with observed (expected) significance: 9.1 (6.7) σ
- ▶ Sensitive to EW couplings of the top quark (esp. top- γ vertex)



Cross section measurement



- ▶ Parton level: $\sigma_{tq\gamma} \times B(t \rightarrow l\nu b) = 580 \pm 19 \text{ (stat.)} \pm 63 \text{ (syst.) fb}$
- ▶ Particle level: $\sigma_{tq\gamma} \times B(t \rightarrow l\nu b) + \sigma_{t(\rightarrow l\nu b\gamma)q} = 287 \pm 8 \text{ (stat.)} \pm 31 \text{ (syst.) fb}$

- ▶ ATLAS measurements consistently higher than the prediction by $\sim 40\%$
- ▶ Major systematic uncertainties come from
 - ▶ background modelling: $t\bar{t}\gamma \sim 6\%$; $t\bar{t} \sim 3\%$
 - ▶ MC statistics: $tq\gamma \sim 3\%$; all other processes $\sim 3\%$

Single top s-channel

arXiv:2209.08990
submitted to JHEP

- ▶ single lepton + 2 b-jets events
- ▶ s-channel : the most challenging at the LHC
- ▶ **not yet observed in pp collisions**
- ▶ Sensitive to anomalous couplings





- ▶ Matrix Element Method (for Signal and Background separation):
 - ▶ **hypothesis that a measured final state X is of a certain process H_{proc}**
 - ▶ **normalised fully differential x-section**
 - ▶ **transfer functions**

$$\mathcal{P}(X | H_{\text{proc}}) = \int d\Phi \frac{1}{\sigma_{H_{\text{proc}}}} \frac{d\sigma_{H_{\text{proc}}}}{d\Phi} T_{H_{\text{proc}}}(X | \Phi)$$

- ▶ Probability for a measured event X to be a signal event S :

$$P(S | X) = \frac{\sum_i P(S_i) \mathcal{P}(X | S_i)}{\sum_i P(S_i) \mathcal{P}(X | S_i) + \sum_j P(B_j) \mathcal{P}(X | B_j)}$$

Distribution of the MEM discriminant in the SR before and after the fit



Source	$\Delta\sigma/\sigma$ [%]
$t\bar{t}$ normalisation	+24/-17
$t\bar{t}$ shape modelling	+18/-15
PS & had.	+12/-10
ME/PS matching	+10/-8
h_{damp}	< 1
s-channel modelling	+18/-8
PS & had.	+18/-8
ISR/FSR	+3/-1
Jet energy resolution	+18/-12
Jet energy scale	+18/-13
MC statistics	+13/-11
Flavour tagging	+12/-10
W+jets normalisation	+11/-8
PDFs	+10/-9
$t\bar{t}$	+10/-9
s-channel	± 1
t-channel	± 1
tW	± 1

► Measured cross-section :

$$\sigma_{\text{obs}} = 8.2^{+3.5}_{-2.9} \text{ pb}$$

$$\sigma_{\text{SM}} = 10.32^{+0.40}_{-0.36} \text{ pb}$$

} 3.3 σ (3.9 σ) observed (expected) significance


Main sources of uncertainty

- ▶ First search for the leptonic charge asymmetry of $t\bar{t}W$ in the 3l final state using the full Run2 dataset

▶ Leptonic Charge Asymmetry:

$$A_C^l = \frac{N(\Delta_\eta^l > 0) - N(\Delta_\eta^l < 0)}{N(\Delta_\eta^l > 0) + N(\Delta_\eta^l < 0)}, \text{ where } \Delta_\eta^l = |\eta_{\bar{l}}| - |\eta_l|$$

- ▶ $t\bar{t}W$ process presents larger A_C^l prediction wrt. $t\bar{t}$:
 - ▶ $q\bar{q}$ dominated initial state
 - ▶ ISR W boson polarizes the top pair
- ▶ Lepton-top association is done using a BDT



▶ Observed A_C^l at reconstruction level:

$$A_C^l (t\bar{t}W) = -0.123 \pm 0.136(\text{stat.}) \pm 0.051(\text{syst.})$$

$$\text{Expected: } A_C^l (t\bar{t}W)_{SM} = -0.084_{-0.003}^{+0.005} (\text{scale}) \pm 0.006 (\text{MC stat.})$$

▶ Unfolding to particle level:


$$A_C^l (t\bar{t}W)^{PL} = -0.112 \pm 0.170 (\text{stat.}) \pm 0.055 (\text{syst.})$$

$$\text{Expected: } A_C^l (t\bar{t}W)_{SM}^{PL} = -0.063_{-0.004}^{+0.007} (\text{scale}) \pm 0.004 (\text{MC stat.})$$

- ▶ Analysis is dominated by statistical uncertainties

Summary

- ▶ Recent ATLAS measurements and searches in the associated production of top quarks were presented
- ▶ New results in the investigation of SM rare top processes:
 - ▶ **Strong evidence** for the $t\bar{t}t\bar{t}$ production
 - ▶ **Observation** of $t\gamma$ production
 - ▶ single top s-channel
 - ▶ $t\bar{t}W$ charge asymmetry
- ▶ Highlights of searches for FCNC processes involving top:
 - ▶ tqg , $tq\gamma$, tqZ and tqH
 - ▶ **Significant improvement** of the limits on the BR and the effective coupling strengths wrt previous results



Backup





Branching ratios of top FCNC decays

Process	SM	2HDM(FV)	2HDM(FC)	MSSM	RPV	RS
$t \rightarrow Zu$	7×10^{-17}	–	–	$\leq 10^{-7}$	$\leq 10^{-6}$	–
$t \rightarrow Zc$	1×10^{-14}	$\leq 10^{-6}$	$\leq 10^{-10}$	$\leq 10^{-7}$	$\leq 10^{-6}$	$\leq 10^{-5}$
$t \rightarrow gu$	4×10^{-14}	–	–	$\leq 10^{-7}$	$\leq 10^{-6}$	–
$t \rightarrow gc$	5×10^{-12}	$\leq 10^{-4}$	$\leq 10^{-8}$	$\leq 10^{-7}$	$\leq 10^{-6}$	$\leq 10^{-10}$
$t \rightarrow \gamma u$	4×10^{-16}	–	–	$\leq 10^{-8}$	$\leq 10^{-9}$	–
$t \rightarrow \gamma c$	5×10^{-14}	$\leq 10^{-7}$	$\leq 10^{-9}$	$\leq 10^{-8}$	$\leq 10^{-9}$	$\leq 10^{-9}$
$t \rightarrow hu$	2×10^{-17}	6×10^{-6}	–	$\leq 10^{-5}$	$\leq 10^{-9}$	–
$t \rightarrow hc$	3×10^{-15}	2×10^{-3}	$\leq 10^{-5}$	$\leq 10^{-5}$	$\leq 10^{-9}$	$\leq 10^{-4}$



FCNC tqg - selection requirements

Observable	Common requirements			
$n_{\text{Tight}}(e) + n_{\text{Medium}}(\mu)$	= 1			
$n_{\text{Loose}}(e) + n_{\text{Loose}}(\mu)$	= 1			
$E_{\text{T}}^{\text{miss}}$	> 30 GeV			
$m_{\text{T}}(W)$	> 50 GeV			
$n(j)$	≥ 1			
$p_{\text{T}}(\ell)$	$> 50 \text{ GeV} \cdot \left(1 - \frac{\pi - \Delta\phi(j_1, \ell) }{\pi - 1}\right)$			
	Analysis regions			
	SR	W+jets VR	$t\bar{t}$ VR	tq VR
$n(\eta(j) < 2.5)$	= 1	= 1	= 2	= 1
$n(b)$	= 1	= 1	= 2	= 1
ϵ_b	30%	60% (veto 30%)	30%	30%
$n(\eta(j) > 2.5)$	≥ 0	≥ 0	≥ 0	= 1
$D_{1(2)}$	–	$0.3 < D_{1(2)} < 0.6$	–	$0.2 < D_{1(2)} < 0.4$

FCNC tqg - postfit discriminants



FCNC $tq\gamma$ - Wilson coefficient and BR limits



Effective coupling	Coefficient limits		Coupling	BRs $[10^{-5}]$	
	Expected	Observed		Expected	Observed
$ C_{uW}^{(13)*} + C_{uB}^{(13)*} $	$0.104^{+0.020}_{-0.016}$	0.103	$t \rightarrow u\gamma$ LH	$0.88^{+0.37}_{-0.25}$	0.85
$ C_{uW}^{(31)} + C_{uB}^{(31)} $	$0.122^{+0.023}_{-0.018}$	0.123	$t \rightarrow u\gamma$ RH	$1.20^{+0.50}_{-0.33}$	1.22
$ C_{uW}^{(23)*} + C_{uB}^{(23)*} $	$0.205^{+0.037}_{-0.031}$	0.227	$t \rightarrow c\gamma$ LH	$3.40^{+1.35}_{-0.95}$	4.16
$ C_{uW}^{(32)} + C_{uB}^{(32)} $	$0.214^{+0.039}_{-0.032}$	0.235	$t \rightarrow c\gamma$ RH	$3.70^{+1.47}_{-1.03}$	4.46

Observable	Vertex	Coupling	Observed	Expected
SR1+CRs				
$\mathcal{B}(t \rightarrow Zq) [10^{-5}]$	tZu	LH	9.7	$8.6^{+3.6}_{-2.4}$
$\mathcal{B}(t \rightarrow Zq) [10^{-5}]$	tZu	RH	9.5	$8.2^{+3.4}_{-2.3}$
SR2+CRs				
$\mathcal{B}(t \rightarrow Zq) [10^{-5}]$	tZu	LH	7.8	$6.1^{+2.7}_{-1.7}$
$\mathcal{B}(t \rightarrow Zq) [10^{-5}]$	tZu	RH	9.0	$6.6^{+2.9}_{-1.8}$
SRs+CRs				
$\mathcal{B}(t \rightarrow Zq) [10^{-5}]$	tZu	LH	6.2	$4.9^{+2.1}_{-1.4}$
$\mathcal{B}(t \rightarrow Zq) [10^{-5}]$	tZu	RH	6.6	$5.1^{+2.1}_{-1.4}$
$\mathcal{B}(t \rightarrow Zq) [10^{-5}]$	tZc	LH	13	11^{+5}_{-3}
$\mathcal{B}(t \rightarrow Zq) [10^{-5}]$	tZc	RH	12	10^{+4}_{-3}
$ C_{uW}^{(13)*} $ and $ C_{uB}^{(13)*} $	tZu	LH	0.15	$0.13^{+0.03}_{-0.02}$
$ C_{uW}^{(31)} $ and $ C_{uB}^{(31)} $	tZu	RH	0.16	$0.14^{+0.03}_{-0.02}$
$ C_{uW}^{(23)*} $ and $ C_{uB}^{(23)*} $	tZc	LH	0.22	$0.20^{+0.04}_{-0.03}$
$ C_{uW}^{(32)} $ and $ C_{uB}^{(32)} $	tZc	RH	0.21	$0.19^{+0.04}_{-0.03}$

LH: left hand

RH: right hand

← Higher sensitivity from SR2





► Upper limits on branching ratios, were improved with respect to the previous results

- by factors of 5 (3): LH expected BR limits for $t \rightarrow Zu$ ($t \rightarrow Zc$)
- by factors of 3 (2): LH observed BR limits for $t \rightarrow Zu$ ($t \rightarrow Zc$)
- Inclusion of prod. mode, MVA technique, and higher lumi.

FCNC tqZ - predicted and observed yields in SR

	SR1 ($D_1 > -0.6$)	SR2 ($D_2^u > -0.7$ or $D_2^c > -0.4$)
$t\bar{t}Z + tWZ$	137 ± 12	36 ± 6
$VV + \text{LF}$	18 ± 7	24 ± 8
$VV + \text{HF}$	114 ± 19	162 ± 26
tZ	46 ± 7	108 ± 18
$t\bar{t} + tW$ fakes	14 ± 4	27 ± 8
Other fakes	7 ± 8	5 ± 6
$t\bar{t}W$	4.2 ± 2.1	3.1 ± 1.6
$t\bar{t}H$	4.8 ± 0.7	0.89 ± 0.17
Other bkg.	2.0 ± 1.0	2.5 ± 2.9
FCNC $(u)tZ$	0.9 ± 1.7	4 ± 8
FCNC $t\bar{t}(uZ)$	5 ± 9	0.8 ± 1.5
Total background	348 ± 15	369 ± 21
Data	345	380

FCNC tqZ - postfit discriminants










FCNC $H \rightarrow \tau + \tau^-$: overview of regions

	Regions	b -jet	light flavour jets	lepton	hadronic taus	charge
SR	$t_\ell \tau_{\text{had}} \tau_{\text{had}}$	1	≥ 0	1	2	$\tau_{\text{had}} \tau_{\text{had}}$ OS
	$t_\ell \tau_{\text{had}}^{-1j}$	1	1	1	1	$t_\ell \tau_{\text{had}}$ SS
	$t_\ell \tau_{\text{had}}^{-2j}$	1	2	1	1	$t_\ell \tau_{\text{had}}$ SS
	$t_h \tau_{\text{lep}} \tau_{\text{had}}^{-2j}$	1	2	1	1	$\tau_{\text{lep}} \tau_{\text{had}}$ OS
	$t_h \tau_{\text{lep}} \tau_{\text{had}}^{-3j}$	1	≥ 3	1	1	$\tau_{\text{lep}} \tau_{\text{had}}$ OS
	$t_h \tau_{\text{had}} \tau_{\text{had}}^{-2j}$	1	2	0	2	$\tau_{\text{had}} \tau_{\text{had}}$ OS
	$t_h \tau_{\text{had}} \tau_{\text{had}}^{-3j}$	1	≥ 3	0	2	$\tau_{\text{had}} \tau_{\text{had}}$ OS
VR	$t_\ell \tau_{\text{had}} \tau_{\text{had}}^{-\text{SS}}$	1	≥ 0	1	2	$\tau_{\text{had}} \tau_{\text{had}}$ SS
CRtt	$t_\ell t_\ell 1b \tau_{\text{had}}$	1	≥ 0	2	1	$t_\ell t_\ell$ OS
	$t_\ell t_\ell 2b \tau_{\text{had}}$	2	≥ 0	2	1	$t_\ell t_\ell$ OS
	$t_\ell t_h 2b \tau_{\text{had}}^{-2j} \text{SS}$	2	2	1	1	$t_\ell \tau_{\text{had}}$ SS
	$t_\ell t_h 2b \tau_{\text{had}}^{-2j} \text{OS}$	2	2	1	1	$t_\ell \tau_{\text{had}}$ OS
	$t_\ell t_h 2b \tau_{\text{had}}^{-3j} \text{SS}$	2	≥ 3	1	1	$t_\ell \tau_{\text{had}}$ SS
	$t_\ell t_h 2b \tau_{\text{had}}^{-3j} \text{OS}$	2	≥ 3	1	1	$t_\ell \tau_{\text{had}}$ OS

FCNC $H \rightarrow \tau + \tau^-$: absolute uncertainties

Source of uncertainty	$\Delta B [10^{-5}]$	
	$t \rightarrow uH$	$t \rightarrow cH$
Lepton ID	0.6	1.0
E_T^{miss}	0.7	0.8
Fake lepton modeling	0.9	1.1
JES and JER	2.4	3.2
Flavour tagging	2.7	3.7
$t\bar{t}$ modeling	2.9	4.3
Other MC modeling	2.1	2.9
Fake τ modeling	3.2	4.6
Signal modeling including $\text{Br}(H \rightarrow \tau\tau)$	5.3	7.0
τ ID	3.3	4.4
Luminosity and Pileup	0.9	1.3
MC statistics	5.1	7.0
Total systematic uncertainty	11.2	15.5
Data statistical uncertainty	14.1	19.6
Total uncertainties	18	25

FCNC $H \rightarrow \tau + \tau^-$: t_uH - BDT output distributions



FCNC $H \rightarrow \tau + \tau^-$: tcH - BDT output distributions

