

Poster: Measurement of θ_{13} using reactor antineutrino events with neutron capture on hydrogen at RENO

Wednesday, October 26, 2022 3:40 PM (5 minutes)

The RENO Collaboration reports a measured value of the smallest neutrino mixing angle (θ_{13}) based on ~ 2900 days of reactor electron antineutrino events with a delayed signal of neutron capture on hydrogen (H). The neutron captures on H emitting a 2.2 MeV γ -ray are not easily detected because of high environmental radioactivity below 3.5 MeV. Due to satisfactory purification of liquid scintillator, use of low-radioactivity photomultiplier tube (PMT) glass, and effective selection criteria, it is possible to extract the reactor neutrino signal against the high backgrounds and observe a clear deficit of the reactor neutrino rate. Based on a rate-only analysis, we obtain $\sin^2 2\theta_{13} = 0.086 \pm 0.006(\text{stat}) \pm 0.010(\text{syst})$. This corresponds to a more precisely measured θ_{13} value of the n-H IBD candidates than the previous measurement from 1500 days of data. With the increased data sample, the statistical error of this measurement is reduced by roughly 40%. Based on improved background uncertainties and additional removal of PMT noise events, the systematic error is reduced by roughly 60%.

Primary author: KIM, Sang Yong (Seoul National University)

Presenter: KIM, Sang Yong (Seoul National University)

Session Classification: Poster under break time