

The two-step mechanism explaining the dibaryon $d^*(2380)$ peak

Raquel Molina, Natsumi Ikeno and E. Oset



Table of contents

1. Introduction
2. Triangle Singularities
3. The $pp \rightarrow \pi^+ d$ reaction
4. Results I
5. The $np \rightarrow \pi^+ \pi^- d$ reaction
6. Results II
7. Conclusions

Intro

The dibaryon peak




Figure 1: Total cross sections obtained from PRL106,242302(2011) WASA-at-COSY on $pd \rightarrow d\pi^0\pi^0 + p_{\text{spectator}}$ for the beam energies $T_p = 1.0$ GeV (triangles), 1.2 GeV (dots) and 1.4 GeV (squares). The drawn lines represent the expected cross sections for the Roper excitation process (dotted) and the t -channel $\Delta\Delta$ contribution (dashed) as well as a calculation for a s -channel resonance with $m = 2.37$ GeV and $\Gamma = 68$ MeV (solid).

The dibaryon peak

- The peak is also seen in $pn \rightarrow d\pi^+\pi^-$, WASA-at-COSY (2013)




Figure 2: Total cross sections of the basic double-pionic fusion reactions. Left: $pp \rightarrow d\pi^+\pi^0$, middle: $pn \rightarrow d\pi^0\pi^0$, bottom: $pn \rightarrow d\pi^+\pi^-$. Filled symbols denote results from PLB721, 229 (2013), open symbols from previous works. The crosses denote the result for the $pn \rightarrow d\pi^0\pi^0$ reaction by using Eq. (1) with the data for the $pn \rightarrow d\pi^+\pi^-$ and $pp \rightarrow d\pi^+\pi^0$ channels as input.

$$\sigma(pn \rightarrow d\pi^+\pi^-) = 2\sigma(pn \rightarrow d\pi^0\pi^0) + \frac{1}{2}\sigma(pp \rightarrow d\pi^+\pi^0). \quad (1)$$

- A possible interpretation has been a $\Delta\Delta$ bound state with $I = 0(3^+)$ quantum numbers which are favored. PRL 106, 242302 (2011).

Is this a new 6-quark state OR it is possible another interpretation?

The dibaryon peak

- I. Bar-Nir et al., Nucl. Phys. B 54, 17-28 (1973):

The dominant two pion production plus fusion mechanism comes from a two step single pion production process, $np \rightarrow \pi^- pp$ followed by $pp \rightarrow \pi^+ d$ (plus $np \rightarrow \pi^+ nn$ followed by $nn \rightarrow \pi^- d$).




Figure 3: Two step mechanism for $np \rightarrow \pi^+ \pi^- d$ investigated ($\pi^+ nn$ in the first step is also considered).

The dibaryon peak

$np(l=0) \rightarrow \pi^- pp$ H. Clement and T. Skorodko 2010.09217 (2020)




Figure 4: WASA-at-COSY (solid circles), earlier results (squares & triangles), Dakhno, PLBB 114 (1982), Sarantsev, EPJ. A21 (2004). t -channel Roper excitation from Alvarez-Ruso, Oset et al. NPA633(1998) (dashed line). Solid line, Lorentzian fit ($m = 2315$ MeV and $\Gamma = 150$ MeV).


$$\frac{1}{3}\sigma_{np(l=0) \rightarrow NN\pi} = \sigma_{np \rightarrow pp\pi^-} - \frac{1}{2}\sigma_{pp \rightarrow pp\pi^0}$$

Indeed, $\sigma(pp \rightarrow \pi^+ d) \sim 3 - 4$
mb. **Too BIG** for a fusion
reaction!

C. Richard-Serre et al., NPB20, 413-440 (1970)

P. W. F. Alons et al., NPA480, 413 (1988)

See also GWU SAID data base for $\pi^+ d \rightarrow pp$



Which kind of structure can cause such a
large cross section?


Can the two-step mechanism explain the
dibaryon peak?

Triangle Singularities

Triangle singularities

Landau, Nucl. Phys. 13, (1959), Coleman, Nuovo Cimento 38 (1965)

Coleman-Norton. A triangle singularity appears when the process visualized in the triangle-loop diagram can occur at the classical level, the three intermediate particles can be placed simultaneously on shell and they are collinear.



Example. Scalar three-point loop integral, $p_{13} \equiv k$, $p_{23} \equiv P - k$

$$I_1 = i \int \frac{d^4 q}{(2\pi)^4} \frac{1}{(q^2 - m_2^2 + i\epsilon) [(P - q)^2 - m_1^2 + i\epsilon] [(P - q - k)^2 - m_3^2 + i\epsilon]}$$

Integral in q^0 , it is sufficient to consider:

$$I(m_{23}) = 2\pi \int_0^\infty dq \frac{q^2}{P^0 - \omega_1(q) - \omega_2(q) + i\epsilon} f(q), \quad (3)$$

where $\omega_{1,2}(q) = \sqrt{m_{1,2}^2 + q^2}$, $\omega_3(\vec{q} + \vec{k}) = \sqrt{m_3^2 + (\vec{q} + \vec{k})^2}$,
 $E_{23} = P^0 - k^0$, and

$$f(q) = \int_{-1}^1 dz \frac{1}{E_{23} - \omega_2(q) - \sqrt{m_3^2 + q^2 + k^2 + 2qkz} + i\epsilon}, \quad (4)$$

where $q \equiv \vec{q}$, $k \equiv |\vec{k}| = \sqrt{\lambda(M^2, m_{13}^2, m_{23}^2)}/(2M)$, with $M = \sqrt{P^2}$ and
 $m_{13,23} = \sqrt{p_{13,23}^2}$.

cut 1: $P^0 - \omega_1(\vec{q}) - \omega_2(\vec{q}) + i\epsilon = 0$; sol. $q_{\text{on}+} = q_{\text{on}} + i\epsilon$ (5)

with $q_{\text{on}} = \frac{1}{2M} \sqrt{\lambda(M^2, m_1^2, m_2^2)}$.

Triangle singularities

The function $f(q)$ has endpoint singularities, which are logarithmic branch points when $z = \pm 1$ and,

$$\text{cut 2: } E_{23} - \omega_2(q) - \sqrt{m_3^2 + q^2 + k^2 \pm 2 q k} + i \epsilon = 0, \quad (6)$$

the momentum of particle 2 being **anti-parallel and parallel** to the momentum of the (2,3) system in the $\vec{P} = 0$ frame, sols. $z = -1, q_{a\pm}, z = +1, q_{b\pm}$.

$$q_{a+} = \gamma(v E_2^* + p_2^*) + i \epsilon$$

$$q_{a-} = \gamma(v E_2^* - p_2^*) - i \epsilon$$

$$q_{b+} = \gamma(-v E_2^* + p_2^*) + i \epsilon$$

$$q_{b-} = -\gamma(v E_2^* + p_2^*) - i \epsilon$$

TS condition:

$$\lim_{\epsilon \rightarrow 0} (q_{0n+} - q_{a-}) = 0$$





Figure 5: (a) no pinching, (b) the two-body threshold singularity, and (c), triangle singularity.

The $pp \rightarrow \pi^+ d$ reaction

The $pp \rightarrow \pi^+ d$ reaction






Figure 6: Mechanism for pion production with Δ excitation and np fusion in the deuteron.

(Coleman-Norton) The pp system produces a Δ and N , back to back in the pp rest frame. The Δ decays into $\pi N'$, with the π in the direction of the Δ and N' in its opposite direction (N direction). The N' goes faster than N and catches up with N fusing to give the deuteron.

$$q_{\text{on}} = q_{a-} \longrightarrow \sqrt{s} = 2179 \text{ MeV for } \Gamma_{\Delta} = 0 \text{ (} n, p \text{ on-shell)} \quad (7)$$

Broad peak around $\sqrt{s} = 2165 \text{ MeV}$ for $\Gamma_{\Delta} \simeq 110 \text{ MeV}$.

Diagrams



- **Antisymmetrized pp** $|pp\rangle \equiv \frac{1}{\sqrt{2}} (|\vec{p}, s_1; -\vec{p}, s_2\rangle - |-\vec{p}, s_2; \vec{p}, s_1\rangle)$
- **Deuteron wave function** $|d\rangle \equiv \frac{1}{\sqrt{2}} |pn - np\rangle \chi_d$

χ_d : three spin 1 states ($\uparrow\uparrow$, $\frac{1}{\sqrt{2}}(\uparrow\downarrow + \downarrow\uparrow)$, $\downarrow\downarrow$).

πNN & $\pi N\Delta$ vertices

Isospin factors

$$-i\delta H_{\pi NN} = \frac{f}{m_\pi} \vec{\sigma} \cdot \vec{q} \tau^\lambda; \quad f = 1.00$$

$$-i\delta H_{\pi N\Delta} = \frac{f^*}{m_\pi} \vec{S}^\dagger \cdot \vec{q} T^{\dagger\lambda}; \quad f^* = 2.13$$

$$h_{\Delta\text{-up}} = \frac{4\sqrt{2}}{3}$$

$$h_{\Delta\text{-down}} = -\frac{4\sqrt{2}}{3}$$

Amplitude

$$-it_{ij}^{\pi} = -g_d \frac{4\sqrt{2}}{3} \left(\frac{f^*}{m_{\pi}} \right)^2 \left(\frac{f}{m_{\pi}} \right) \int \frac{d^3 q}{(2\pi)^3} \mathcal{F}_{\pi}(\vec{q}) \left\{ Q_{ij}^{(u)} F(\vec{p}, \vec{q}, \vec{p}_{\pi}) - Q_{ij}^{(d)} F(-\vec{p}, \vec{q}, \vec{p}_{\pi}) \right\}$$

Momentum dependence

$$\begin{aligned} F(\vec{p}, \vec{q}, \vec{p}_{\pi}) = & \frac{M_N}{E_N(-\vec{p} + \vec{q})} \frac{M_N}{E_N(\vec{p} - \vec{q} - \vec{p}_{\pi})} \frac{M_{\Delta}}{E_{\Delta}(\vec{p} - \vec{q})} \frac{1}{2\omega(q)} \\ & \times \frac{\theta(q_{\max} - |\vec{p} - \vec{q} - \frac{\vec{p}_{\pi}}{2}|)}{2p^0 - E_{\pi} - E_N(-\vec{p} + \vec{q}) - E_N(\vec{p} - \vec{q} - \vec{p}_{\pi}) + i\epsilon} \\ & \times \left\{ \frac{1}{p^0 - \omega(q) - E_{\Delta}(\vec{p} - \vec{q}) + i\frac{\Gamma_{\Delta}}{2}} \frac{1}{p^0 - \omega(q) - E_{\pi} - E_N(\vec{p} - \vec{q} - \vec{p}_{\pi}) + i\epsilon} \right. \\ & + \frac{1}{p^0 - \omega(q) - E_{\Delta}(\vec{p} - \vec{q}) + i\frac{\Gamma_{\Delta}}{2}} \frac{1}{2p^0 - E_{\Delta}(\vec{p} - \vec{q}) - E_N(-\vec{p} + \vec{q}) + i\frac{\Gamma_{\Delta}}{2}} \\ & \left. + \frac{1}{p^0 - \omega(q) - E_N(-\vec{p} + \vec{q}) + i\epsilon} \frac{1}{2p^0 - E_{\Delta}(\vec{p} - \vec{q}) - E_N(-\vec{p} + \vec{q}) + i\frac{\Gamma_{\Delta}}{2}} \right\}, \end{aligned}$$

with $\vec{p}_d^{\text{CM}} = \vec{p} - \vec{q} - \frac{\vec{p}_{\pi}}{2}$. **Explicit cuts (TS):** $\sqrt{s} - E_{\Delta} - E_N + i\Gamma_{\Delta}/2 = 0$,
 $(2p^0 = \sqrt{s})$ and $\sqrt{s} - E_{\pi} - E_N - E_{N'} + i\epsilon = 0$.

Amplitude

Pion form factor: $\mathcal{F}_\pi(\vec{q}) = \left(\frac{\Lambda_\pi^2 - m_\pi^2}{\Lambda_\pi^2 + \vec{q}^2} \right)^2$, with values of Λ_π around 1 – 1.2 GeV.

FULL pion propagator. **NOT** done the static approximation

Deuteron, $S = 1$

Transitions, (i) $\uparrow\uparrow$ (1), $\uparrow\downarrow$ (2) to (f) $\uparrow\uparrow$ (1), $\frac{1}{\sqrt{2}}(\uparrow\downarrow + \downarrow\uparrow)$ (2), $\downarrow\downarrow$ (3)

Spin transitions:

$$\begin{aligned} Q_{11}^{\text{up}} &= \left(\frac{2}{3} \vec{p}_\pi \cdot \vec{q} - \frac{i}{3} a_z \right) q_z \\ Q_{11} : \uparrow\uparrow &\rightarrow \uparrow\uparrow \\ Q_{12} : \uparrow\uparrow &\rightarrow \frac{1}{\sqrt{2}}(\uparrow\downarrow + \downarrow\uparrow) \\ Q_{12}^{\text{up}} &= \frac{1}{\sqrt{2}} \left\{ \left(\frac{2}{3} \vec{p}_\pi \cdot \vec{q} - \frac{i}{3} a_z \right) q_+ - \frac{i}{3} a_+ q_z \right\} \\ Q_{13} : \uparrow\uparrow &\rightarrow \downarrow\downarrow \\ Q_{13}^{\text{up}} &= -\frac{i}{3} a_+ q_+ \\ Q_{21} : \uparrow\downarrow &\rightarrow \uparrow\uparrow \\ Q_{21}^{\text{up}} &= \left(\frac{2}{3} \vec{p}_\pi \cdot \vec{q} - \frac{i}{3} a_z \right) q_- \\ Q_{22} : \uparrow\downarrow &\rightarrow \frac{1}{\sqrt{2}}(\uparrow\downarrow + \downarrow\uparrow) \\ Q_{22}^{\text{up}} &= \frac{1}{\sqrt{2}} \left\{ - \left(\frac{2}{3} \vec{p}_\pi \cdot \vec{q} - \frac{i}{3} a_z \right) q_z - \frac{i}{3} a_+ q_- \right\} \\ Q_{23} : \uparrow\downarrow &\rightarrow \downarrow\downarrow \\ Q_{23}^{\text{up}} &= \frac{i}{3} a_+ q_z \end{aligned}$$

Transitions from $\downarrow\uparrow$ and $\downarrow\downarrow$ give a factor 2. Similarly for the Δ -down diagram but $\vec{p} \rightarrow -\vec{p}$.

The deuteron in Field Theory

Deuteron wave function Gammernann, Nieves, Arriola, Oset, PRD(2010)

$$\begin{aligned}V(q, q') &= V\theta(q_{\max} - |\vec{q}|)\theta(q_{\max} - |\vec{q}'|) \\t(q, q') &= t\theta(q_{\max} - |\vec{q}|)\theta(q_{\max} - |\vec{q}'|)\end{aligned}\quad (8)$$

$$t(E) = (V^{-1} - G(E))^{-1}. \quad \tilde{g}_d^2 = \lim_{E \rightarrow M_d} (E - M_d)t = -\frac{1}{\partial G(E)/\partial E}.$$

$$\psi(\vec{p}) \equiv \langle \vec{p} | \psi \rangle = \tilde{g}_d \frac{\theta(q_{\max} - |\vec{p}|)}{E_d - E_N(p) - E_N(p)}, \quad \int d^3p |\langle \vec{p} | \psi \rangle|^2 = 1. \quad (9)$$

We obtain q_{\max} , from the experimental scattering length value, $a = 5.377$ fm,

$$t(E) = \frac{1}{G(M_d) - G(E)}; \quad a = 2\pi^2 M_N \frac{1}{G(M_d) - G(2M_N)}. \quad (10)$$

$$q_{\max} = 240 \text{ MeV and } \tilde{g}_d^2 = (2.68 \times 10^{-3})^2 \text{ MeV}^{-1}, \quad g_d = (2\pi)^{3/2} \tilde{g}_d.$$

Standard formula of **Weinberg**, $g_d^2 = (2\pi)^3 \tilde{g}_d^2 = \frac{8\pi\gamma}{M_N^2}$; $\gamma = \sqrt{M_N B}$, gives $g_d = (2\pi)^{3/2} 2.30 \times 10^{-3} \text{ MeV}^{-1/2}$.

The deuteron in Field Theory

Deuteron wave function: Bonn potential, R. Machleid, PRC(2001)

$$\langle \vec{p} | \psi \rangle = \frac{1}{N} \sum_j \frac{C_j}{\vec{p}^2 + m_j^2}; \quad N^2 = \int d^3 p \left(\sum_j \frac{C_j}{\vec{p}^2 + m_j^2} \right)^2 \quad (11)$$

In our normalization, ($\int d^3 p / (2\pi)^3$ integrations instead of $\int d^3 p$) and including also the weight factors of Field Theory (M/E close to unity),

$$\begin{aligned} & \frac{M_N}{E_N} \frac{M_N}{E_{N'}} \frac{1}{2p^0 - E_\pi - E_N - E_{N'} + i\epsilon} g_d \theta(q_{\max} - |\vec{p} - \vec{q} - \frac{\vec{p}_\pi}{2}|) \\ & \rightarrow (-)(2\pi)^{3/2} \psi(|\vec{p} - \vec{q} - \frac{\vec{p}_\pi}{2}|) \end{aligned} \quad (12)$$

Short range correlations: Oset and Weise, NPA (1979)

In π exchange terms, $S_i^\dagger q_i \sigma_j q_j \frac{1}{q^2 - m_\pi^2}$,

$$q_i q_j \frac{1}{q^2 - m_\pi^2} \simeq q_i q_j \frac{1}{-\vec{q}^2} = (q_i q_j - \frac{1}{3} \vec{q}^2 \delta_{ij}) \frac{1}{-\vec{q}^2} + \frac{1}{3} \frac{\vec{q}^2}{-\vec{q}^2} \delta_{ij}. \quad (13)$$

The NN and $N\Delta$ wave functions have a correlation factor at short distances vanishing at $r \rightarrow 0$ that kills the δ function. Realistic case: $+ g' \delta_{ij}$, $g' \simeq 0.6$ (π and ρ exchange). $\vec{S} \cdot \vec{p}_\pi \vec{S}^\dagger \cdot \vec{q} \vec{\sigma} \cdot \vec{q} \frac{1}{q^2 - m_\pi^2} \rightarrow g' \vec{S} \cdot \vec{p}_\pi \vec{S}^\dagger \cdot \vec{\sigma}$.



Rho exchange Brack, Riska and Weise, NPA (1977)

$$\frac{f^*}{m_\pi} \frac{f}{m_\pi} \longrightarrow \frac{f_\rho^*}{m_\rho} \frac{f_\rho}{m_\rho}; \quad \mathcal{F}_\pi(\vec{q}) \longrightarrow \mathcal{F}_\rho(\vec{q}) = \left(\frac{\Lambda_\rho^2 - m_\rho^2}{\Lambda_\rho^2 + \vec{q}^2} \right)^2; \quad Q_{ij}^\pi \longrightarrow Q_{ij}^\rho$$

$$\frac{\vec{S}_1^\dagger \cdot \vec{q} \vec{\sigma}_2 \cdot \vec{q}}{q^2 - m_\pi^2 + i\epsilon} \longrightarrow \frac{\vec{S}_1^\dagger \times \vec{q} \vec{\sigma}_2 \times \vec{q}}{q^2 - m_\rho^2 + i\epsilon}; \quad F(\text{mom}, m_\pi) \longrightarrow F(\text{mom}, m_\rho)$$

and $f_\rho = 7.96$; $f_\rho^* = 13.53$; $\Lambda_\pi \sim 1.1$ GeV; $\Lambda_\rho \sim 1.8$ GeV.

Impulse approximation



π^+ emission vertex,

$$-i\delta H_{\pi^+ np} = -\sqrt{2} \frac{f}{m_\pi} \vec{\sigma} \cdot \vec{p}_\pi.$$

Amplitude, $\uparrow\uparrow \rightarrow \uparrow\uparrow$

$$-it'_{11} = -F_I p_{\pi,z} [\psi(|\vec{p} - \frac{\vec{p}_\pi}{2}|)$$

$$-\psi(|-\vec{p} - \frac{\vec{p}_\pi}{2}|)]$$

$$\boxed{\frac{d\sigma}{d \cos \theta_\pi} = \frac{1}{4\pi} \frac{1}{s} (M_N)^2 M_d \frac{p_\pi}{p} \overline{\sum} \sum |t|^2} \quad (14)$$

with $\sum |t|^2 \equiv \sum_{ij} |t_{ij}^\pi + t_{ij}^\rho + t_{ij}^{\text{corr.}} + t_{ij}'|^2$ (**Full amplitude**)

Results I




Figure 7: Cross section of $pp \rightarrow \pi^+ d$ as a function of the kinetic energy in the lab frame of the proton. The variable s is $s = 4M_N^2 + 2M_N K_p^{\text{lab}}$.

The peak appears around $\sqrt{s} = 2165$ MeV for fit (d).

Results

	Range of K_p^{lab} and restrictions [MeV]	Parameters [MeV]	N of data	χ^2	χ^2/dof
(a)	$[450:800]$ $M_\Delta \in [1150:1240]$ $\Gamma_\Delta \in [50:200]$ $\Lambda_\pi \in [800:1300]$ $\Lambda_\rho \in [1400:1900]$	$M_\Delta = 1216$ $\Gamma_\Delta = 200$ $\Lambda_\pi = 1175$ $\Lambda_\rho = 1400$	50	415.29	9.03
(b)	$[450:800]$ M_Δ, Γ_Δ fixed $\Lambda_\pi \in [900:1200]$ $\Lambda_\rho \in [1400:1900]$	$M_\Delta = 1232$ $\Gamma_\Delta = 117$ $\Lambda_\pi = 1152$ $\Lambda_\rho = 1900$	50	15883.25	330.90
(c)	$[450:800]$ $M_\Delta \in [1200:1250]$ $\Gamma_\Delta \in [100:150]$ $\Lambda_\pi \in [900:1200]$ $\Lambda_\rho \in [1400:1900]$	$M_\Delta = 1208$ $\Gamma_\Delta = 150$ $\Lambda_\pi = 1014$ $\Lambda_\rho = 1400$	50	792.50	17.23
(d)	$[525:700]$ $M_\Delta \in [1200:1250]$ $\Gamma_\Delta \in [100:150]$ $\Lambda_\pi \in [900:1200]$ $\Lambda_\rho \in [1400:1900]$	$M_\Delta = 1215$ $\Gamma_\Delta = 150$ $\Lambda_\pi = 1015$ $\Lambda_\rho = 1428$	29	101.15	4.05
(e)	$[550:700]$ $M_\Delta \in [1200:1250]$ $\Gamma_\Delta \in [100:150]$ $\Lambda_\pi \in [900:1200]$ $\Lambda_\rho \in [1400:1900]$	$M_\Delta = 1213$ $\Gamma_\Delta = 117$ $\Lambda_\pi = 966$ $\Lambda_\rho = 1552$	25	43.10	2.05

Table 1: We perform fits to Richard-Serre, NPB20 & SAID data, with M_Δ , Γ_Δ , Λ_π , Λ_ρ as free pars. except for (b). $\sigma_{pp \rightarrow \pi+d} = 2 \frac{3}{4} \left(\frac{p_\pi}{p} \right)^2 \sigma_{\pi+d \rightarrow pp}$

Results

With the two initial protons antisymmetrical, $d\sigma/d\cos\theta$ gives the same for \vec{p}_π or $-\vec{p}_\pi$, thus, it depends on $\cos^2\theta_\pi$.






Figure 8: The differential cross section, $d\sigma/d\Omega = \frac{1}{2\pi} \frac{d\sigma}{d\cos\theta_\pi}$, as a function of $\cos^2\theta_\pi$ for $K_p^{\text{lab}} = 570, 616$ and 660 MeV.



Contribution of the different terms




Figure 9: Contribution of the different terms: The π , $\pi + \rho$, $\pi + \rho + \text{corr}$ and $\pi + \rho + \text{corr} + \text{imp}$. Inset: all terms except π -exchange to see the small differences. Pion exchange dominates!. Effect of g' negligible $\implies L = 2$.






Figure 10: Contribution of the different spin transitions. A factor two is included to account for transitions from $\downarrow\uparrow$ and $\downarrow\downarrow$. $\uparrow\downarrow \rightarrow \frac{1}{\sqrt{2}}(\uparrow\downarrow + \downarrow\uparrow)$ dominates!. The initial state combination $\frac{1}{\sqrt{2}}(\uparrow\downarrow - \downarrow\uparrow)$ is the one responsible for the transitions. Altogether, there is dominance of $L = 2, S = 0$ in $pp!$.

The $np \rightarrow \pi^+ \pi^- d$ reaction

The $np \rightarrow \pi^+ \pi^- d$ reaction




The $np \rightarrow \pi^+ \pi^- d$ reaction



$$\begin{aligned}
 (\pi^-) L &= 1, L_z = 0 \\
 J^{(pp)} &= 2, J_z^{(pp)} = 0 \quad \longrightarrow \quad J^{(\pi^+ d)} = 2, J_z^{(\pi^+ d)} = 0
 \end{aligned}$$

The $np \rightarrow \pi^+ \pi^- d$ reaction




$$(\pi^-)L = 1, L_z = 0$$

$$J^{(pp)} = 2, J_z^{(pp)} = 0 \rightarrow J^{(\pi^+ d)} = 2, J_z^{(\pi^+ d)} = 0$$

$$\Rightarrow np^{(I=0)} \equiv |2, 0\rangle|1, 0\rangle = \sqrt{\frac{3}{5}}|(J^{\text{tot}} = 3, 0\rangle - \sqrt{\frac{2}{5}}|1, 0\rangle \equiv |\pi^- \pi^+ d\rangle$$

The $np \rightarrow \pi^+ \pi^- d$ reaction




$$(\pi^-) L = 1, L_z = 0$$

$$J^{(pp)} = 2, J_z^{(pp)} = 0 \rightarrow J^{(\pi^+ d)} = 2, J_z^{(\pi^+ d)} = 0$$

$$\Rightarrow np^{(I=0)} \equiv |2, 0\rangle |1, 0\rangle = \sqrt{\frac{3}{5}} |J^{\text{tot}} = 3, 0\rangle - \sqrt{\frac{2}{5}} |1, 0\rangle \equiv |\pi^- \pi^+ d\rangle$$

$$(np) I = 0, L = 2 \Rightarrow S = 1; |2, 0\rangle |1, 0\rangle = \sqrt{\frac{3}{5}} |3, 0\rangle - \sqrt{\frac{2}{5}} |1, 0\rangle$$

The $np \rightarrow \pi^+ \pi^- d$ reaction



$$(\pi^-)L = 1, L_z = 0$$

$$J^{(pp)} = 2, J_z^{(pp)} = 0 \rightarrow J^{(\pi^+d)} = 2, J_z^{(\pi^+d)} = 0$$

$$\Rightarrow np^{(I=0)} \equiv |2,0\rangle|1,0\rangle = \sqrt{\frac{3}{5}}|(J^{\text{tot}}=)3,0\rangle - \sqrt{\frac{2}{5}}|1,0\rangle \equiv |\pi^- \pi^+ d\rangle$$

$$(np) I = 0, L = 2 \Rightarrow S = 1; |2,0\rangle|1,0\rangle = \sqrt{\frac{3}{5}}|3,0\rangle - \sqrt{\frac{2}{5}}|1,0\rangle$$

$$2s+1 L_J : (np)^3 D_3 \quad (pp)^1 D_2 \quad (\pi^+ d)^3 P_2 \quad \boxed{J^{\text{tot}, P} = 3^+}$$

The $np \rightarrow \pi^+ \pi^- d$ reaction




Figure 11: Two step mechanism for $np \rightarrow \pi^+ \pi^- d$ with explicit Δ excitation in the $pp \rightarrow \pi^+ d$. The mechanism with the nn intermediate state is considered in addition. (Previous work on the two step mechanism: I. Bar-Nir, NPB(1973))

We use experim. data on the $np \rightarrow \pi^- pp$ and $pp \rightarrow \pi^+ d$ reactions.

$$np (I = 0) \rightarrow \pi^- pp: \frac{d\sigma'_{np \rightarrow \pi^- pp}}{dM_{\text{inv}}(p_1 p'_1)} = \frac{1}{4ps} (2M_N)^4 \frac{1}{16\pi^3} p_\pi \tilde{p}_1 |\bar{t}|^2 \frac{1}{2}$$

$M_{\text{inv}}(p_1 p'_1)$: invariant mass of pp ; p : CM momentum of in. $n(p)$; \tilde{p}_1 : momentum of the final p in the pp rest frame.

$$pp \rightarrow \pi^+ d: \sigma_{pp \rightarrow \pi^+ d} = \frac{1}{16\pi M_{\text{inv}}^2(p_1 p'_1)} \frac{p'_\pi}{\tilde{p}_1} |\bar{t}'|^2 (2M_N)^2 (2M_d)$$

p'_π : π^+ momentum in the pp rest frame; $|\bar{t}'|^2$, angle averaged $|t'|^2$.

The $np \rightarrow \pi^+ \pi^- d$ reaction

Amplitude for the two step process

$$-it'' = \frac{1}{2} \int \frac{d^4 p_1}{(2\pi)^4} \frac{(2M_N)^2}{2E_N(p_1)2E_N(p'_1)} \frac{i}{p_1^0 - E_N(p_1) + i\epsilon} \frac{i(-i)t(-i)t'}{\sqrt{s} - p_1^0 - \omega_\pi - E_N(p'_1) + i\epsilon}$$

($\frac{1}{2}$ for the two identical particles). Approximation: take tt' outside the dp_1^0 integration with their on-shell values.

$$t'' \simeq \frac{1}{2} \int \frac{d^3 p_1}{(2\pi)^3} \frac{(2M_N)^2}{2E_N(p_1)2E_N(p'_1)} \frac{tt'}{\sqrt{s} - E_N(p_1) - E_N(p'_1) - \omega_\pi + i\epsilon}$$

Since

$$\frac{1}{M_{\text{inv}}(p_1 p'_1) - 2E_N(p_1) + i\epsilon} \equiv \mathcal{P} \left[\frac{1}{M_{\text{inv}}(p_1 p'_1) - 2E_N(p_1)} \right] - i\pi \delta(M_{\text{inv}}(p_1 p'_1) - 2E_N(p_1))$$

In the on-shell approx. (Bar-Nir), this becomes, $t''_{\text{on}} = -i \frac{1}{2} \frac{\tilde{p}_1}{8\pi} \frac{(2M_N)^2}{M_{\text{inv}}(p_1 p'_1)} t\bar{t}'$.

Off-shell effects can be taken into account by means of,

$$\left(\frac{\tilde{p}_1}{2\pi M_{\text{inv}}(p_1 p'_1)} \right)^2 \rightarrow |G(M_{\text{inv}})|^2 \quad (15)$$

The $np \rightarrow \pi^+ \pi^- d$ reaction

Thus, for the two-step mechanism,

$$\frac{d\sigma_{np \rightarrow \pi^+ \pi^- d}}{dM_{\text{inv}}(\pi^+ \pi^-)} = (2M_N)^2 (2M_d) p_d \tilde{p}_\pi \frac{1}{4} \frac{\tilde{p}_1^2}{64\pi^2} \frac{1}{M_{\text{inv}}^2(p_1 p_1')} \frac{1}{p_\pi \tilde{p}_1} 2|\bar{t}'|^2 \frac{d\sigma_{np \rightarrow \pi^- p p}}{dM_{\text{inv}}(p_1 p_1')}$$

Approximation: $|\bar{t}\bar{t}'|^2 \sim |\bar{t}|^2 |\bar{t}'|^2$, the amplitudes t, t' have smooth angular structure. Adlarson, PLB774, 599 (2017), Richard-Serre, NPB20, 413 (1970)


Bose enhancement

For the two pions in $I = 0$, L must be even. Preferred direction for $L = 0$.

$$\frac{d\sigma_{np \rightarrow \pi^+ \pi^- d}}{dM_{\text{inv}}(\pi^+ \pi^-)} = \sigma_{np \rightarrow \pi^+ \pi^- d} \delta(M_{\text{inv}}(\pi^+ \pi^-) - \bar{M}_{\pi\pi})$$

$$E_{2\pi} = \frac{s + M_{\text{inv}}^2(\pi\pi) - M_d^2}{2\sqrt{s}}, E_\pi \simeq E_{2\pi}/2$$

$$M_{\text{inv}}^2(p_1 p_1') = (P(np) - p_{\pi^-})^2 = s + m_\pi^2 - 2\sqrt{s}E_\pi$$



Bashkanov,
PRL102(2009)

The $np \rightarrow \pi^+ \pi^- d$ reaction

Using

$$2M_{\text{inv}}(p_1 p'_1) dM_{\text{inv}}(p_1 p'_1) = -2\sqrt{s} dE_\pi = -M_{\text{inv}}(\pi\pi) dM_{\text{inv}}(\pi\pi),$$

we obtain

$$\sigma_{np \rightarrow \pi^+ \pi^- d} = \frac{M_{\text{inv}}(p_1 p'_1)}{4\pi} \frac{\sigma_{np \rightarrow \pi^- pp} \sigma_{pp \rightarrow \pi^+ d}}{M_{\text{inv}}(\pi\pi)} \frac{\tilde{p}_1^2}{p_\pi p'_\pi} p_d \tilde{p}_\pi. \quad (16)$$

The amplitudes $np(I=0) \rightarrow \pi^- pp$ and $np(I=0) \rightarrow \pi^+ nn$ are identical up to the phase of π^+ and the same happens for $pp \rightarrow \pi^+ d$ and $nn \rightarrow \pi^- d$. Isospin symmetry gives $\sigma_{np(I=0) \rightarrow pp\pi^-} = \frac{1}{3} \sigma_{np(I=0) \rightarrow NN\pi}$.

$$\sigma_{np \rightarrow \pi^+ \pi^- d} = \frac{M_{\text{inv}}(p_1 p'_1)}{6\pi} \frac{\sigma_{np \rightarrow NN\pi}^I \sigma_{pp \rightarrow \pi^+ d}}{M_{\text{inv}}(\pi\pi)} \frac{\tilde{p}_1^2}{p_\pi p'_\pi} p_d \tilde{p}_\pi$$

with $\sigma_{np \rightarrow NN\pi}^I = \sigma_{np(I=0) \rightarrow NN\pi}$.

Adlarson, PLB774(2017), Clement, 2010.09217 (2020).

Results II

The $np \rightarrow \pi^+ \pi^- d$ reaction

Fit to data

Data for $\sigma_{np(I=0) \rightarrow NN\pi}$ from Fig. 1 of [Clement, Skorodko, 2010.09217 \(2020\)](#). Isospin conservation assumed.

$$\sigma_{np(I=0) \rightarrow pp\pi^-} = \frac{1}{3} \sigma_{np(I=0) \rightarrow NN\pi} = \frac{1}{6} \sigma_{NN(I=0) \rightarrow NN\pi} = \frac{1}{6} 3(2\sigma_{np \rightarrow pp\pi^-} - \sigma_{pp \rightarrow pp\pi^0})$$

5 % isospin symmetry violation give systematic errors of 0.5 mb, which we add in quadrature.

The $np(I=0) \rightarrow NN\pi$ cross section is parameterized as,

$$\sigma_i = \left| \frac{\alpha_i}{\sqrt{s} - \tilde{M}_i + i\tilde{\Gamma}_i/2} \right|^2 \quad (17)$$

- **Set I:** $\tilde{M}_1 = 2326$ MeV, $\tilde{\Gamma}_1 = 70$ MeV, $\alpha_1^2 = 2.6 \left(\frac{\tilde{\Gamma}_1}{2}\right)^2$ mb MeV².
- **Set II:** $\tilde{M}_2 = 2335$ MeV, $\tilde{\Gamma}_2 = 80$ MeV, $\alpha_2^2 = 2.5 \left(\frac{\tilde{\Gamma}_2}{2}\right)^2$ mb MeV².

$$pp \rightarrow \pi^+ d: \sigma_3 = \left| \frac{\alpha_3}{M_{\text{inv}(p_1 p'_1)} - \tilde{M}_3 + i\tilde{\Gamma}_3/2} \right|^2 \text{ with } \tilde{M}_3 = 2165 \text{ MeV, } \tilde{\Gamma}_3 = 123.27$$

MeV, $\alpha_3^2 = 3.186 \left(\frac{\tilde{\Gamma}_3}{2}\right)^2$ mb MeV². Data from Richard-Serre, NPB20 (1970).

The $np \rightarrow \pi^+ \pi^- d$ reaction




Figure 12: Left: Results with set I for $\sigma_{np \rightarrow \pi^- pp}(I=0)$ and $\sigma_{pp \rightarrow \pi^+ d}$. The results with $\sigma_{np \rightarrow \pi^+ \pi^- d}$ in $I=0$ are multiplied by 10 for a better comparison. Data for $np(I=0) \rightarrow \pi NN$ are taken from Dakhno et al., and WASA-at-COSY^(*), including systematic errors from isospin violation. Inset: $\sigma_{pp \rightarrow \pi^+ d}$ in comparison with data for $pp \rightarrow \pi^+ d$ from Richard-Serre. Right: Experimental data $pn \rightarrow d\pi^0\pi^0$. Filled symbols denote results from Adlarson (2013), open symbols from previous works, and crosses the result with Eq. (1).

The $np \rightarrow \pi^+ \pi^- d$ reaction




Figure 13: Left: Results for set II; Right: Experimental data $pn \rightarrow d\pi^+\pi^-$. Filled symbols denote results from Adlarson, PLB721, 229 (2013), open symbols from previous works.

The $np \rightarrow \pi^+\pi^-d$ reaction

	$\delta\bar{M}_{\pi\pi}$ (MeV)			$p_{1,\max}^{\text{O.s.}}$ (MeV)	
Set I	40	60	80	700	800
strength (mb)	0.72	0.76	0.75	0.82	0.95
position (MeV)	2332	2332	2332	2332	2332
width (MeV)	76	76	81	75	75
Set II					
strength (mb)	0.75	0.80	0.80	0.85	0.96
position (MeV)	2342	2345	2345	2343	2342
width (MeV)	86	87	88	87	84

Table 2: Peak strength, position, and width, for intermediate particles on shell ($\bar{M}_{\pi\pi} = 2m_\pi + \delta\bar{M}_{\pi\pi}$), and off-shell (“o.s.”), $\delta\bar{M}_{\pi\pi} = 60$ MeV.

We obtain the peak strength $\sim 0.76 - 0.96$ mb, the position, 2332 – 2345 MeV, and the width, 75 – 88 MeV, to be compared with the experimental one, with strength ~ 0.5 mb, at 2370 MeV.

Conclusions

Conclusions

- We showed that the $pp \rightarrow \pi^+ d$ reaction has a **triangle singularity**
- The sequential one pion production in $np(l=0) \rightarrow \pi^- pp$ followed by $pp \rightarrow \pi^+ d$ offers a natural explanation of the peak so far associated to the dibaryon $d^*(2380)$
- The mechanism naturally gives rise to the dominant **quantum numbers** observed in the experiments (3^+).