

The η and η' physics at JLab

Liping Gan

University of North Carolina Wilmington
(for the GlueX Collaboration)


Outline

1. Overview of η and η' physics
2. Two experiments in Hall D at JLab
 - PrimEx-eta experiment
 - JLab Eta Factory (JEF) experiment
3. Summary

Why η is a unique probe for QCD and BSM physics?

- ◆ A **Goldstone** boson due to spontaneous breaking of chiral symmetry in QCD

→ η plays important role in bridging our understanding of low-energy hadron dynamics and underlying QCD



- ◆ All its possible strong and EM decays are forbidden in the lowest order so that η has **narrow** decay width ($\Gamma_\eta = 1.3 \text{ KeV}$ compared to $\Gamma_\omega = 8.5 \text{ MeV}$)

→ Enhanced sensitivity to the higher order contributions (by a factor of ~ 7000 compared to ω decays) for new physics search

- ◆ Eigenstate of P, **C**, CP, and G: $I^G J^{PC} = 0^+ 0^{-+}$

→ tests for **C, CP**

- ◆ All its additive quantum numbers are zero and its decays are **flavor-conserving**

→ effectively “free” of SM backgrounds for new physics search.

Rich η and η' Physics

Standard Model Tests:

- Chiral symmetry and anomalies
- Extract η - η' mixing angle and quark mass ratio
- Theory inputs to HLbL for $(g-2)_\mu$
- Scalar dynamics in ChPT

Fundamental Symmetry Tests:

- C, CP violations
- P, CP violations

BSM Physic Searches:


- Vector bosons (B boson, dark photon and X boson)
- Dark scalars
- Pseudoscalars (ALPs)
- BSM weak decays

Channel	Expt. branching ratio	Discussion
$\eta \rightarrow 2\gamma$	39.41(20)%	chiral anomaly, η - η' mixing
$\eta \rightarrow 3\pi^0$	32.68(23)%	$m_u - m_d$
$\eta \rightarrow \pi^0\gamma\gamma$	$2.56(22) \times 10^{-4}$	χ PT at $O(p^6)$, leptophobic B boson, light Higgs scalars
$\eta \rightarrow \pi^0\pi^0\gamma\gamma'$	$< 1.2 \times 10^{-3}$	χ PT, axion-like particles (ALPs)
$\eta \rightarrow 4\gamma$	$< 2.8 \times 10^{-4}$	$< 10^{-11}$ [54]
$\eta \rightarrow \pi^+\pi^-\pi^0$	22.92(28)%	$m_u - m_d$, C/CP violation, light Higgs scalars
$\eta \rightarrow \pi^+\pi^-\gamma$	4.22(8)%	chiral anomaly, theory input for singly-virtual TFF and $(g-2)_\mu$, P/CP violation
$\eta \rightarrow \pi^+\pi^-\gamma\gamma$	$< 2.1 \times 10^{-3}$	χ PT, ALPs
$\eta \rightarrow e^+e^-\gamma$	$6.9(4) \times 10^{-3}$	theory input for $(g-2)_\mu$, dark photon, protophobic X boson
$\eta \rightarrow \mu^+\mu^-\gamma$	$3.1(4) \times 10^{-4}$	theory input for $(g-2)_\mu$, dark photon
$\eta \rightarrow e^+e^-$	$< 7 \times 10^{-7}$	theory input for $(g-2)_\mu$, BSM weak decays
$\eta \rightarrow \mu^+\mu^-$	$5.8(8) \times 10^{-6}$	theory input for $(g-2)_\mu$, BSM weak decays, P/CP violation
$\eta \rightarrow \pi^0\pi^0\ell^+\ell^-$		C/CP violation, ALPs
$\eta \rightarrow \pi^+\pi^-\ell^+e^-$	$2.68(11) \times 10^{-4}$	theory input for doubly-virtual TFF and $(g-2)_\mu$, P/CP violation, ALPs
$\eta \rightarrow \pi^+\pi^-\mu^+\mu^-$	$< 3.6 \times 10^{-4}$	theory input for doubly-virtual TFF and $(g-2)_\mu$, P/CP violation, ALPs
$\eta \rightarrow e^+e^-e^+e^-$	$2.40(22) \times 10^{-5}$	theory input for $(g-2)_\mu$
$\eta \rightarrow e^+e^-\mu^+\mu^-$	$< 1.6 \times 10^{-4}$	theory input for $(g-2)_\mu$
$\eta \rightarrow \mu^+\mu^-\mu^+\mu^-$	$< 3.6 \times 10^{-4}$	theory input for $(g-2)_\mu$
$\eta \rightarrow \pi^+\pi^-\pi^0\gamma$	$< 5 \times 10^{-4}$	direct emission only
$\eta \rightarrow \pi^+e^-\nu_e$	$< 1.7 \times 10^{-4}$	second-class current
$\eta \rightarrow \pi^+\pi^-$	$< 4.4 \times 10^{-6}$ [55]	P/CP violation
$\eta \rightarrow 2\pi^0$	$< 3.5 \times 10^{-4}$	P/CP violation
$\eta \rightarrow 4\pi^0$	$< 6.9 \times 10^{-7}$	P/CP violation

arXiv:2007.00664

Primakoff Program at JLab 6 & 12 GeV

Precision measurements of electromagnetic properties of π^0 , η , η' via Primakoff effect



a) Two-Photon Decay Widths:

- 1) $\Gamma(\pi^0 \rightarrow \gamma\gamma)$ @ 6 GeV
- 2) $\Gamma(\eta \rightarrow \gamma\gamma)$
- 3) $\Gamma(\eta' \rightarrow \gamma\gamma)$

Input to Physics:

- precision tests of chiral symmetry and anomalies
- determination of light quark mass ratio
- η - η' mixing angle
- input to calculate HLbL in $(g-2)_\mu$

b) Transition Form Factors

at Q^2 of 0.001-0.5 GeV^2/c^2 :


$$F(\gamma\gamma^* \rightarrow \pi^0), F(\gamma\gamma^* \rightarrow \eta), F(\gamma\gamma^* \rightarrow \eta')$$

Input to Physics:

- π^0, η and η' electromagnetic interaction radii
- is the η' an approximate Goldstone boson?
- input to calculate HLbL in $(g-2)_\mu$

Primakoff Program at JLab 6 & 12 GeV

Precision measurements of electromagnetic properties of π^0 , η , η' via Primakoff effect




a) Two-Photon Decay Widths:

- 1) $\Gamma(\pi^0 \rightarrow \gamma\gamma)$ @ 6 GeV
- 2) $\Gamma(\eta \rightarrow \gamma\gamma)$
- 3) $\Gamma(\eta' \rightarrow \gamma\gamma)$


Input to Physics:

- precision tests of chiral symmetry and anomalies
- determination of light quark mass ratio
- η - η' mixing angle
- input to calculate HLbL in $(g-2)_\mu$



Primakoff Program at JLab 6 & 12 GeV

Precision measurements of electromagnetic properties of π^0 , η , η' via Primakoff effect



a) Two-Photon Decay Widths:

- 1) $\Gamma(\pi^0 \rightarrow \gamma\gamma)$ @ 6 GeV
- 2) $\Gamma(\eta \rightarrow \gamma\gamma)$ (PrimEx-eta)
- 3) $\Gamma(\eta' \rightarrow \gamma\gamma)$

Input to Physics:

- precision tests of Chiral symmetry and anomalies
- determination of light quark mass ratio
- η - η' mixing angle
- input to calculate HLbL in $(g-2)_\mu$

b) Transition Form Factors

at Q^2 of 0.001-0.5 GeV^2/c^2 :


$$F(\gamma\gamma^* \rightarrow \pi^0), F(\gamma\gamma^* \rightarrow \eta), F(\gamma\gamma^* \rightarrow \eta')$$

Input to Physics:


- π^0, η and η' electromagnetic interaction radii
- is the η' an approximate Goldstone boson?
- input to calculate HLbL in $(g-2)_\mu$

Physics for $\Gamma(\eta \rightarrow \gamma\gamma)$ Measurement

1. Resolve long standing discrepancy between collider and Primakoff measurements:



2. Extract η - η' mixing angle:



3. Improve calculation of the η -pole contribution to Hadronic Light-by-Light (HLbL) scattering in $(g-2)_\mu$

4. Improve all partial decay widths in the η -sector


Precision Determination Light Quark Mass Ratio

A clean probe for quark mass ratio:
$$Q^2 = \frac{m_s^2 - \hat{m}^2}{m_d^2 - m_u^2}, \quad \text{where } \hat{m} = \frac{1}{2}(m_u + m_d)$$

➤ $\eta \rightarrow 3\pi$ decays through isospin violation: $A = (m_u - m_d)A_1 + a_{em}A_2$

➤ a_{em} is small

➤ Amplitude:
$$A(h \rightarrow 3p) = \frac{1}{Q^2} \frac{m_K^2}{m_p^2} (m_p^2 - m_K^2) \frac{M(s, t, u)}{3\sqrt{3}F_p^2}$$




arXiv:2007.00664

Precision Determination Light Quark Mass Ratio

A clean probe for quark mass ratio:
$$Q^2 = \frac{m_s^2 - \hat{m}^2}{m_d^2 - m_u^2}, \quad \text{where } \hat{m} = \frac{1}{2}(m_u + m_d)$$


- $\eta \rightarrow 3\pi$ decays through isospin violation: $A = (m_u - m_d)A_1 + a_{em}A_2$
- a_{em} is small
- Amplitude:
$$A(h \rightarrow 3p) = \frac{1}{Q^2} \frac{m_K^2}{m_p^2} (m_p^2 - m_K^2) \frac{M(s, t, u)}{3\sqrt{3}F_p^2}$$




- Critical input to extract Cabibbo Angle, $V_{us} = \sin(q_c)$ from kaon or hyperon decays.
- V_{us} is a cornerstone for test of CKM unitarity:
$$|V_{ud}|^2 + |V_{us}|^2 + |V_{ub}|^2 = 1$$

arXiv:2007.00664

Primakoff Method



$$\frac{d\sigma_{\text{Pr}}}{d\Omega} = \Gamma_{\gamma\gamma} \frac{8\alpha Z^2}{m_\pi^3} \frac{\beta^3 E^4}{Q^4} |F_{e.m.}(Q)|^2 \sin^2 \theta_\pi$$



Requirement:

- Photon flux
 - Beam energy
 - Compact nuclear target
- ← Tagged photon beam

- Peaked at very small forward angle

$$\langle \theta_{\text{Pr}} \rangle_{\text{peak}} \propto \frac{m^2}{2E^2}$$

- Beam energy sensitive:

$$\left\langle \frac{d\sigma_{\text{Pr}}}{d\Omega} \right\rangle_{\text{peak}} \propto E^4, \quad \int d\sigma_{\text{Pr}} \propto Z^2 \log(E)$$

- Coherent process

Experimental Challenges

Compared to π^0 :

- η mass is a factor of 4 larger
- smaller Primakoff cross section

$$\left\langle \frac{dS_{\text{Pr}}}{dW} \right\rangle_{\text{peak}} \propto \frac{E^4}{m^3}$$

- larger overlap between Primakoff and hadronic processes;

$$\langle q_{\text{Pr}} \rangle_{\text{peak}} \propto \frac{m^2}{2E^2} \quad \langle q_{\text{NC}} \rangle_{\text{peak}} \propto \frac{2}{E \cdot A^{1/3}}$$

- larger momentum transfer (coherency, form factors, FSI,...)

Experimental Challenges

Compared to π^0 :

- η mass is a factor of 4 larger
- smaller Primakoff cross section

$$\left\langle \frac{dS_{\text{Pr}}}{dW} \right\rangle_{\text{peak}} \propto \frac{E^4}{m^3}$$

- larger overlap between Primakoff and hadronic processes

$$\langle q_{\text{Pr}} \rangle_{\text{peak}} \propto \frac{m^2}{2E^2} \quad \langle q_{\text{NC}} \rangle_{\text{peak}} \propto \frac{2}{E \cdot A^{1/3}}$$

- larger momentum transfer (coherency, form factors, FSI,...)



1. Higher beam energy
2. Light targets

Advantage of Light Targets

Low A targets to control:

- Coherency: compact nucleus
- Separate background

$$\langle q_{Pr} \rangle_{peak} \propto \frac{m^2}{2E^2} \quad \langle q_{NC} \rangle_{peak} \propto \frac{2}{E \cdot A^{1/3}}$$


- Well known form factors

Hydrogen:


- No inelastic hadronic contribution
- No nuclear final state interactions

^4He :

- Higher Primakoff cross section: $\sigma_{Pr} \propto Z^2$





PrimEx-eta Experiment on $\Gamma(\eta \rightarrow \gamma\gamma)$ in Hall D



- Tagged photon beam (~8.0-11.7 GeV)
- Pair spectrometer and a TAC detector for the photon flux control
- Liquid Hydrogen (3.5% R.L.) and ^4He targets (~4% R.L.)
- Forward Calorimeter (FCAL) detects the $\eta \rightarrow \gamma\gamma$ decay photons
- **CompCal** and FCAL to measure electron Compton scattering for control of overall systematics.

Status of PrimEx-eta


- A engineering run with a new CompCal in fall 2018
- The first production run (Feb 22 - April 15, 2019)
 - ~50% ^4He data was collected
- Second run will start in a week
- Third run is scheduled for 2022




Time Spent on each Target during Spring-2019

Preliminary Result from the First Run

$\eta \rightarrow \gamma\gamma$ yield [$8.0 \text{ GeV} < E_\gamma < 11.2 \text{ GeV}$]




Control of Systematics with Compton Scattering



(By D. Smith)

JLab Eta Factory (JEF) Experiment



Simultaneously measure η/η' decays: $\eta \rightarrow \pi^0 \gamma \gamma$, $\eta \rightarrow 3\gamma$, and ...

- ◆ η/η' produced on LH₂ target with **8.4-11.7 GeV tagged photon beam**:
 $\gamma + p \rightarrow \eta/\eta' + p$
- ◆ Reduce non-coplanar backgrounds by **detecting recoil protons** with GlueX detector
- ◆ Upgraded Forward Calorimeter with **High resolution, high granularity PWO** insertion (**FCAL-II**) to detect multi-photons from the η/η' decays

Production Rate

JEF for 100 days of beam:

	η	η'
Tagged mesons	6.5×10^7	4.9×10^7

Previous Experiments:

Experiment	Total η	Total η'
CB at AGS	10^7	-
CB MAMI-B	2×10^7	-
CB MAMI-C	6×10^7	10^6
WASA-COSY	$\sim 3 \times 10^7$ (p+d), $\sim 5 \times 10^8$ (p+p)	-
KLOE-II	3×10^8	5×10^5
BESIII	$\sim 10^7$	$\sim 5 \times 10^7$

JEF offers a competitive η/η' factory


Uniqueness of JEF Experiment

1. Highly suppressed background in decay channels:
 - a) η/η' energy boost; b) an upgraded calorimeter (FCAL-II)

A2 at MAMI: $\gamma p \rightarrow \eta p$ ($E_\gamma = 1.5$ GeV)

(P.R. C90, 025206)

JEF: $\gamma p \rightarrow \eta p$ ($E_\gamma = 8.4-11.7$ GeV)



2. Simultaneously produce tagged η and η' with similar rates
($\sim 5 \times 10^7$ per 100 beam days)

Main JEF Physics Objectives

1. Search for sub-GeV hidden bosons

vector:

- Leptophobic vector B'

$$\eta, \eta' \rightarrow B' \gamma \rightarrow \pi^0 \gamma \gamma, (0.14 < m_{B'} < 0.62 \text{ GeV});$$

$$\eta' \rightarrow B' \gamma \rightarrow \pi^+ \pi^- \pi^0 \gamma, (0.62 < m_{B'} < 1 \text{ GeV}).$$

- Hidden or dark photon: $\eta, \eta' \rightarrow X \gamma \rightarrow e^+ e^- \gamma$.

scalar S: $\eta \rightarrow \pi^0 S \rightarrow \pi^0 \gamma \gamma, \pi^0 e^+ e^-, (10 \text{ MeV} < m_S < 2m_\pi);$

$$\eta, \eta' \rightarrow \pi^0 S \rightarrow 3\pi, \eta' \rightarrow \eta S \rightarrow \eta \pi \pi, (m_S > 2m_\pi).$$

Axion-Like Particles (ALP): $\eta, \eta' \rightarrow \pi \pi a \rightarrow \pi \pi \gamma \gamma, \pi \pi e^+ e^-$


2. Directly constrain CVPC new physics: $\eta^{(\prime)} \rightarrow 3\gamma, \eta^{(\prime)} \rightarrow 2\pi^0 \gamma, \eta^{(\prime)} \rightarrow \pi^+ \pi^- \pi^0$

3. Precision tests of low-energy QCD:

- Interplay of VMD & scalar dynamics in ChPT: $\eta \rightarrow \pi^0 \gamma \gamma \quad \eta' \rightarrow \pi^0 \gamma \gamma$
- Transition Form Factors of $\eta^{(\prime)}$: $\eta^{(\prime)} \rightarrow e^+ e^- \gamma$

4. Improve the quark mass ratio via Dalitz distributions of $\eta \rightarrow 3\pi$

Landscape of BSM Physics Search



Complementary to other types of experiments, η/η' decays in JEF offer unique sensitivity for new physics that are flavor-conserving, light quark-coupling, PC-conserving.

A Key Channel: $\eta \rightarrow \pi^0 \gamma \gamma$

❖ Search for sub-GeV gauge bosons

- A leptophobic **vector** B' :


$$\eta \rightarrow \gamma B', B' \rightarrow \pi^0 \gamma$$

PL, B221, 80
PR, D89, 114008

- An **scalar** S :


$$\eta \rightarrow \pi^0 S, S \rightarrow \gamma \gamma$$

PR, D100, 095020
Nucl. Phys. B, 114638



❖ A rare window to probe interplay of VMD & scalar resonance in ChPT


PR, C78, 015206



Search for CVPC Interaction via $\eta \rightarrow 3\gamma$

- ◆ SM contribution:
 $BR(\eta \rightarrow 3\gamma) < 10^{-19}$ via P-violating weak interaction.
- ◆ **A new C- and T-violating, and P-conserving interaction** was proposed by Bernstein, Feinberg and Lee
Phys. Rev., 139, B1650 (1965)
- ◆ A calculation due to such new physics by Tarasov suggests:
 $BR(\eta \rightarrow 3\gamma) < 10^{-2}$

Sov.J.Nucl.Phys., 5, 445 (1967)



Improve BR upper limit by one order of magnitude to directly tighten the constraint on CVPC new physics

Improve Quark-Mass Ratio via $\eta \rightarrow 3\pi$ Dalitz Distributions


◆ A clean probe for quark mass ratio: $Q^2 = \frac{m_s^2 - \hat{m}^2}{m_d^2 - m_u^2} \quad \hat{m} = \frac{m_u + m_d}{2}$

➤ decays through isospin violation: $A = (m_u - m_d)A_1 + a_{em}A_2$


➤ a_{em} is small

➤ Amplitude: $A(s, t, u) = \frac{1}{Q^2} \frac{m_K^2}{m_\pi^2} (m_\pi^2 - m_K^2) \frac{M(s, t, u)}{3\sqrt{3}F_\pi^2}$

◆ Uncertainties in quark mass ratio



e-Print: [2007.00664](https://arxiv.org/abs/2007.00664)




Status of JEF



1. JEF was approved by PAC in 2017 to run concurrently with GlueX-II, and it passed PAC jeopardy review on Sept 25, 2020.

2. Development of a 40x40 array of PWO insert for FCAL-II is in progress:

- Constructed a 12x12 array of PWO prototype calorimeter and successfully tested and used for the PrimEx-eta experiment in 2019.
- Engineering design for calorimeter frame is finalized
- PWO module ($2 \times 2 \times 20 \text{ cm}^3$) mechanical design is complete and mass production is in progress
- Installation is anticipated in 2023



prototype calorimeter



PWO modules



3. Data taking is expected in 2024


Summary

- ◆ The JLab η/η' decay measurements offer a rich physics program:
 - **Precision tests of Standard Model:** Chiral symmetry and anomalies; inputs to HLbL in $(g-2)_\mu$ calculation; improve the light quark mass and η - η' mixing angle; scalar dynamics in ChPT
 - **Search for sub-GeV hidden bosons:** vectors, scalars, and ALPs
 - **Fundamental symmetries tests:** directly constrain CVPC new physics
- ◆ The PrimEx-eta experiment on $\Gamma(\eta \rightarrow \gamma\gamma)$ is on-going.
- ◆ The JEF experiment will measure η and η' decays simultaneously, with **two orders of magnitude background reduction** in the rare neutral modes compared to other facilities. Upgrade of Forward calorimeter with a PWO insert is currently under construction. JEF run will be expected in 2024.

Thanks for support by NSF PHY-1812396 and PHY-211181



Uncertainties of on $\Gamma(\pi^0 \rightarrow \gamma\gamma)$

Item	PrimEx-II
Beam parameters	0.2%
Photon flux	0.8%
Target	0.3%
DAQ	0.1%
Event selection	0.2%
Monte-Carlo simulation	0.6%
Yield extraction	1.0%
Photoproduction theory parameters	0.4%
Systematics	1.5%
Statistical	0.7%
Total	1.6%





Differential Cross Sections

PrimEx I:



PrimEx II:






Fitting data with new theoretical calculations to extract $\Gamma(\pi^0 \rightarrow \gamma\gamma)$

Phys.Rev. C80, 055201 (2009); Phys.Part.Nucl.Lett.,9,3 (2012)

Verification of Overall Experimental Systematics with Compton Scattering

Phys.Lett. B797 (2019) 4884



Systematic uncertainties of measured cross section are controlled at 1.5%