The equation of state form Lattice QCD with finite μ_B and μ_S

arXiv:2202.05574, [Borsanyi:2022qlh]

Szabolcs Borsanyi, Zoltan Fodor, Jana N. Guenther, Ruben Kara, Paolo Parotto, Attila Pasztor, Claudia Ratti and Kalman K. Szabo

Avril 6th 2022

BERGISCHE UNIVERSITÄT WUPPERTAL

WB collaboration

BMW collaboration

lattice simulations | real chemical potentials

The sign problem

The QCD partition function:

$$Z(V,T,\mu) = \int \mathcal{D}U\mathcal{D}\psi\mathcal{D}\bar{\psi}\; e^{-S_F(U,\psi,\bar{\psi})-\beta S_G(U)}$$
$$= \int \mathcal{D}U \det M(U) e^{-\beta S_G(U)}$$

- For Monte Carlo simulations $\det M(U)e^{-\beta S_G(U)}$ is interpreted as Boltzmann weight
- If there is particle-antiparticle-symmetry $\det M(U)$ is real
- If $\mu^2 > 0$ $\det M(U)$ is complex

Analytic continuation from imaginary chemical potential



Common technique:

- [deForcrand:2002hgr]
- [Bonati:2015bha]
- [Cea:2015cya]
- [DElia:2016jqh]
- [Bonati:2018nut]
- [Borsanyi:2018grb]
- [Borsanyi:2020fev]
- [Bellwied:2021nrt]
- . . .

Trouble with the equation of state



[Borsanyi:2021sxv], [Borsanyi:2018grb], $N_t = 12$

Trouble with the equation of state







[Borsanyi:2021sxv], [Borsanyi:2018grb], $N_t = 12$



Taylor method

[Bazavov:2017dus]

[Bollweg:2022rps]

Trouble with the equation of state







[Borsanyi:2021sxv]



Taylor method

[Bazavov:2017dus]

[Bollweg:2022rps]

Trouble with the equation of state







[Borsanyi:2021sxv]



Taylor method

[Bazavov:2017dus]

[Bollweg:2022rps]



Results at $\mu_S = 0$

Find a different extrapolation scheme for extrapolating to higher μ_B.





- [Borsanyi:2021sxv]
- $N_t = 10, 12, 16$

Strangeness Neutrality

Enforcing the conditions $\mu_Q = 0$ and $\chi_1^S = 0$:

$$\frac{\mathrm{d}\mu_S}{\mathrm{d}\mu_B} = -\frac{\chi_{11}^{BS}}{\chi_2^S}.$$

On this line, total derivatives with respect to the baryochemical potential read

$$\frac{\mathrm{d}}{\mathrm{d}\hat{\mu}_B} = \frac{\partial}{\partial\hat{\mu}_B} + \frac{\mathrm{d}\hat{\mu}_S}{\mathrm{d}\hat{\mu}_B}\frac{\partial}{\partial\hat{\mu}_S} = \frac{\partial}{\partial\hat{\mu}_B} - \frac{\chi_{11}^{BS}}{\chi_2^S}\frac{\partial}{\partial\hat{\mu}_S}.$$

For the pressure we get:

$$c_n^B(T,\hat{\mu}_B) \equiv \left.\frac{\mathrm{d}^n\hat{p}(T,\hat{\mu}_B)}{\mathrm{d}\hat{\mu}_B^n}\right|_{\substack{\mu_Q=0\\ \chi_1^S=0}}.$$

The net baryon density is given by:

$$c_1^B(T,\hat{\mu}_B) = \chi_1^B - \frac{\chi_{11}^{BS}}{\chi_2^S}\chi_1^S = \chi_1^B$$

c_1^B



This rescaling will break down at large $T \longrightarrow$ rescaling with SBL

c_1^B



This rescaling will break down at large $T \longrightarrow$ rescaling with SBL

Why does the rescaling work?

- It is an observation that it works
- It could be related to the critical scaling in the chiral limit
- If the universal contribution to EoS is large → single scaling variable
- If strength of transition is strongly Influenced by light quark masses → curves keep there shape
- Fits with the observation of constant width of the transition



[Borsanyi:2020fev]

Measuring the shift



c_1^B: net baryon density

$\overline{c_1^B}$: SBL of net baryon density

$$\Pi(T, \hat{\mu}_B, N_\tau) = \frac{T'(T, \hat{\mu}_B, N) - T}{T\hat{\mu}_B}$$

Measuring the shift



c_1^B: net baryon density

$\overline{c_1^B}$: SBL of net baryon density

$$\Pi(T, \hat{\mu}_B, N_\tau) = \frac{T'(T, \hat{\mu}_B, N) - T}{T\hat{\mu}_B}$$

Measuring the shift



c_1^B: net baryon density

$\overline{c_1^B}$: SBL of net baryon density

$$\Pi(T, \hat{\mu}_B, N_\tau) = \frac{T'(T, \hat{\mu}_B, N) - T}{T\hat{\mu}_B}$$

Measuring the shift



c_1^B: net baryon density

$\overline{c_1^B}$: SBL of net baryon density

$$\Pi(T, \hat{\mu}_B, N_\tau) = \frac{T'(T, \hat{\mu}_B, N) - T}{T\hat{\mu}_B}$$

Measuring the shift



c_1^B: net baryon density

$\overline{c_1^B}$: SBL of net baryon density

$$\Pi(T, \hat{\mu}_B, N_\tau) = \frac{T'(T, \hat{\mu}_B, N) - T}{T\hat{\mu}_B}$$

Measuring the shift



c_1^B: net baryon density

$\overline{c_1^B}$: SBL of net baryon density

$$\Pi(T, \hat{\mu}_B, N_\tau) = \frac{T'(T, \hat{\mu}_B, N) - T}{T\hat{\mu}_B}$$

Measuring the shift



c_1^B: net baryon density

$\overline{c_1^B}$: SBL of net baryon density

$$\Pi(T, \hat{\mu}_B, N_\tau) = \frac{T'(T, \hat{\mu}_B, N) - T}{T\hat{\mu}_B}$$

Measuring the shift



c_1^B: net baryon density

$\overline{c_1^B}$: SBL of net baryon density

$$\Pi(T, \hat{\mu}_B, N_\tau) = \frac{T'(T, \hat{\mu}_B, N) - T}{T\hat{\mu}_B}$$

Lattice Setup



- Action: tree-level Symanzik improved gauge action, with four times stout smeared staggered fermions
- 2+1+1 flavour, on LCP with pion and kaon mass
- Simulation at $\langle n_S \rangle = 0$
- Continuum estimate from lattice sizes: $32^3 \times 8$, $40^3 \times 10$, $48^3 \times 12$ and $64^3 \times 16$
- $\frac{\mu_B}{T} = \mathrm{i}\frac{j\pi}{8}$ with $j = 0, 3, 4, 5, (5.5), 6$ and 6.5
- Two methods of scale setting: f_π and w_0, $Lm_\pi > 4$

Systematic Errors

- 3 different sets of spline node points at μ_B=0
- 2 different sets of spline node points at finite imaginary μ_B
- w_0 or f_π based scale setting
- 2 different chemical potential ranges in the global fit: $\hat{\mu}_B \leq 5.5$ or $\hat{\mu}_B \leq 6.5$
- 2 functions for the chemical potential dependence of the global fit: linear or parabola
- including the coarsest lattice, $N_\tau = 8$, or not, in the continuum extrapolation.



histogram for λ_2 at $T = 155$ MeV

In total we perform 96 Fits. We weight every result with a $Q > 0.01$ uniformly

The expansion coefficients



$$\Pi(T, \hat{\mu}_B, N_\tau) = \frac{T'(T, \hat{\mu}_B, N) - T}{T\hat{\mu}_B}$$

$$\Pi(T, \hat{\mu}_B, N_\tau) = \lambda_2^A + \lambda_4^A \hat{\mu}_B^2 + \lambda_6^A \hat{\mu}_B^4 + \frac{1}{N_\tau^2}\left(\alpha^A + \beta^A \hat{\mu}_B^2 + \gamma^A \hat{\mu}_B^4\right)$$

We make a fit to calculate derivatives and constrain it with the HRG.

Results at $n_S = 0$ and $\mu_Q = 0$





More strangeness

Two more observables:





More strangeness

Two more expansion:





Beyond strangeness neutrality

$$\Delta\hat{\mu}_S \equiv \hat{\mu}_S - \hat{\mu}_S^\star,$$

the dimensionless strangeness and baryon densities become:

$$\chi_1^S(\hat{\mu}_S) \approx \chi_2^S(\hat{\mu}_S^\star)\Delta\hat{\mu}_S$$
$$\chi_1^B(\hat{\mu}_S) \approx \chi_1^B(\hat{\mu}_S^\star) + \chi_{11}^{BS}(\hat{\mu}_S^\star)\Delta\hat{\mu}_S,$$

where we only kept the linear leading order terms in $\Delta\hat{\mu}_S$. We will express thermodynamic quantities in terms of the strangeness-to-baryon fraction:

$$R = \frac{\chi_1^S}{\chi_1^B} = \frac{\chi_2^S(\hat{\mu}_S^\star)\Delta\hat{\mu}_S}{\chi_1^B(\hat{\mu}_S^\star)\Delta\hat{\mu}_S + \chi_{11}^{BS}(\hat{\mu}_S^\star)}.$$

Inverting this equation we get:

$$\Delta\hat{\mu}_S = \frac{R\hat{\chi}_1^B(\hat{\mu}_S^\star)}{\chi_2^S(\hat{\mu}_S^\star) - R\chi_{11}^{BS}(\hat{\mu}_S^\star)}.$$

Beyond strangeness neutrality



$$R = \frac{\chi_1^S}{\chi_1^B}$$

$$\Delta\hat{\mu}_S = \frac{R\hat{\chi}_1^B(\hat{\mu}_S^\star)}{\chi_2^S(\hat{\mu}_S^\star) - R\chi_{11}^{BS}(\hat{\mu}_S^\star)}$$

Strange Baryon density



Expanding the baryon density:

$$\frac{\chi_1^B(T,\hat{\mu}_B,R)}{\chi_1^B(T,\hat{\mu}_B,R=0)} \approx 1 + R\frac{\chi_{11}^{BS}(T,\hat{\mu}_B,R=0)}{\chi_2^S(T,\hat{\mu}_B,R=0)}$$

where all quantities on the right hand side are along the strangeness neutral line.

Strange Pressure



At the strangeness neutral line the $\mathcal{O}(R)$ correction of the pressure vanishes. The leading order correction gives:

$$\hat{p}(T, \hat{\mu}_B, R) \approx \hat{p}(T, \hat{\mu}_B, R) + \frac{1}{2}\frac{\mathrm{d}^2\hat{p}}{\mathrm{d}R^2}(T, \hat{\mu}_B)\, R^2,$$

where

$$\frac{\mathrm{d}^2\hat{p}}{\mathrm{d}R^2}(T, \hat{\mu}_B) = \frac{\left(\chi_1^B(T, \hat{\mu}_B)\right)^2}{\chi_2^S(T, \hat{\mu}_B)}.$$

Summary


μB = 0
μB = i π/8
μB = i 2π/8
μB = i 3π/8
μB = i 4π/8
μB = i 5π/8
μB = i 6π/8
T rescaled using κ=0.0205
T χ2B/μB (T)
T (1+κ (μB/T)2) [MeV]


lattice: 48³ × 12
Spline
measured difference
data
μ̂B = 4π/3
μ̂B = 4π/4
μ̂B = 4π/5
T/MeV


λ2 cont. est.
λ4 cont. est.
λ2 fit
λ4 fit
λ2 HRG
λ4 HRG
λn BB (T)
T [MeV]


μB/T = 3.5
μB/T = 3
μB/T = 2.5
μB/T = 2
μB/T = 1.5
μB/T = 1
μB/T = 0.5
μB/T = 0
nB/T3 (T)
T [MeV]


4stout, 48³ × 12
μ̂B = 0
μ̂B = i3π/8
μ̂B = i4π/8
μ̂B = i5π/8
μ̂B = i6π/8
λ2S
T[MeV]


R = 0.50
R = 0.25
R = 0.10
R = -0.10
R = -0.25
R = -0.50
μB/T = 2
nB/T3 (T)
T [MeV]

μ_Q


0.7
0.6
0.5
0.4
0.3
0.2
0.1
0
120
140
160
180
200
220
240
260
280
300
T [MeV]
d_2^B [n_S=0, n_Q=0.4n_B]
c_2^B [n_S=0, n_Q=0.5n_B]
T d d_2^B / d T [n_S=0, n_Q=0.4n_B]
T d c_2^B / d T [n_S=0, n_Q=0.5n_B]

κ vs. λ



Thermodynamics

$$\frac{p(T,\hat{\mu}_B)}{T^4} = \frac{p(T,0)}{T^4} + \int_0^{\hat{\mu}_B} c_1^B(T,\hat{\mu}_B')\mathrm{d}\hat{\mu}_B',$$

with

$$c_1^B(T,\hat{\mu}_B) = c_2^B(T',0)\frac{\overline{c_1^B}(\hat{\mu}_B)}{\overline{c_2^B}(0)},$$

and $\frac{p(T,0)}{T^4}$ from [Borsanyi:2013bia] The entropy density is defined as $s = \left.\frac{\partial p}{\partial T}\right|_{\mu_B,\mu_S}$, which can be rewritten in terms of dimensionless quantities as:

$$\hat{s} = 4\hat{p} + T\left.\frac{\partial \hat{p}}{\partial T}\right|_{\mu_B} = 4\hat{p} + T\left.\frac{\partial \hat{p}}{\partial T}\right|_{\hat{\mu}_B} - \hat{\mu}_B\chi_1^B,$$

where $\hat{s} \equiv \frac{s}{T^3}$ and we took into account the difference between derivatives at fixed μ_B versus at fixed $\hat{\mu}_B$.

Thermodynamics II

By noticing that on the strangeness neutral line

$$\frac{\mathrm{d}\hat{p}(T,\hat{\mu}_B,\hat{\mu}_S(T,\hat{\mu}_B))}{\mathrm{d}T} = \chi_1^S \frac{\partial \hat{\mu}_S}{\partial T} + \frac{\partial \hat{p}}{\partial T} = \frac{\partial \hat{p}(T,\hat{\mu}_B,\hat{\mu}_S(T,\hat{\mu}_B))}{\partial T},$$

we can write the logarithmic temperature derivative of the pressure as:

$$T\left.\frac{\partial \hat{p}(T,\hat{\mu}_B)}{\partial T}\right|_{\hat{\mu}} = T\frac{\partial \hat{p}(T,0)}{\partial T}$$

$$+\frac{1}{2}\int_0^{\hat{\mu}_B^2} T\left.\frac{\mathrm{d}c_2^B(T',0)}{\mathrm{d}T'}\right|_{T'} \times \left[1+\lambda_2^{BB}y+\lambda_4^{BB}y^2+T\left(\frac{\mathrm{d}\lambda_2^{BB}}{dT}y+\frac{\mathrm{d}\lambda_4^{BB}}{dT}y^2\right)\right]dy$$

where $\frac{\mathrm{d}c_2^B(T)}{\mathrm{d}T}$ is calculated at $\mu_B = 0$ and $T' = T\left(1+\lambda_2^{BB}y+\lambda_4^{BB}y^2\right)$
Given the pressure and the entropy, the dimensionless energy density is given by:

$$\hat{\epsilon} = \hat{s} - \hat{p} + \hat{\mu}_B \chi_1^B,$$

where $\hat{\epsilon} = \frac{\epsilon}{T^4}$.