



Contribution ID: 796

Type: Poster

Dilepton Signature of a First-Order Phase Transition

Wednesday 6 April 2022 19:10 (4 minutes)

Dileptons provide a unique way to access the properties of the fireball created in heavy-ion collisions. Since the dilepton yield directly depends on the duration of the fireball evolution, dileptons are valuable probes of the phase structure of strongly interacting matter. We study dilepton production in the SIS18 energy range by calculating and comparing the thermal dilepton emission for coarse grained microscopic transport models and hydrodynamical approaches with different equations of state, applying state-of-the-art in-medium spectral functions from hadronic many-body theory, chiral mixing in a chiral mean field model as well as from functional renormalization group methods. Presenting systematic comparisons of the space-time evolution of the fireball by means of various coarse-grained transport simulations and collision systems with respect to energy and participant number, we will show that the influence of the system size on the spectra is well under control and enhancements in heavy-ion collisions at SIS18 energies stem primarily from phase transitions. Supported by VH-NG-823, DFG CRC-TR 211 and GSI.

Author: WIEST, Maximilian**Co-authors:** STEINHEIMER, Jan; GALATYUK, Tetyana; RAPP, Ralf; STROTH, Joachim (Goethe-University and GSI); TRIPOLT, Ralf-Arno (University of Giessen); SECK, Florian (TU Darmstadt)**Presenter:** WIEST, Maximilian**Session Classification:** Poster Session 2 T05 / T13**Track Classification:** Electroweak probes