



Contribution ID: 249

Type: Oral presentation

Detailed study of Upsilon suppression with the measurement of the $\Upsilon(3S)$ meson in PbPb collisions at 5.02 TeV in CMS

Thursday, 7 April 2022 09:40 (20 minutes)

Because of the different binding energies, bottomonium mesons are particularly useful probes to understand the thermal properties of quark-gluon plasma. Previously, CMS observed the sequential suppression of $\Upsilon(1S)$, $\Upsilon(2S)$, and $\Upsilon(3S)$ in heavy ion (AA) collisions, which was widely accepted as evidence for the QGP formation. However, the $\Upsilon(3S)$ yield was excessively low, thus allowing us to report only statistical upper limits. In this talk, we present a detailed study of the measurement of excited bottomonium states with improved analysis technique and high-statistics data that enables us to observe the $\Upsilon(3S)$ meson in AA for the first time. The results are discussed together with the previous measurements in pPb collisions, which finally provides a full scan of all $\Upsilon(nS)$ states over the whole phase space.

Primary author: CMS

Presenter: LEE, Soohwan (Korea University (KR))

Session Classification: Parallel Session T11: Heavy flavors, quarkonia, and strangeness production

Track Classification: Heavy flavors, quarkonia, and strangeness production