QM 2022



Contribution ID: 92

Type: Poster

Measurement of long-range correlations and $v_2(\eta)$ in pp and p–Pb collisions with ALICE

Wednesday 6 April 2022 18:30 (4 minutes)

Measurements of long-range correlations are one of the important tools for evaluation of the interplay between the initial- and final-state effects. Recently, non-vanishing second Fourier harmonic v_2 has been observed in small collision systems such as pp and p–Pb collisions. In this contribution, we present measurements of v_2 extracted from long-range two-particle correlations for different charged particles multiplicities in pp at $\sqrt{s} = 13$ TeV and in p–Pb collisions at $\sqrt{s_{\rm NN}} = 5.02$ TeV.These measurements utilize the Forward Multiplicity Detector, which allows for unprecedented pseudorapidity ($\Delta \eta$) ranges to be explored (up to $\Delta \eta \sim 8$). We show the result of the η dependence of v_2 over a wide pseudorapidity range ($-3.1 < \eta < 4.8$). Results are compared with the AMPT and hydrodynamical calculations.

Author: Ms SEKIGUCHI, Yuko (University of Tokyo (JP))Presenter: Ms SEKIGUCHI, Yuko (University of Tokyo (JP))Session Classification: Poster Session 2 T07_2

Track Classification: Correlations and fluctuations