



Contribution ID: 459

Type: Poster

## Measurement of non-prompt $\Lambda_c^+$ production in pp collisions at $\sqrt{s} = 13$ TeV with ALICE

Friday 8 April 2022 14:04 (4 minutes)

### Measurement of non-prompt $\Lambda_c^+$ production in pp collisions at $\sqrt{s} = 13$ TeV with ALICE

Daniel Battistini for the ALICE Collaboration

In proton-proton (pp) collisions, the production of HF hadrons is typically described as a convolution of the parton distribution functions of the colliding protons, the partonic cross section, and the Fragmentation Functions (FFs). The latter describes the hadronisation of the heavy quarks in the different hadron species, and, since this process is non-perturbative, it is usually parametrised from measurements in  $e^+e^-$  collisions. However, recent studies by the ALICE Collaboration show that the ratio between the production of charm baryons with respect to mesons is significantly higher in hadronic collisions compared to  $e^+e^-$  interactions, invalidating the assumption that the FFs are independent of the collision system.

This contribution presents an extension of the studies on HF-baryon production in hadronic collisions to the beauty sector, via the measurement of the transverse-momentum-differential production cross section of  $\Lambda_c^+$ -baryon originating from beauty-hadron decays in pp collisions at  $\sqrt{s} = 13$  TeV. The measurement will also be compared to theoretical predictions based on fixed order plus next to the leading logarithm pQCD calculations folded with the beauty-hadron to  $\Lambda_c^+$  decay kinematics from PYTHIA8 simulations.

**Author:** BATTISTINI, Daniel (Universita e INFN Torino (IT))

**Presenter:** BATTISTINI, Daniel (Universita e INFN Torino (IT))

**Session Classification:** Poster Session 3 T11\_2

**Track Classification:** Heavy flavors, quarkonia, and strangeness production