

# 5th ComHEP: Colombian Meeting on High Energy Physics



## Report of Contributions

Contribution ID: 1

Type: **Regular Talk (15'+5')**

## Algebraic approach to scattering amplitudes

*Tuesday, 1 December 2020 09:35 (20 minutes)*

In this talk I will present our recent advances in the calculation of scattering amplitudes using strong homotopy algebras. This prescription inspired from string field theory allows to construct a generating function for all the tree-level scattering amplitudes. Several examples will be presented.

**Primary author:** LOPEZ ARCOS, Cristhiam

**Presenter:** LOPEZ ARCOS, Cristhiam

**Session Classification:** Formal theory

**Track Classification:** Formal theory

Contribution ID: 3

Type: **Regular Talk (15'+5')**

# Anisotropic Einstein Yang-Mills Higgs Dark Energy

*Thursday, 3 December 2020 15:40 (20 minutes)*

In the context of the dark energy scenario, the Einstein Yang-Mills Higgs model in the  $SO(3)$  representation was studied for the first time by M. Rinaldi (see JCAP **1510**, 023 (2015)) in a homogeneous and isotropic spacetime. We revisit this model, finding in particular that the interaction between the Higgs field and the gauge fields generates contributions to the momentum density, anisotropic stress and pressures, thus making the model inconsistent with the assumed background. We instead consider a homogeneous but anisotropic Bianchi-I spacetime background in this paper and analyze the corresponding dynamical behaviour of the system. We find that the only attractor point corresponds to an isotropic accelerated expansion dominated by the Higgs potential. However, the model predicts non-negligible anisotropic shear contributions nowadays, i.e. the current universe can have hair although it will lose it in the future. We investigate the evolution of the equation of state for dark energy and highlight some possible consequences of its behaviour related to the process of large-scale structure formation. As a supplement, we propose the “Higgs triad” as a possibility to make the Einstein Yang-Mills Higgs model be consistent with a homogeneous and isotropic spacetime.

**Primary authors:** ORJUELA-QUINTANA, JOHN BAYRON (UNIVERSIDAD DEL VALLE); VALENZUELA-TOLEDO, Cesar A. (Universidad del Valle); RODRIGUEZ GARCIA, Yeinzon (UAN & UIS (Colombia)); ÁLVAREZ, Miguel (Universidad Industrial de Santander)

**Presenter:** ORJUELA-QUINTANA, JOHN BAYRON (UNIVERSIDAD DEL VALLE)

**Session Classification:** Cosmology / Astroparticles

**Track Classification:** Cosmology / Astroparticles

Contribution ID: 5

Type: **Regular Talk (15'+5')**

## Prediction for Neutrino Masses, CP Violation, Leptogenesis and Neutrinoless Double Beta Decay from $\mathcal{T}_{13}$ Flavor Symmetry

*Thursday, 3 December 2020 10:55 (20 minutes)*

We propose an  $SU(5) \times \mathcal{T}_{13}$  model for both quarks and leptons. With tribimaximal (TBM) mixing for the seesaw matrix, symmetric Yukawa textures for down-quarks and charged leptons fail to explain the experimentally observed reactor angle. We derive the minimal asymmetric texture that not only reproduces the three lepton mixing angles assuming TBM mixing with a single phase, but also predicts the CP violating angle  $\delta_{CP} = 1.32\pi$ , consistent with the current global fit. We show that a straightforward origin of the “asymmetric texture” is traced back to the family symmetry  $\mathcal{T}_{13}$ , an order 39 discrete subgroup of  $SU(3)$ . With minimal number of familons having simple vacuum expectation values, the  $\mathcal{T}_{13}$  symmetry makes very definite prediction for the light neutrino masses of 28, 29 and 58 meV and neutrinoless double beta decay with  $|m_{\beta\beta}| = 25$  meV. We demonstrate that the CP violation predicted in the model can explain the baryon asymmetry of the universe through flavored leptogenesis with right-handed neutrino masses of  $\mathcal{O}(10^9 - 10^{12})$  GeV.

**Primary author:** RAHAT, Moinul Hossain (University of Florida)

**Presenter:** RAHAT, Moinul Hossain (University of Florida)

**Session Classification:** Neutrinos

**Track Classification:** Neutrinos

Contribution ID: 7

Type: **Short Talk (5')**

## Search for Long-Lived Heavy Neutrinos at the LHC with a VBF Trigger

*Friday, 4 December 2020 15:00 (5 minutes)*

The charged current production of long-lived heavy neutrinos at the LHC can use a prompt charged lepton for triggering the measurement of the process. However, in order to fully characterize the heavy neutrino interactions, it is necessary to also probe Higgs or  $Z$  mediated neutral current production. In this case the charged lepton is not available, so other means of triggering are required.

In this work, we explore the possibility of using a vector boson fusion trigger in the context of a GeV-scale Type I Seesaw model. We consider a minimal model, where both Higgs and  $Z$ -mediated contributions produce one heavy neutrino, as well as an extended model where the Higgs can decay into two heavy ones. Both scenarios are tested through displaced dilepton and displaced multitrack jet searches

**Primary author:** MASIAS TEVES, JOAQUIN (PUCP)

**Co-authors:** RUIZ, Jose (Universidad de Antioquia (CO)); Dr JONES-PEREZ, Joel (PUCP)

**Presenter:** MASIAS TEVES, JOAQUIN (PUCP)

**Session Classification:** Common Session

Contribution ID: 8

Type: **Regular Talk (15'+5')**

## Non-linear Regge trajectories with AdS/QCD

*Tuesday, 1 December 2020 09:55 (20 minutes)*

In this work, we consider a non-quadratic dilaton  $\Phi(z) = (\kappa z)^{2-\alpha}$  in the context of the static soft wall model to describe the mass spectrum of a wide range of vector mesons from the light up to the heavy sectors. The effect of this non-quadratic approach is translated into non-linear Regge trajectories with the generic form  $M^2 = a(n+b)^\nu$ . We apply this sort of fits for the isovector states of  $\omega$ ,  $\phi$ ,  $J/\psi$ , and  $\Upsilon$  mesons and compare with the corresponding holographic duals. We also extend these ideas to the heavy-light sector by using the isovector set of parameters to extrapolate the proper values of  $\kappa$  and  $\alpha$  through the average constituent mass  $\bar{m}$  for each mesonic specie considered. In the same direction, we address the description of possible non- $q\bar{q}$  candidates using  $\bar{m}$  as a holographic threshold, associated with the structure of the exotic state, to define the values of  $\kappa$  and  $\alpha$ . We study the  $\pi_1$  mesons in the light sector and the  $Z_c$ ,  $Y$ , and  $Z_b$  mesons in the heavy sector as possible exotic vector states. Finally, the RMS error for describing these twenty-seven states with fifteen parameters (four values for  $\kappa$  and  $\alpha$  respectively and seven values for  $\bar{m}$ ) is 12.61%.

**Primary authors:** MARTÍN CONTRERAS, Miguel Angel (Universidad de Valparaíso); VEGA, Alfredo

**Presenter:** MARTÍN CONTRERAS, Miguel Angel (Universidad de Valparaíso)

**Session Classification:** Formal theory

**Track Classification:** Formal theory

Contribution ID: 10

Type: **Regular Talk (15'+5')**

## Towards the one-loop galaxy bispectrum in the weak field approximation

*Thursday, 3 December 2020 16:35 (20 minutes)*

The large scale structure bispectrum contains information about the dynamics of the early universe in the squeezed limit since it couples large scales and short scales. Therefore, we compute the dark matter and galaxy bispectrum in a framework that is non-linear and based on General Relativity. To start, we use the weak-field approximation and standard perturbation theory to find the fluid evolution for dark matter density contrast to fourth-order in perturbations. Hereafter, we compute the galaxy density contrasts through the bias expansion. To do that, we evolve the Lagrangian bias expansion. We write the Lagrangian bias expansion in terms of operators built on the curvature of early time hypersurface of a comoving observer and the dark matter density contrast. Before computing the galaxy bispectrum, we properly renormalized the bias operators by adding the counter-terms that cancels the cut-off dependencies in the one-loop integrals. We find that in the squeezed limit, the non-linear relativistic corrections to the bispectrum are as large as the Newtonian ones and degenerates with the primordial non-gaussianity signal. Still, we have to add projection effects to our results. So, these are the first steps towards determining the gauge-invariant galaxy bispectrum.

**Primary authors:** CASTIBLANCO, Lina (Pontificia Universidad Católica de Valparaíso); NOREÑA, Jorge

**Presenter:** CASTIBLANCO, Lina (Pontificia Universidad Católica de Valparaíso)

**Session Classification:** Cosmology / Astroparticles

**Track Classification:** Cosmology / Astroparticles

Contribution ID: 11

Type: **Regular Talk (15'+5')**

## Noncommutative momentum and torsional regularization

*Tuesday, 1 December 2020 10:15 (20 minutes)*

We show that in the presence of the torsion tensor  $S^k_{ij}$ , the quantum commutation relation for the four-momentum, traced over spinor indices, is given by  $[p_i, p_j] = 2i\hbar S^k_{ij} p_k$ .

In the Einstein–Cartan theory of gravity, in which torsion is coupled to spin of fermions, this relation in a coordinate frame reduces to a commutation relation of noncommutative momentum space,  $[p_i, p_j] = i\epsilon_{ijk} U p^3 p_k$ , where  $U$  is a constant on the order of the squared inverse of the Planck mass.

We propose that this relation replaces the integration in the momentum space in Feynman diagrams with the summation over the discrete momentum eigenvalues.

We derive a prescription for this summation that agrees with convergent integrals:

$$\int \frac{d^4 p}{(p^2 + \Delta)^s} \rightarrow 4\pi U^{s-2} \sum_{l=1}^{\infty} \int_0^{\pi/2} d\phi \frac{\sin^4 \phi n^{s-3}}{[\sin \phi + U \Delta n]^s},$$

where  $n = \sqrt{l(l+1)}$  and  $\Delta$  does not depend on  $p$ .

We show that this prescription regularizes ultraviolet-divergent integrals in loop diagrams.

We extend this prescription to tensor integrals.

We derive a finite, gauge-invariant vacuum polarization tensor and a finite running coupling.

Including loops from all charged fermions, we find a finite value for the bare electric charge of an electron:  $\approx -1.22 e$ .

This torsional regularization may therefore provide a realistic, physical mechanism for eliminating infinities in quantum field theory and making renormalization finite.

**Primary author:** POPLAWSKI, Nikodem

**Presenter:** POPLAWSKI, Nikodem

**Session Classification:** Formal theory

**Track Classification:** Formal theory



Contribution ID: 16

Type: **Regular Talk (15'+5')**

## On the very nature of neutrinos: the inverse beta-decay as a test bench

*Thursday, 3 December 2020 10:35 (20 minutes)*

The lifetime of a particle is usually considered as one of its inherent features. On one hand, the lifetime of non-elementary quantum objects like pions or muons can be calculated from the knowledge of the interaction which rules their decay. On the other hand, other particles such as the electron are regarded as stable, in the sense that they do not decay at all (at least within Standard Model framework). In spite of this, decay properties are less fundamental than commonly thought, since they depend on the reference frame where they are evaluated. Along this line, the decay of accelerated protons was analyzed both in the laboratory frame (where the proton is accelerated) and in the comoving frame (where the proton is at rest and interacts with the Unruh thermal bath of electrons and neutrinos). The equality between the two rates was exhibited as a theoretical proof of the necessity of Unruh effect for the consistency of QFT. Recently, this formalism was refined, embedding the phenomenon of mixing for the emitted neutrinos. However, the question of whether to consider mass or flavor states for asymptotic neutrinos remained open. Here, we show that the only scenario compatible with: i) the general covariance of QFT, ii) the description of the phenomenologically observed neutrino oscillations, iii) CP-violation in neutrino oscillations is the one build upon flavor eigenstates. We further point out that Unruh radiation must be made up of oscillating neutrinos. For this purpose, we exploit previous results showing that Unruh bath for mixed fields loses its characteristic thermal spectrum. Possible experimental implications are investigated in connection with recent neutrino experiments such as *Katrin* and *Ptolemy*.

**Primary author:** LUCIANO, Giuseppe Gaetano (INFN - National Institute for Nuclear Physics)

**Presenter:** LUCIANO, Giuseppe Gaetano (INFN - National Institute for Nuclear Physics)

**Session Classification:** Neutrinos

**Track Classification:** Neutrinos

Contribution ID: 19

Type: **Regular Talk (15'+5')**

## Explaining muon $g-2$ anomaly in a non-universal $U(1)_X$ extended SUSY theory

*Monday, 30 November 2020 15:20 (20 minutes)*

It is presented a non-universal  $U(1)_X$  extension to the Standard Model with two scalar doublets, two scalar singlets, three additional quark singlets, two lepton singlets and three generations of right-handed and Majorana neutrinos is made to explain lepton mass hierarchy and muon anomalous magnetic moment in a chiral anomaly free framework. Besides, neutrino mass generation is achieved by implementing an inverse seesaw mechanism and it is found that the lightest fermions are tree-level massless but massive at one-loop level. Finally, the conditions that recreate the PMNS matrix and the Higgs boson as the lightests scalar are obtained to make a numerical approach to muon  $g - 2$ . It is found that only contributions due to exotic neutrinos interacting with charged scalars are relevant, though they are negative. Nevertheless, they might explain the anomaly depending on the order of magnitude of  $v_\chi$  which is achieved in the supersymmetric scenario when the contributions of exotic neutrinos interacting with  $W$  gauge bosons are positive and no longer negligible..

**Primary authors:** ALVARADO GALEANO, Juan Sebastian (Universidad Nacional de Colombia); Prof. MARTÍNEZ MARTÍNEZ , Roberto Enrique (Universidad Nacional de Colombia)

**Presenter:** ALVARADO GALEANO, Juan Sebastian (Universidad Nacional de Colombia)

**Session Classification:** Formal theory

**Track Classification:** Higgs / Standard model

Contribution ID: 20

Type: **Regular Talk (15'+5')**

## Quantum Machine Learning concepts for HEP

*Monday, 30 November 2020 16:00 (20 minutes)*

This contribution explores the Machine Learning techniques and Quantum Computing concepts and applications in High Energy Physics considering a phenomenological and theoretical view. Besides, we show the main tools to explore the Standard Model extensions, decay process and the parameter space. With this set of tools, we want to explore the bounds and define exclusion regions, those bounds might be relevant to explore in the next generation of colliders and could be tested in order to understanding of phenomena.

**Primary authors:** Prof. ORDUZ-DUCUARA, Javier (UNAM); Ms IYER, Vijayasri (Department of Information Technology)

**Presenter:** Ms IYER, Vijayasri (Department of Information Technology)

**Session Classification:** Formal theory

**Track Classification:** Higgs / Standard model

Contribution ID: 23

Type: **Regular Talk (15'+5')**

## Effective Theory of Freeze-in Dark Matter

*Wednesday, 2 December 2020 15:20 (20 minutes)*

We perform a model independent study of *freeze-in* of massive particle dark matter (DM) by adopting an effective field theory framework. Considering the dark matter to be a gauge singlet Majorana fermion, odd under a stabilising symmetry  $Z_2$  under which all standard model (SM) fields are even, we write down all possible DM-SM operators upto and including mass dimension eight. For simplicity of the numerical analysis we restrict ourselves only to the scalar operators in SM as well as in the dark sector. We calculate the DM abundance for each such dimension of operator considering both *UV and IR freeze-in* contributions which can arise before and after the electroweak symmetry breaking respectively. After constraining the cut-off scale and reheat temperature of the universe from the requirement of correct DM relic abundance, we also study the possibility of connecting the origin of *neutrino mass* to the same cut-off scale by virtue of lepton number violating Weinberg operators. We thus compare the bounds on such cut-off scale and corresponding reheat temperature required for UV freeze-in from the origin of light neutrino mass as well as from the requirement of correct DM relic abundance. We also briefly comment upon the possibilities of realising such DM-SM effective operators in a UV complete model.

**Primary authors:** BARMAN, Basabendu; Dr BORAH, Debasish (Indian Institute of Technology Guwahati); ROSHAN, Rishav (Indian Institute of Technology Guwahati)

**Presenter:** BARMAN, Basabendu

**Session Classification:** Dark matter

**Track Classification:** Dark matter

Contribution ID: 24

Type: **Regular Talk (15'+5')**

## Dark Matter in the Time of Primordial Black Holes

*Wednesday, 2 December 2020 17:55 (20 minutes)*

The evaporation of primordial black holes (PBH) with masses ranging from  $\sim 10^{-1}$  to  $\sim 10^9$  g could have generated the whole observed dark matter (DM) relic density. It is typically assumed that after being produced, its abundance freezes and remains constant. However, thermalization and number-changing processes in the dark sector can have a strong impact, in particular enhancing the DM population by several orders of magnitude. Here we estimate the boost from general arguments such as the conservation of energy and entropy, independently from the underlying particle physics details of the dark sector. Two main consequences can be highlighted: i) As the DM abundance is increased, a smaller initial energy density of PBHs is required. ii) Thermalization in the dark sector decreases the mean DM kinetic energy, relaxing the bound from structure formation and hence, allowing light DM with mass in the keV ballpark.

**Primary authors:** BERNAL, Nicolás (Universidad Antonio Nariño); ALBERTO ZAPATA NOREÑA, Oscar

**Presenter:** BERNAL, Nicolás (Universidad Antonio Nariño)

**Session Classification:** Dark matter

**Track Classification:** Dark matter

Contribution ID: 25

Type: **Regular Talk (15'+5')**

## Light pseudoscalar and axial meson spectroscopy via an AdS/QCD modified soft wall model

*Friday, 4 December 2020 10:35 (20 minutes)*

We describe the mass spectrum of light pseudoscalars and axial mesons using a modified softwall model with an UV cutoff. These mesons are included using an anomalous dimension that shift the conformal dimension of the non-interacting bulk fields. Using the extra UV cutoff approach, we can fit six eta and six a1 states organized in radial trajectories with an error close to 21.1%. We also confirm that chiral symmetry is restored in this model after checking that highly excited rho and a1 states become degenerate.

**Primary authors:** CORTES, Santiago (Universidad de los Andes); Dr MARTIN CONTRERAS, Miguel Angel (Universidad de Valparaíso); Prof. VEGA, Alfredo (Universidad de Valparaíso)

**Presenter:** CORTES, Santiago (Universidad de los Andes)

**Session Classification:** Heavy flavour

**Track Classification:** Heavy flavour

Contribution ID: 26

Type: **Regular Talk (15'+5')**

## Probing the properties of relic neutrinos using the cosmic microwave background, the Hubble Space Telescope and galaxy clusters

*Tuesday, 1 December 2020 16:55 (20 minutes)*

We investigate the observational constraints on the cosmic neutrino background (CNB) given by the extended  $\Lambda$ CDM scenario ( $\Lambda$ CDM +  $N_{\text{eff}}$  +  $\sum m_\nu$  +  $c_{\text{eff}}^2$  +  $c_{\text{vis}}^2$  +  $\xi_\nu$ ) using the latest observational data from *Planck* CMB (temperature power spectrum, low-polarisation and lensing reconstruction), baryon acoustic oscillations (BAOs), the new recent local value of the Hubble constant from *Hubble Space Telescope* (*HST*) and information of the abundance of galaxy clusters (GCs). We study the constraints on the CNB background using CMB + BAO + *HST* data with and without the GC data. We find  $\Delta N_{\text{eff}} = 0.614 \pm 0.26$  at 68 per cent confidence level when the GC data are added in the analysis. We do not find significant deviation for sound speed in the CNB rest frame. We also analyze the particular case  $\Lambda$ CDM +  $N_{\text{eff}}$  +  $\sum m_\nu$  +  $\xi_\nu$  with the observational data. Within this scenario, we find  $\Delta N_{\text{eff}} = 0.60 \pm 0.28$  at 68 per cent confidence level. In both the scenarios, no mean deviations are found for the degeneracy parameter.

**Primary author:** BONILLA RIVERA, Alexander (UFJF)**Presenter:** BONILLA RIVERA, Alexander (UFJF)**Session Classification:** Neutrinos**Track Classification:** Neutrinos

Contribution ID: 27

Type: **Regular Talk (15'+5')**

## Flavored axions and the flavor problem

*Friday, 4 December 2020 10:55 (20 minutes)*

A Peccei-Quinn (PQ) symmetry is proposed, in order to generate in the Standard Model (SM) quark sector a realistic mass matrix ansatz with five texture-zeros. Limiting our analysis to Hermitian mass matrices we show that this requires a minimum of 4 Higgs doublets. This model allows assigning values close to 1 for several Yukawa couplings, giving insight into the origin of the mass scales in the SM. Since the PQ charges are non-universal the model features Flavor-Changing Neutral Currents (FCNC) at the tree level. We calculate the FCNC couplings of the most general low-energy effective Lagrangian for the axion in a procedure valid for an arbitrary number of Higgs doublets. Finally, we report the allowed region in the parameter space obtained from the measurements of branching ratios of semileptonic meson decays.

**Primary authors:** ROJAS, Eduardo (Universidad de Nariño); MARTINEZ, Roberto (Universidad Nacional de Colombia); Prof. GIRALDO, Yithsbey (Universidad de Nariño); Prof. SALAZAR, Juan Carlos (Universidad de Nariño)

**Presenter:** ROJAS, Eduardo (Universidad de Nariño)

**Session Classification:** Heavy flavour

**Track Classification:** Heavy flavour



Contribution ID: 28

Type: **Regular Talk (15'+5')**

## A Guidance for Building Dark Energy Models

We discuss in this talk several physical principles as well as emerging observational constraints, as the speed of gravitational waves, that any dark energy model has to face in order to ensure its cosmological viability. In particular dark energy models based on modified gravity theories find a major challenge for addressing the late-time dynamics in a consistent way due to the presence of instabilities. We present a guidance for building dark energy models with some necessary conditions that must be fulfilled, and illustrate this point showing some simple but suitable examples.

**Primary author:** GABRIEL , Gómez (Departamento de Física, Universidad de Santiago de Chile)

**Presenter:** GABRIEL , Gómez (Departamento de Física, Universidad de Santiago de Chile)

**Session Classification:** Cosmology / Astroparticles

**Track Classification:** Cosmology / Astroparticles

Contribution ID: 29

Type: **Regular Talk (15'+5')**

## Diophantine scotogenic models

*Wednesday, 2 December 2020 17:15 (20 minutes)*

We propose a general method to find anomaly free Abelian standard model extensions with radiative Dirac neutrino masses and DM matter candidates

**Primary author:** Prof. RESTREPO, Diego (Universidad de Antioquia)

**Presenter:** Prof. RESTREPO, Diego (Universidad de Antioquia)

**Session Classification:** Dark matter

**Track Classification:** Dark matter

Contribution ID: 30

Type: **Regular Talk (15'+5')**

## External Momentum Dependence from the Higgs Boson Mass

*Monday, 30 November 2020 15:40 (20 minutes)*

The three-loop corrections to the Higgs boson mass are usually computed in the on-shell scheme using the effective potential approach, where the external momentum are put equal to zero. In this presentation we are going to discuss the technical details for the Feynman diagrammatic computation of the external momentum dependence coming from the three-loop self-energy corrections to the Higgs boson mass in both the Standard Model (SM) as in the minimal supersymmetric standard model with real parameters (rMSSM).

**Primary author:** REYES ROJAS, Edilson Alfonso (UNAL, DESY)

**Presenter:** REYES ROJAS, Edilson Alfonso (UNAL, DESY)

**Session Classification:** Formal theory

**Track Classification:** Higgs / Standard model

Contribution ID: 31

Type: **Short Talk (5')**

## Three level FCNC from Models with a flavored Peccei-Quinn Symmetry

*Friday, 4 December 2020 15:50 (5 minutes)*

A Peccei-Quinn (PQ) symmetry is proposed, in order to generate a realistic mass matrix ansatz with five texture-zeros for both quark and lepton sector in the Standard Model (SM). Limiting our analysis to Hermitian mass matrices we show that this requires a minimum of 4 Higgs doublets. The price we pay is to have Yukawa values in the lepton sector much lower than 1, but consistent with the experimental values. Since the PQ charges are non-universal the model features Flavor-Changing Neutral Currents (FCNC) at the tree level. We calculate the FCNC couplings of the most general low-energy effective Lagrangian for the axion in a procedure valid for an arbitrary number of Higgs doublets. Finally, we report the allowed region in the parameter space obtained from the measurements of branching ratios of semileptonic meson decays.

**Primary authors:** Prof. GIRALDO, Yithsbey (Universidad de Nariño); ROJAS, Eduardo (Universidad de Nariño); Dr MARTÍNEZ, Roberto (Universidad Nacional de Colombia); Dr SALAZAR, Juan Carlos (Universidad de Nariño)

**Presenter:** Prof. GIRALDO, Yithsbey (Universidad de Nariño)

**Session Classification:** Common Session

Contribution ID: 33

Type: **Regular Talk (15'+5')**

## Addressing the $R(D)$ and $R(D^*)$ anomalies within a charged scalar boson scenario

*Friday, 4 December 2020 11:15 (20 minutes)*

The experimental measurements collected by the BABAR, Belle and LHCb experiments on the  $R(D)$  and  $R(D^*)$  observables indicate the existence of disagreement with respect to the Standard Model (SM) predictions. This discrepancy has been referred to as the “ $R(D)$  and  $R(D^*)$  anomalies”, and consequently, a large number of studies considering the effects of new physics (NP) beyond MS have been discussed in the literature. In this talk, we will present a review of the current state (experimental and theoretical), and a possible scenario to address the  $R(D)$  and  $R(D^*)$  anomalies through a charged scalar boson will be discussed. We will show the recent results of the phenomenological study to the parameter space allowed by these NP models. We will discuss the implications of parametric space that would be obtained from the projections of the Belle II experiment.

**Primary author:** QUINTERO, Nestor (Universidad Santiago de Cali)

**Co-authors:** MUÑOZ, J. H. (Universidad del Tolima); CARDOZO, J. (Universidad del Tolima); ROJAS, E. (Universidad de Nariño)

**Presenter:** QUINTERO, Nestor (Universidad Santiago de Cali)

**Session Classification:** Heavy flavour

**Track Classification:** Heavy flavour

Contribution ID: 34

Type: **Regular Talk (15'+5')**

## Generation of Carroll-Field-Jackiw term in Horava-Lifshitz $z=3$ CPT-violating QED

Tuesday, 1 December 2020 10:35 (20 minutes)

Theories with anisotropy between space and time, also called Horava-Lifshitz (HL) theories, has been studied in many works since the seminal paper of P. Horava was published, where the Lifshitz scaling is used to construct a power counting renormalizable gravity model. The Lifshitz scaling is described by the transformation  $x \rightarrow bx$ ,  $t \rightarrow b^z t$ , where the integer  $z$  is so-called critical exponent. In the case of quantum gravity, if  $z = 3$ , the dimension of gravitational constants turns out to be zero, a necessary condition for the renormalizability. Then, other theories started to be studied within the Horava-Lifshitz context, for example, theories with four fermion interaction and spinor QED.

An interesting problem to be studied is the HL-like theory where CPT symmetry is broken and the calculating the generation of a Carroll-Field-Jackiw (CFJ) term. The aim of this study could be the search for a generalized description of Lorentz breaking theories involving HL-like theories and where the 'usual' Lorentz-breaking theories as particular cases.

We start with the action of HL  $z = 3$  CPT-violating QED where a Lorentz and CPT violating term is included in the fermionic sector

$$S = \int d^4x [\bar{\psi}(i\not{D}_0 + (i\not{D}_i)^3 - m^3 - b_0\gamma_5 + (b\not{D}\not{D})_i\gamma_5)\psi],$$

where  $\not{D}_0 = D_0\gamma^0$ ,  $\not{D}_i = D_i\gamma^i$ ,  $b_0 = b_0\gamma^0$ , and  $(b\not{D}\not{D})_i = (bDD)_{ijk}\gamma^i\gamma^j\gamma^k$ , with  $D_{0,i} = \partial_{0,i} + ieA_{0,i}$ .

To calculate the generated CFJ term, first we use the simplest prescription for Dirac matrices, where we find that the CFJ term is finite and reproduces one of the values obtained in the usual Lorentz-breaking QED

$$\mathcal{L}_{CFJ} = -\frac{e^2}{4\pi^2} b_\kappa \epsilon^{\kappa\lambda\mu\nu} A_\lambda \partial_\mu A_\nu.$$

Using the 't Hooft and Veltman prescription, we find that the CFJ term is not generated.

$$\mathcal{L}_{CFJ} = 0$$

So, we can conclude that the generated CFJ term is finite, and gauge and Lorentz invariant, but indetermined because of its dependence on the calculation scheme. This result is natural, because the chiral anomaly is related with the ambiguity of the triangle graph, and since the chiral anomalies also exist in  $z = 3$  HL-QED, the presence of ambiguity in the triangle graph in our theory seems to be a direct analogy of the situation taking place in the usual Lorentz-breaking QED.

**Primary authors:** Mr MARTÍNEZ, Ricardo (Universidade Federal de Alagoas); Dr MARIZ, Tiago (Universidade Federal de Alagoas); Dr NASCIMENTO, José Roberto (Universidade Federal da Paraíba); Dr PETROV, Albert (Universidade Federal da Paraíba)

**Presenter:** Mr MARTÍNEZ, Ricardo (Universidade Federal de Alagoas)

**Session Classification:** Formal theory

**Track Classification:** Formal theory

Contribution ID: 35

Type: **Short Talk (5')**

## Matriz de mezcla de neutrinos en el marco de la simetría $\mu - \tau$

*Friday, 4 December 2020 15:45 (5 minutes)*

Mientras que experimentos de oscilaciones de neutrinos han dado evidencia de que los neutrinos tienen masa y se mezclan, aún no se tiene una teoría que explique la estructura de estas mezclas, plasmada en la conocida matriz PMNS. Antes de la medida del ángulo de reactor ( $\theta_{13}$ ), la matriz PMNS presentaba una simetría de intercambio  $\mu$ - $\tau$  consistente con un valor de cero para  $\theta_{13}$  y para la fase de Dirac de violación CP. Después de la medida de gran precisión del ángulo  $\theta_{13}$  se ha explorado la posibilidad de que la matriz PMNS surja de la perturbación de dicha simetría. Un caso especial de dicha simetría es el esquema TBM (Tri-Bi-Maximal), la cual establece  $\sin^2 \theta_{12} = 1/3$ . En este trabajo se utilizan diferentes desviaciones del patrón de mezcla TBM, mediante rotaciones unitarias y ortogonales, para estudiar la violación de CP, mostrando que es posible generar predicciones para las fases de Majorana y la fase de Dirac, estudiando sus relaciones con los parámetros de mezcla. Adicionalmente se estudian los efectos que tienen estas predicciones sobre el observable  $|m_{ee}|$  relevante en el decaimiento doble beta sin neutrinos.

**Primary authors:** CARDONA, David; Dr RIVERA AGUDELO, Diana Carolina (Universidad Santiago de Cali)

**Presenter:** CARDONA, David

**Session Classification:** Common Session

Contribution ID: 36

Type: **Regular Talk (15'+5')**

## Production and properties of the $B_c(2S)$ and $B^*c(2S)$ states

*Friday, 4 December 2020 10:15 (20 minutes)*

We reported the production and properties studies of the  $B_c(2S)$  and  $B^*c(2S)$  states, based on an event sample of pp collisions at a center-of-mass energy of 13 TeV, collected by CMS detector and corresponding to an integrated luminosity of  $143 \text{ fb}^{-1}$ . The excited states are reconstructed in the  $B_c\pi+\pi$ -decay. The production rate of  $B_c(2S)$  and  $B^*c(2S)$  respect to the base state  $B_c$  as estimated within the acceptance of the detector. The relative rate of the two mesons respect to each other is also estimated as well as the shape of the invariant mass of the pion-pion system.

**Primary author:** RODRIGUEZ GIRALDO, Manuel Alejandro (Universidad de Antioquia (CO))

**Co-author:** MEJIA GUISAO, Jhovanny Andres (Universidad de Antioquia (CO))

**Presenter:** RODRIGUEZ GIRALDO, Manuel Alejandro (Universidad de Antioquia (CO))

**Session Classification:** Heavy flavour

**Track Classification:** Heavy flavour



Contribution ID: 38

Type: **Regular Talk (15'+5')**

## Anisotropic solid dark energy

*Thursday, 3 December 2020 16:00 (20 minutes)*

In this work, we study a triplet of inhomogeneous scalar fields, known as “solid”, as a source of anisotropic dark energy. By using a dynamical system approach, we find that anisotropic accelerated solutions can be realized as attractor points for suitable parameters of the model. We complement the dynamical analysis with a numerical solution whose initial conditions are set in the deep radiation epoch. The model predicts a spatial shear within the observational bounds nowadays, even when it is set to zero as an initial condition. The hairy attractors and an ultra slowly varying equation of state of dark energy close to -1 are key features of this scenario. We also analyzed the isotropic limit of the model, finding that the solid can be described by a constant equation of state and thus being able to simulate the behaviour of a cosmological constant.

**Primary authors:** MOTOA MANZANO, Josue (Universidad del valle); VALENZUELA-TOLEDO, Cesar A. (Universidad del Valle); ORJUELA-QUINTANA, JOHN BAYRON (UNIVERSIDAD DEL VALLE); Prof. PEREIRA, Thiago (Universidad estadual de Londrina)

**Presenter:** MOTOA MANZANO, Josue (Universidad del valle)

**Session Classification:** Cosmology / Astroparticles

**Track Classification:** Cosmology / Astroparticles

Contribution ID: 39

Type: **Short Talk (5')**

## K-matrix formalism in light-meson spectroscopy

*Friday, 4 December 2020 15:35 (5 minutes)*

In this work we study the  $K$ -matrix formalism and show how this can be applied to Dalitz plot analyses of charm-meson decays, such as  $D^0 \rightarrow K_s^0 \pi^+ \pi^-$ . The  $K$ -matrix, in contrast to the typical Isobar Model (IM), allows to properly include in the decay amplitude broad-overlapping resonances and non-resonant background, which are features of some of the known experimental results in light-meson spectroscopy. The  $\pi\pi$  S-wave scattering presented in this work is a good example of this underlying dynamics. Moreover, the  $K$ -matrix formalism imposes by construction a unitarity constraint, which is not ensured by other approaches as IM.

**Primary author:** ORDONEZ, Sebastian (National University of Colombia)

**Co-author:** Prof. MILANÉS, Diego (Universidad Nacional de Colombia)

**Presenters:** ORDONEZ, Sebastian (National University of Colombia); Prof. MILANÉS, Diego (Universidad Nacional de Colombia)

**Session Classification:** Common Session

Contribution ID: 40

Type: **Short Talk (5')**

## Correlación Energía-Energía en Electrodinámica Cuántica Supersimétrica

*Friday, 4 December 2020 15:55 (5 minutes)*

### Resumen:

La física moderna ha entrado en una nueva era de precisión donde los observables físicos, como las masas de las partículas o la correlación energía-energía de átomos hidrogenoides, deben ser medidos experimentalmente y calculados teóricamente con una mayor precisión. La búsqueda de nuevos grados de libertad físicos es también una prioridad en la agenda de los aceleradores de partículas modernos como el LHC, el ILC y el FCC. En particular, la supersimetría es el escenario de nueva física con mayor fenomenología en la actualidad y recibirá gran parte de la atención en los experimentos futuros. En nuestro proyecto vamos a estudiar las implicaciones de las correcciones provenientes de la supersimetría a la correlación energía-energía en el sistema de dos partículas electrón-positrón, con el fin de dar una descripción alternativa de la cuantización de la radiación electromagnética en la electrodinámica cuántica y establecer un esquema de renormalización que permita encontrar restricciones fenomenológicas al valor numérico de la escala de nueva física, donde se espera que los nuevos grados de libertad aparezcan.

**Primary authors:** GAMBOA, Alexander; REYES ROJAS, Edilson Alfonso (UNAL, DESY)

**Presenter:** GAMBOA, Alexander

**Session Classification:** Common Session

Contribution ID: 43

Type: **Regular Talk (15'+5')**

# Theory of Fast Flavor Conversion for Supernova neutrinos

*Thursday, 3 December 2020 11:15 (20 minutes)*

We all know that in the dense anisotropic interior of the star, neutrino- neutrino forward-scattering can lead to fast collective neutrino oscillations, which has striking consequences on flavor dependent neutrino emission and can be crucial for the evolution of a supernova and its neutrino signal. Although the triggering and initial growth of fast oscillations are understood, owing to its complicated nonlinear evolution, the final impact is not yet known. Interestingly, stellar explosion and the neutrino signal are sensitive to the processed flavor-dependent fluxes, but the required neutrino theory prediction is still lacking. In my talk I will address this crucial theoretical and phenomenological obstacle and present a theory of fast flavor conversions that will explain how, when and to what extent do the flavor differences change. Finally, I will give a method and a simple formula for computing the final fluxes that can be a crucial input for supernova theory and neutrino phenomenology.

**Primary author:** BHATTACHARYYA, Soumya (TIFR)**Presenter:** BHATTACHARYYA, Soumya (TIFR)**Session Classification:** Neutrinos**Track Classification:** Neutrinos

Contribution ID: 45

Type: **Regular Talk (15'+5')**

## The Z5 model of two-component dark matter

*Wednesday, 2 December 2020 15:40 (20 minutes)*

In this talk we present the phenomenology of the Z5 model for two-component dark matter. This model, which can be seen as an extension of the well-known singlet scalar model, features two complex scalar fields—the dark matter particles—that are Standard Model singlets but have different charges under a Z5 symmetry. The interactions allowed by the Z5 give rise to novel processes between the dark matter particles that affect their relic densities and their detection prospects. Dark matter masses below the TeV are still compatible with present data, and current and future direct detection experiments may be sensitive to signals from both dark matter particles.

**Primary authors:** ZAPATA, Oscar (Universidad de Antioquia); E. YAGUNA, Carlos

**Presenter:** ZAPATA, Oscar (Universidad de Antioquia)

**Session Classification:** Dark matter

**Track Classification:** Dark matter

Contribution ID: 46

Type: **Regular Talk (15'+5')**

## Two component dark matter model in light of Beam Dump experiments.

*Wednesday, 2 December 2020 16:15 (20 minutes)*

We study a scalar dark matter (DM) model with two DM species coupled to standard model (SM) particles via a sub-GeV dark photon. The two DM candidates can be produced at fixed-target experiment a la Beam-Dump. Predictions for signal and backgrounds are obtained with the help of MadDump and NuWro Montecarlo generators. We explore the potential reach on the sensitivity of DUNE near detector and SHiP experiment and analyze the constraints coming from current limits on DM observables such as relic density and DM conversion in the early universe.

**Primary authors:** Mr PALACIO, Guillermo (Universidad EIA); Dr BETANCUR, Amalia (Universidad EIA); Dr CASTILLO RAMIREZ, Andrés Fernando (Universidad Sergio Arboleda); Dr SUAREZ, Juan Guillermo (Universidad EIA)

**Presenter:** Mr PALACIO, Guillermo (Universidad EIA)

**Session Classification:** Dark matter

**Track Classification:** Dark matter

Contribution ID: 47

Type: **Regular Talk (15'+5')**

## Multicomponent dark matter and the IDM

*Wednesday, 2 December 2020 17:35 (20 minutes)*

We present a multicomponent dark sector model, comprised of a fermionic and a scalar dark matter candidate. In the scalar sector, the dark matter candidate is the Inert Doublet Model, while the fermionic candidate is a mix of fields in different representations of the  $SU(2)_L$  group. We present the results from imposing restrictions on relic density, direct detection and other observables.

**Primary authors:** Dr RIVERA ROMERO, Andrés Felipe (Universidad de Antioquia); PALACIO, Guillermo; BETANCUR RODRÍGUEZ, Amalia

**Presenter:** BETANCUR RODRÍGUEZ, Amalia

**Session Classification:** Dark matter

**Track Classification:** Dark matter

Contribution ID: 48

Type: Short Talk (5')

## Search for new physics in the final state of $\tau, b, \nu$

Friday, 4 December 2020 15:05 (5 minutes)

The Standard model (SM) is one of the most successful physics theory, however it presents disagreements between some predictions and the observations. One of them has been studied in the last decade and it's related to the lepton universality. The  $R_{D^*}$  Anomaly appear to be enhanced respect to the standard model by roughly 30% with a global significance of  $\approx 4\sigma$ . This has been studied by BaBar, Belle and LHC-b experiments. As the measure it's close to the traditional  $5\sigma$  deviations, it could be a sign of new physics. Nevertheless an approaching to the problem considering the  $b$  quark and the  $\tau$  in the final state has never been measured. The aim of this project is discarding values from the mass parameter space for extensions to the SM as the  $W'$  model.

The standard model is the ruling theory for the composition and interaction of the known universe so far. It's composed by different type of particles, the quarks and leptons, each one with 3 generations, then we have the gauge bosons, responsible for the electric, weak and strong interactions, and finally the Higgs boson, responsible of giving mass to the rest of the massive particles from the SM. However, it's far from being a definitive theory, with problems as the Strong CP problem, and the neutrino masses, it is still in tests in experiments as the Tevatron and the LHC. One of this problem is associated with particles from the third generation of quarks and leptons, punctually with the  $b$  quarks, and the  $\tau$  leptons. \\

The  $R_{D^*}$  is the ratio between the branching fraction of  $B$  decaying semileptonically to a  $\tau$  with respect to the decay to other charged leptons, i.e:

$$R_{D^*} = \frac{\Gamma(B \rightarrow D^* \tau \nu)}{\Gamma(B \rightarrow D^* l \nu)}$$

With  $l = e$  or  $\mu$  \cite{greljo2019mono}. \\ \\

This measure appears to be enhanced respect to the standard model by roughly 30% with a global significance of  $\approx 4\sigma$  in experiments as BaBar, LHCb, and Belle \cite{PhysRevLett.120.171802,PhysRevD.92.072014,PhysRevL

This is a problem because the standard model predicts for such process a characteristic known as the lepton Universality. Which states that in a process sufficiently energetic with a leptonic the 33.3% of the times the process will end with a  $\tau$  in the final state, 33.3% of the the times we will find a  $\mu$  in the final state and the other times will be a  $e$ . In that sense many models related to Beyond Standard Model physics as  $W', Z'$  model predict stronger interactions to the third generation of leptons and quarks compared to the other generations. \\

Many Studies had search new physics signs in process as  $b \rightarrow \tau, \nu$ , but in this sense the crossing symmetry can be used to study the phenomena by considering the three particles in the final state and the distribution of variables as the Transverse mass between the  $b, \nu$  and  $\tau, \nu$ , and the total mass between the 3 particles.

**Primary authors:** ATEHORTUA GARCES, Tomas (Universidad de Antioquia (CO)); Dr RUIZ ÁLVAREZ, José David (Universidad de Antioquia)

**Presenter:** ATEHORTUA GARCES, Tomas (Universidad de Antioquia (CO))

**Session Classification:** Common Session



Contribution ID: 49

Type: **Short Talk (5')**

## SpaceMath version 1.0: A Mathematica package for beyond the standard model parameter space searches.

*Friday, 4 December 2020 15:40 (5 minutes)*

We present a Mathematica package, called SpaceMath, for Beyond the Standard Model (BSM) parameter space searches which agree with the most up-to-date experimental measurements. The physical observables implemented in SpaceMath are classified in five categories, namely, LHC Higgs boson data (LHC-HBD), Flavor-Violating Processes (FVP), Oblique Parameters (OP), Unitarity and perturbativity (UP) and Meson Physics (MP). Nevertheless, SpaceMath version 1.0 (SpaceMath v1.0) works only with LHC-HBD and with extended scalar sector models. Future versions will implement the observables previously mentioned.

SpaceMath v1.0 is able to find allowed regions for free parameters of extension models by using LHC-HBD within a friendly interface and an intuitive environment in which users enter the couplings, set parameters and execute `Mathematica` in the traditional way. We present examples, step by step, in order to start new users in a fast and efficient way. To validate SpaceMath v1.0, we reproduce results reported in the literature.

**Primary authors:** ARROYO UREÑA, Marco Antonio; Dr GAITÁN-LOZANO, Ricardo (UNAM); Dr VALENCIA-PÉREZ, Tomás Antonio (BUAP)

**Presenter:** ARROYO UREÑA, Marco Antonio

**Session Classification:** Common Session

Contribution ID: 50

Type: **Regular Talk (15'+5')**

## Front-end readout electronics for the PDS-SP of the DUNE Experiment

*Tuesday, 1 December 2020 15:35 (20 minutes)*

The Deep Underground Neutrino Experiment (DUNE) is one of the next generation of neutrino experiments in construction in the USA under an international scheme. Colombian institutions are involved in the development of the DUNE experiment as an example of how Latin American countries can help design one mega-science experiment. This talk summarizes the Colombian activities in the design of the front-end readout electronics (DAPHNE) for the photon detection system of DUNE and presents the perspectives for the future regional working around the project.

**Primary author:** MORENO LOPEZ, Deywis (Universidad Antonio Narino (CO))

**Presenter:** MORENO LOPEZ, Deywis (Universidad Antonio Narino (CO))

**Session Classification:** Neutrinos

**Track Classification:** Neutrinos

Contribution ID: 51

Type: **Regular Talk (15'+5')**

## Heavy Sterile Neutrino Decay at Short-Baseline Experiments

*Thursday, 3 December 2020 09:20 (20 minutes)*

We studied a neutrino decay scenario as a potential solution to conciliate the tension between appearance and disappearance data at the short-baseline experiments. Particularly, we considered a heavy neutrino mass-eigenstate that decays into a usual light neutrino plus a massless scalar. Under this neutrino decay hypothesis, we fitted LSND and MiniBooNE electron neutrino appearance data assuming Dirac or Majorana neutrinos. We obtained reasonable results for both cases. Including muon neutrino disappearance searches, and also current bounds on the new decay coupling constant, we noticed that the heavy neutrino decay scenario is compatible as long as  $1 \text{ MeV} \leq m_4 \leq 10 \text{ keV}$ . Finally, we showed that the future SBN program at Fermilab has the potential to definitively test the considered decay hypothesis.

**Primary authors:** VITTI STENICO, Gabriela (State University of Campinas (UNICAMP)); DE GOUVEA, Andre (Northwestern University); PRAKASH, Suprabh (Sun Yat-Sen University, Guangzhou); PERES, Orlando L. G. (UNICAMP)

**Presenter:** VITTI STENICO, Gabriela (State University of Campinas (UNICAMP))

**Session Classification:** Neutrinos

**Track Classification:** Neutrinos

Contribution ID: 52

Type: **Regular Talk (15'+5')**

## Description and constraints of the back-end trigger of the Photon Detection System in DUNE

*Tuesday, 1 December 2020 15:55 (20 minutes)*

The photon detection system has been of great importance at DUNE (Deep Underground Neutrino Experiment). It can improve the sensitivity of the instrument by adding photon traces of the events, it's possible to increase data on CC interactions, and get information of the less common NC scattering on Ar. Supernova events are the high-level constraint DUNE must achieve in terms of bandwidth because it is impossible to know the rate of neutrinos in such events. For this reason, we present a description of the triggering scheme on each of the channels of the photon detection system, the region of zero-delay digitization, and the bandwidth of the readout system, for further trigger schemes descriptions of higher level.

**Primary author:** AM, Manuel (EIA)

**Presenter:** AM, Manuel (EIA)

**Session Classification:** Neutrinos

**Track Classification:** Neutrinos

Contribution ID: 53

Type: **Regular Talk (15'+5')**

## Systematically building the generalized Proca and SU(2) Proca theories of gravity and beyond

*Thursday, 3 December 2020 15:20 (20 minutes)*

To date, different alternative theories of gravity involving Proca and SU(2) Proca fields have been proposed. Unfortunately, the procedure to obtain the relevant terms has not been systematic enough or exhaustive, thus resulting in some missing terms or ambiguity in the process carried out. In this talk, we will show a systematic procedure to build the generalized Proca and SU(2) Proca field theories and beyond in four dimensions. In our approach, we employ all the possible Lorentz-invariant Lagrangian pieces made out of the Proca and SU(2) Proca fields and their first-order derivatives, and find the relevant combinations that propagate only three degrees of freedom and have healthy dynamics for the longitudinal mode. The key step in our procedure is to retain the flat space-time divergences of the currents in the theory during the covariantization process. In the curved space-time theory, some of the retained terms are no longer current divergences so that they induce the new terms that identify the beyond generalized Proca and SU(2) Proca field theories.

**Primary author:** GALLEGO CADAVID, ALEXANDER

**Presenter:** GALLEGO CADAVID, ALEXANDER

**Session Classification:** Cosmology / Astroparticles

**Track Classification:** Cosmology / Astroparticles

Contribution ID: 54

Type: **Regular Talk (15'+5')**

## Vector Boson Fusion Topology and Simplified Models for Dark Matter searches at colliders

*Wednesday, 2 December 2020 16:35 (20 minutes)*

A possibility to attempt at observing Dark Matter is to produce it at high energy colliders such as the Large Hadron Collider (LHC). LHC proton-proton collisions might result in the production of WIMPs in association with one or more QCD jets, photons as well as other detectable SM debris. Since WIMPs are electrically neutral and cosmologically stable massive particles, they manifest at colliders as missing transverse energy  $E_T$ . We explore prospects for dark matter at the LHC via the VBF topology. Particularly, we focus on scalar and vector-like mediators to look for better constraints on the masses and couplings regarding those previously found via the mono-jet strategy.

**Primary author:** OCAMPO HENAO, Daniel (Universidad de Antioquia (CO))

**Co-authors:** RUIZ-ÁLVAREZ, José D (Universidad de Antioquia); DUQUE-ESCOBAR, Santiago (Universidad de Antioquia)

**Presenter:** OCAMPO HENAO, Daniel (Universidad de Antioquia (CO))

**Session Classification:** Dark matter

**Track Classification:** Dark matter

Contribution ID: 55

Type: **Regular Talk (15'+5')**

## Five texture zeros for Dirac neutrino mass matrices

*Thursday, 3 December 2020 10:15 (20 minutes)*

In this work, we propose new five texture zeros for the mass matrices in the lepton sector in order to predict neutrino masses. In our approach, we extend beyond the standard model (SM) by assuming Dirac masses for the neutrinos, a feature which allows us to make a theoretical prediction for the lightest neutrino mass in the normal ordering. The textures that were analyzed have enough free parameters to adjust the VPMNS mixing matrix including the CP-violating phase, the neutrino mass squared differences, and the three charged lepton masses. we used two different procedures, the first method was based on a least-squares analysis and the second approach was algebraic.

**Primary authors:** BEANVIDES, Richard; Prof. GIRALDO, Yitsbhey; Prof. MUÑOZ, Luis; Prof. PONCE, William; Prof. ROJAS, Eduardo

**Presenter:** BEANVIDES, Richard

**Session Classification:** Neutrinos

**Track Classification:** Neutrinos

Contribution ID: 57

Type: **Regular Talk (15'+5')**

## WIMP Dark Matter in a Type-II Scotogenic model

*Wednesday, 2 December 2020 16:55 (20 minutes)*

Dark Matter and neutrinos are one of the most puzzling components of the Universe. Neutrino masses can be explained by radiative processes where Dark Matter particles are involved. Such models are known as Scotogenic DM models. The Dark Matter candidate in these models are stable thanks to the same symmetry that protect the radiative process. We present a realization of the scotogenic model using as inspiration model the Type-II seesaw. We show the model has a good DM candidate at the TeV scale and its phenomenology can be tested by CTA and Darwin.

**Primary author:** LINEROS, Roberto (Universidad Católica del Norte)

**Presenter:** LINEROS, Roberto (Universidad Católica del Norte)

**Session Classification:** Dark matter

**Track Classification:** Dark matter



Contribution ID: 58

Type: **Short Talk (5')**

## Large extra dimension at JUNO

*Friday, 4 December 2020 15:30 (5 minutes)*

We use the JUNO experiment which is a reactor neutrino experiment to constrain the parameters of the Large Extra Dimension (LED) model. The parameters of this model are the radius of extra dimension  $R_{LED}$  and the lightest neutrino mass  $m_0$ . We select the JUNO experiment because its aim is to determine the hierarchy of the neutrino masses and for this it will simultaneously measure the oscillations due to the parameters  $\Delta m_{21}^2$  and  $|\Delta m_{32}^2|$  and also the mixing angles  $\theta_{12}$  and  $\theta_{13}$  using a resolution on the visible energy of the positrons of 1% at 1 MeV. The LED model used in this work considers that neutrinos are Dirac neutrinos, the space-time structure of our universe is  $4 + 1$  that is: four flat spatial dimensions where the extra spatial dimension is compactified in a circle of radius  $R_{LED}$  and the 1 indicates a time dimension. We expect the results obtained in the parameters  $R_{LED}$  and  $m_0$  from this analysis will be slightly more restrictive than previous work already done due to the energy resolution of the JUNO experiment.

**Primary author:** BASTO-GONZALEZ, Victor (Universidad del Valle/ Universidad de Pamplona)

**Presenter:** BASTO-GONZALEZ, Victor (Universidad del Valle/ Universidad de Pamplona)

**Session Classification:** Common Session

Contribution ID: 59

Type: **Regular Talk (15'+5')**

## Cosmological implications of electroweak vacuum instability: constraints on the Higgs curvature coupling from inflation

*Thursday, 3 December 2020 16:55 (20 minutes)*

The current experimentally measured parameters of the Standard Model (SM) suggest that our universe lies in a metastable electroweak vacuum, where the Higgs field could decay to a lower vacuum state with catastrophic consequences. Our measurements dictate that such an event has not happened yet, despite the many different mechanisms that could have triggered it during our past light-cone. Via this observation, we can establish a promising link between cosmology and particle physics and thus constrain important parameters of our theories. The focus of our work has been to explore this possibility by calculating the probability of the false vacuum to decay during the period of inflation and using it to constrain the last unknown renormalisable SM parameter  $\xi$ , which couples the Higgs field with space-time curvature. In our latest study, we derived lower bounds for the Higgs-curvature coupling from vacuum stability in three inflationary models: quadratic and quartic chaotic inflation, and Starobinsky-like power-law inflation. In contrast to most previous studies we took the time-dependence of the Hubble rate into account both in the geometry of our past light-cone and in the Higgs effective potential, which is approximated with three-loop renormalisation group improvement supplemented with one-loop curvature corrections. We find that in all three models, the lower bound is  $\xi \gtrsim 0.051 \dots 0.066$  depending on the top quark mass. We also demonstrated that vacuum decay is most likely to happen a few e-foldings before the end of inflation.

**Primary author:** MANTZIRIS, Andreas (Imperial College London)

**Co-authors:** MARKKANEN, Tommi; Prof. RAJANTIE, Arttu (Imperial College (GB))

**Presenter:** MANTZIRIS, Andreas (Imperial College London)

**Session Classification:** Cosmology / Astroparticles

**Track Classification:** Cosmology / Astroparticles

Contribution ID: 60

Type: **Regular Talk (15'+5')**

## Estimation of the contribution of Muon Forward Scattering to the signal registered by the MuTe detector

*Thursday, 3 December 2020 17:35 (20 minutes)*

Muography is a technique that allows us to estimate the distribution of matter in the inner structure of geophysical formations like volcanoes. This technique is based on the measurement of the absorption of atmospheric muons that go through the structure, so that with a hodoscope instrument it is possible to build a map of density of matter, similar to a medical X-ray image. The MuTe (Muon-Telescope) collaboration, from *Universidad Industrial de Santander*, has designed and deployed a hybrid detector, with the same name. This detector consists of both a hodoscope and a water Cherenkov detector (WCD), and it has been conceived with two scientific aims in mind: applying Muography to study volcanic structures in Colombia, and working as a muon counting detector. In Muography, one source of noise is the Muon Forward Scattering (MFS) phenomenon, i.e. muons coming from the ground after a deflection caused by processes such as Coulomb interactions, that can be erroneously interpreted as muons that have crossed the structure of interest, when they actually have not. In this work, we quantify this effect using Monte Carlo simulations via the Geant4 toolkit, where we modeled the ground as standard rock. Thus, as a main result, we have estimated the MFS effect on the MuTe signal using an array of computer simulations: CORSIKA, to generate a realistic background of muons and other particles created in extensive air showers from cosmic rays; MAGNETOCOSMICS, to account for the effect that Earth's magnetic field has on the trajectories of primary cosmic rays; and a model of MuTe in Geant4, to estimate the superposition of the signal from MFS over the signal produced by the background of particles.

**Primary authors:** Mr DE LEÓN, Ricardo (Universidad Industrial de Santander); SUÁREZ, Mauricio (Universidad Industrial de Santander); NÚÑEZ, Luis A. (University of Los Andes)

**Presenter:** Mr DE LEÓN, Ricardo (Universidad Industrial de Santander)

**Session Classification:** Cosmology / Astroparticles

**Track Classification:** Cosmology / Astroparticles

Contribution ID: 61

Type: **Short Talk (5')**

## $Z'$ to $t\bar{t}$ in the single lepton-final state with Atlas Open Data

*Friday, 4 December 2020 15:10 (5 minutes)*

Different models of physics Beyond the Standard Model (BSM) predict different kinds of  $Z'$  bosons, motivating the search for these particles at the Large Hadron Collider (LHC). In this work, a  $Z' \rightarrow t\bar{t}$  analysis in the semileptonic top quark pair channel is presented, using the ATLAS Open Data Dataset at a centre-of-mass energy of 13TeV and corresponding to an integrated luminosity of  $10\text{fb}^{-1}$ . An event selection criteria is implemented, where acceptable agreement between the data and Monte Carlo simulated predictions is observed.

**Primary authors:** GOMEZ CRUZ, Nicolas; ORDONEZ, Sebastian (National University of Colombia)

**Co-author:** SANDOVAL USME, Carlos (Universidad Nacional de Colombia)

**Presenters:** GOMEZ CRUZ, Nicolas; ORDONEZ, Sebastian (National University of Colombia)

**Session Classification:** Common Session

Contribution ID: 62

Type: **Short Talk (5')**

## **Simulation of a Water Cherenkov Detector Response to the Background of Cosmic Rays at Pamplona Norte de Santander altitude (2300 m a.s.l.) for the LAGO Collaboration**

*Friday, 4 December 2020 15:15 (5 minutes)***KEYWORDS:**

Cosmic Rays, Water Cherenkov Detector, Monte Carlo Simulations.

**ABSTRACT:**

The studies of cosmic rays (CR) is mainly performed by using detectors installed at the ground; this detectors registered the secondary radiation produced by the interaction of CR with Earth atmosphere. Since 2010 Colombia is an active member of the large scale observatory LAGO (Latin American Giant Observatory). This observatory operated an array of Water Cherenkov Detectors (WCDs) spanned over Latin America in a wide range of latitudes. The installation of a new WCD requires the perform of a series of simulation, from the calculation of the flux of primaries that arrive to the Pamplona atmosphere until the WCD response to the secondary radiation. Here we show the signal expected at Pamplona, Norte de Santander city. For this we have use the AGO's toolkit ARTI, a framework of computational techniques and codes (CORSIKA, GEANT4, C++, Python) to estimated the WCD signals produced by cosmic radiation at ground. As main results, the spectrum of secondaries here calculated fits with the expected and the WCD response allows to us to separate the signal deposited by muons from the electrons/positron ones. In particular, our calculations shows the muon signal far away from the noise signal, so we expected that a WCD installed at Pamplona can works as a muon counting. As a final results, we have checked the capability of the WCD simulated to measured the muon half life time, this implies that a WCD deployed at the University of Pamplona can be used as Lab equipment to teach particle physics.

**Primary author:** PARADA VILLAMIZAR, Heidar Marcel**Co-author:** Dr SUAREZ DURAN, Mauricio**Presenter:** PARADA VILLAMIZAR, Heidar Marcel**Session Classification:** Common Session

Contribution ID: 63

Type: **Short Talk (5')**

# Characterization of the signal produced by muons into one Water Cherenkov Detectors of the LAGO Observatory

*Friday, 4 December 2020 15:20 (5 minutes)*

**The Latin American Giant Observatory (LAGO) is an observatory array of Water Cherenkov**

Detectors (WCDs) expanded throughout Latin America, from Mexico to Antarctica, with scientific objectives to study the cosmic rays, space weather phenomena, and ground-level atmospheric radiation. The quality of the data recorded by LAGO is guaranteed based on a protocol that includes the installation process and periodic monitoring of the instrument's calibration. This calibration is based on the signal produced by muon-like particles that completely and vertically pass through the WCD, it is called VEM (Vertical Equivalent Muon); so, VEM identification is important to the trustworthiness and correct interpretation of the LAGO data. Currently, the LAGO's algorithm to process and analysis the data allows us to identify the VEM signal, but this algorithm fails occasionally for some data, mainly because the water into the WCD becomes more absorbing to Cherenkov photons, so this data is rejected. Here we show a statistical analysis of the signals produced by muons into one of the WCD of LAGO with the aim of recovery the rejected data. In this study, we have used a full month of accepted data to characterize the properties of the muon signal, i.e. peak and integral of the signal, the decay time of the pulse, among others, later we built a pulse model for muon signal and we have tested it throughout Monte Carlo simulations, using the simulation framework of LAGO based on Geant4. Finally, we have validated this model with another set of accepted data.

Keywords: Background Cosmic radiation, Muon signal, Water Cherenkov Detector

**Primary author:** Mr PEREZ CUEVAS, YORLAN (Universidad de Pamplona)

**Co-author:** Dr SUAREZ DURAN, MAURICIO (Universidad de Pamplona )

**Presenter:** Mr PEREZ CUEVAS, YORLAN (Universidad de Pamplona)

**Session Classification:** Common Session

Contribution ID: 64

Type: **Regular Talk** (15'+5')

## Estimation of the effective dose from cosmic radiation received during a commercial flight passing through the South Atlantic anomaly

*Thursday, 3 December 2020 17:55 (20 minutes)*

Exposure to ionizing radiation during commercial flights has repercussions on human health, in particular for flights pass through the South Atlantic Geomagnetic Anomaly (SAA). This radiation is originated from the interaction of cosmic rays (CR) with Earth's atmosphere. Likewise, the flux of CR with energies around GeV is affected by the Geomagnetic field, this establishes a dependence between ionizing radiation and geographical position. Currently, tools such as CARI-7A estimates the effective dose for commercial flights. In general, this kind of code uses semi-empirical models to estimate the radiation at the flight altitude and ignore the contribution of particles like muons on the calculation of the dose; this because of the low probability of interaction with matter and the low dose of muons compare with neutron particles. Nonetheless, a better estimation of the radiation at flight altitude and the contribution of muons for the effective dose can improve this calculation due to the bigger flux of CR into the SAA. Here we show an improvement of the estimation of the effective dose during a commercial flight that crosses the AAS by calculating the secondary radiation produced by CR between 1 GeV and  $10^6$  GeV, via CORSIKA and GATE codes. In this work, we have included a model (metallic cylinder) of the plane fuselage. In addition, we have evaluated the influence of the geomagnetic field on secondary radiation through a method that uses its local and instantaneous configuration. In this way, this work is a contribution to the understanding of how high-energy RC ( $<10^6$  GeV), the aircraft fuselage, and the geomagnetic field can improve the estimation of the effective dose estimated for commercial flights that cross the SAA.

Keywords: ionizing radiation, effective dose, commercial flights, Geomagnetic field.

**Primary author:** Prof. BECERRA VILLAMIZAR, Daniel Camilo (Universidad de Pamplona)

**Co-author:** SUÁREZ, Mauricio (Universidad Industrial de Santander)

**Presenter:** Prof. BECERRA VILLAMIZAR, Daniel Camilo (Universidad de Pamplona)

**Session Classification:** Cosmology / Astroparticles

**Track Classification:** Cosmology / Astroparticles

Contribution ID: 65

Type: **Short Talk (5')**

# Estimation of energy levels for cosmic rays affected during a Forbush decrease from data from the LAGO Observatory

*Friday, 4 December 2020 15:25 (5 minutes)*

## Abstract

The space weather phenomena have an important impacts in science and current technological structure. This work is focused on the interaction between solar transient events and cosmic rays, specifically on Forbush Decrease (FD) events. A FD consists in a decrease/affectation on the counts of radiation a ground level during an Interplanetary Coronal Mass Ejection (ICME) crossing the Geomagnetic field. Here we show a method to estimated the cosmic rays affected during the FD of March of 2012 in terms of the Geomagnetic rigidity cut-off. For this, we have used the data registered at San Carlos de Bariloche, Argentina, by one of the detectors of the LAGO observatory (an extended cosmic radiation observatory over Latin American, from México to the Antarctic). This detector is a water Cherenkov detector that works as muon counting, i.e. we could followed the rate of muon at ground and because these muon coming from hadronic interactions, we built a Rigidity cut-off function to estimate which primaries, according with their magnetic rigidity,were deflected out of their track to the atmosphere. To verify this approach, we validity this function through the LAGO's toolkit ARTI, a framework of computational techniques and codes (CORSIKA, GEANT4, C++, Python), i.e. we calculated the secondaries produced by this rate of cosmic rays and then, the signal produced by those secondaries into the Geant4 detector model.

**Primary author:** PARADA JAIME, Karoll Michely

**Co-author:** Dr SUAREZ DURAN, Mauricio

**Presenter:** PARADA JAIME, Karoll Michely

**Session Classification:** Common Session



Contribution ID: 66

Type: **Regular Talk (15'+5')**

## Machine Learning Applications to Reactor Antineutrino Detection with PROSPECT

*Tuesday, 1 December 2020 16:15 (20 minutes)*

The Precision Reactor Oscillation and Spectrum Experiment (PROSPECT) is an above-ground antineutrino experiment at short baselines located at the High Flux Isotope Reactor (HFIR) at Oak Ridge National Laboratory (ORNL). This experiment's physics goals include searching for the existence of sterile neutrinos and precisely measuring the antineutrino energy spectrum. At PROSPECT, antineutrinos are detected via the inverse beta decay (IBD) interaction. This process provides a near-unique space-time correlated signal pair consisting of a positron energy deposition and a delayed neutron capture in the liquid scintillator (LS). The correlation between prompt and delayed pulses/signals is an excellent handle for background suppression. The ORNL group is currently exploring several applications of machine learning techniques for the reconstruction and analysis of antineutrino events. In this presentation, I will focus on one of these efforts which attempts to improve the current positron identification capability by identifying a subset of positrons through orthopositronium formation.

**Primary authors:** VENEGAS VARGAS, Diego (The University of Tennessee Knoxville/Oak Ridge National Laboratory); HEFFRON, Blaine (The University of Tennessee Knoxville/Oak Ridge National Laboratory)

**Presenter:** VENEGAS VARGAS, Diego (The University of Tennessee Knoxville/Oak Ridge National Laboratory)

**Session Classification:** Neutrinos

**Track Classification:** Neutrinos

Contribution ID: 67

Type: **Regular Talk (15'+5')**

## El Sistema Global de Asimilación de Datos GDAS, para la estimación del flujo de fondo de astropartículas a nivel del suelo en la colaboración LAGO

*Thursday, 3 December 2020 17:15 (20 minutes)*

Presentamos una metodología que permite obtener perfiles atmosféricos mensuales para el estudio del flujo de astropartículas a nivel del suelo, usando el Sistema Global de Asimilación de Datos GDAS. Estos perfiles pueden ser creados para cualquier ubicación geográfica, y su implementación permitió cuantificar los efectos de las variaciones de densidad, en el flujo de partículas.

Creamos 12 perfiles mensuales para la ciudad de Bucaramanga y observamos diferencias entre estos y el perfil predeterminado por la herramienta de simulación CORSIKA. Las diferencias fueron significativas para el flujo a nivel del suelo generado por núcleos primarios, con energías entre 10 GeV y  $10^6$  GeV, considerando el campo magnético terrestre. Encontramos que el flujo simulado con la atmósfera predeterminada, difiere hasta por un  $\approx 10\%$  del flujo mes a mes. Además, también se observa una correlación entre las variaciones de flujo y los cambios de temperatura a lo largo del año, permitiendo determinar que la implementación de estos perfiles es necesaria para obtener estimaciones más realistas incluso en regiones tropicales.

**Primary author:** GRISALES, Jennifer

**Co-authors:** NÚÑEZ, Luis A. (University of Los Andes); SARMIENTO-CANO, Christian (Universidad Industrial de Santander)

**Presenter:** GRISALES, Jennifer

**Session Classification:** Cosmology / Astroparticles

**Track Classification:** Cosmology / Astroparticles

Contribution ID: 68

Type: **Regular Talk (15'+5')**

## Neutrino oscillation probabilities due to possible non-standard interactions

*Thursday, 3 December 2020 09:40 (20 minutes)*

In this work, we studied the modifications of the neutrino oscillation probabilities due to possible non-standard interactions (NSI) with matter (taking into account a constant matter density profile), focused on the channels  $\nu_\mu \rightarrow \nu_\mu$  and  $\nu_\mu \rightarrow \nu_e$ . First, the neutrino oscillation probability in matter without NSI was studied and then we proceeded with NSI. The most relevant NSI parameters that affect the oscillation phenomenon were also studied. To carry out this study, a code was written using the Mathematica™ software, in which the Hamiltonian was defined in terms of non-standard interaction, the evolution operator and the initial and final states of the neutrinos, then the parameters  $\Delta m_{231}^2$  and  $\theta_{23}$  were changed at  $1\sigma$  and  $3\sigma$  of Confidence Level (C.L) for the particular configuration of the NOvA and DUNE experiments. As a result, the  $\nu_e$  appearance probability is affected by the parameters  $\epsilon_{ee}$ ,  $\epsilon_{\mu\mu}$ ,  $\epsilon_{\tau\tau}$  and  $\mu\tau$ , and their range of allowed values are found, taking into account the current experimental oscillation measurements and uncertainties. Unlike the channel of  $\nu_e$  appearance, in muon neutrino disappearance channel there is only one NSI parameter that affects this probability and this is  $\mu\tau$  that takes the range  $-0.07 \leq \mu\tau \leq 0.07$  at  $1\sigma$  and  $-0.176 \leq \mu\tau \leq 0.132$  at  $3\sigma$  for the NOvA experiment and for the DUNE experiment a range was found within the values  $-0.032 \leq \mu\tau \leq 0.032$  at  $1\sigma$  and  $-0.083 \leq \mu\tau \leq 0.090$  at  $3\sigma$  of C.L.

**Primary authors:** BARROS SARMIENTO, Andrea Carolina (Universidad del Atlántico); ACERO, Mario A (Universidad del Atlántico)

**Presenter:** ACERO, Mario A (Universidad del Atlántico)

**Session Classification:** Neutrinos

**Track Classification:** Neutrinos

Contribution ID: 69

Type: **Short Talk (5')**

## LFV signals with Dirac neutrinos and axions

In this work we present a phenomenological analysis of a WIMP/axion dark matter model with radiatively generated neutrino masses. In this model the PQ mechanism is responsible for Dirac neutrino masses through the realization of a five-dimension effective operator at one-loop level, as well as for the stability of all mediators in the loop by means of a remaining  $Z_2$  symmetry originating from the spontaneous breaking of the PQ symmetry. As a result, mixed WIMP-axion dark matter scenarios are obtained. The model is implemented in high energy physics packages such as SARAH, SPheno and micrOMEGAS for its study, and constraints from LFV processes and direct detection searches are used to bound the valid parameter space of the model.

**Primary author:** RODRIGUEZ CIFUENTES, OSCAR ANDRES (Universidad de Antioquia)

**Presenter:** RODRIGUEZ CIFUENTES, OSCAR ANDRES (Universidad de Antioquia)

**Session Classification:** Common Session

Contribution ID: 70

Type: **Regular Talk (15'+5')**

## 3-3-1 model with right neutrinos, implemented in SARAH and SPheno

*Monday, 30 November 2020 15:00 (20 minutes)*

In this work, for the first time the implementation of the 3-3-1 model with right-handed neutrinos without exotic electric charges in SARAH package of MATHEMATICA is presented, and it shows how it correctly reproduces the analytical results of the model. As proof of this, we present the mass matrices for the quark sectors, where a Higgs sector with three scalar triplets has been used. Then, using the SPheno program, a numerical analysis of the analytical outputs obtained with SARAH is performed; this with the objective of determining if the model with three triplets generates the mass values of all quarks adequately when comparing them with those accepted in the literature. This implementation is presented in a didactic and accessible way to students who are interested in this field of physics.

**Primary authors:** Dr TAPIA, Alex (University of Medellin); BEANVIDES, Richard

**Presenter:** Dr TAPIA, Alex (University of Medellin)

**Session Classification:** Formal theory

**Track Classification:** Higgs / Standard model

Contribution ID: 71

Type: **not specified**

## **50 years of the GIM mechanism**

*Monday, 30 November 2020 09:00 (1 hour)*

**Presenter:** MAIANI, Luciano (Sapienza Universita e INFN, Roma I (IT))

**Session Classification:** Opening session

Contribution ID: 72

Type: **not specified**

## **Black Holes in the Milky Way**

*Monday, 30 November 2020 10:00 (1 hour)*

**Presenter:** LU, Jessica (University of California-Berkeley)

**Session Classification:** Opening session

Contribution ID: 73

Type: **not specified**

## Gauge Theories on Quantum Computers

*Monday, 30 November 2020 11:15 (1 hour)*

**Presenter:** BERENSTEIN, David (University of California at Santa Barbara)

**Session Classification:** Opening session



Contribution ID: 74

Type: **not specified**

## **Higgs Physics at the LHC: status and prospects**

*Monday, 30 November 2020 12:15 (1 hour)*

**Presenter:** CAMACHO TORO, Reina Coromoto (Centre National de la Recherche Scientifique (FR))

**Session Classification:** Opening session

Contribution ID: 75

Type: **not specified**

## Challenges for the 3-3-1 models

*Friday, 4 December 2020 09:20 (30 minutes)*

**Presenter:** PLEITEZ, VICENTE (UNIVERSIDADE ESTADUAL PAULISTA-UNESP)

**Session Classification:** Heavy flavour

Contribution ID: 76

Type: **not specified**

## **EFTs: Bounding the systematic uncertainties of the Inverse Amplitude Method**

*Tuesday, 1 December 2020 08:00 (30 minutes)*

**Presenter:** LLANES-ESTRADA, Felipe J.

**Session Classification:** Formal theory

Contribution ID: 77

Type: **not specified**

## Scattering Amplitudes: QCD and gravity applications

*Tuesday, 1 December 2020 08:40 (30 minutes)*

**Presenter:** Dr TORRES BOBADILLA, William J. (IFIC CSIC-UV)

**Session Classification:** Formal theory

Contribution ID: 78

Type: **not specified**

## Neutrinoless double beta decay searches

*Tuesday, 1 December 2020 14:00 (30 minutes)*

**Presenter:** CARDANI, Laura (INFN - National Institute for Nuclear Physics)

**Session Classification:** Neutrinos

Contribution ID: 79

Type: **not specified**

## Neutrino oscillations

*Tuesday, 1 December 2020 14:40 (30 minutes)*

**Presenter:** CATANO-MUR, Erika (William & Mary)

**Session Classification:** Neutrinos

Contribution ID: **80**

Type: **not specified**

## **SM measurements**

*Wednesday, 2 December 2020 09:00 (30 minutes)*

**Presenter:** MANJARRES, Joany (Technische Universitaet Dresden (DE))

**Session Classification:** LHC results

Contribution ID: **81**

Type: **not specified**

## **Searches for Higgs boson pairs with the ATLAS detector**

*Wednesday, 2 December 2020 11:00 (30 minutes)*

**Presenter:** PALACINO, Gabriel (Indiana University (US))

**Session Classification:** LHC results



Contribution ID: **82**

Type: **not specified**

## **SUSY searches**

*Wednesday, 2 December 2020 09:40 (30 minutes)*

**Presenter:** MARTINEZ RUIZ DEL ARBOL, Pablo (Universidad de Cantabria and CSIC (ES))

**Session Classification:** LHC results

Contribution ID: **83**

Type: **not specified**

## **Exotica searches**

*Wednesday, 2 December 2020 10:20 (30 minutes)*

**Presenter:** DE LA CRUZ, Begona (Centro de Investigaciones Energéticas Medioambientales y Tecnol.)

**Session Classification:** LHC results

Contribution ID: 84

Type: **not specified**

## Unraveling the origin of Black Holes from effective spin measurements

*Wednesday, 2 December 2020 14:00 (30 minutes)*

The remarkable discovery of gravitational waves from binary black hole mergers has given us a new way to study our universe. The origin of the black hole binaries remains unclear, I investigate whether information on the effective spin of binary black hole mergers from the LIGO-Virgo gravitational wave detections can be used to discriminate primordial versus astrophysical black holes. I will also present the posterior probability density for a possible mixture of astrophysical and primordial BHs as emerging from current data, and calculate the number of future mergers needed to discriminate different spin and alignment models at a given level of significance.

**Presenter:** FERNANDEZ, Nicolas (University of Illinois at Urbana–Champaign)

**Session Classification:** Dark matter

Contribution ID: 85

Type: **not specified**

## Searching for dark matter emission with highly energetic cosmic messengers

*Wednesday, 2 December 2020 14:40 (30 minutes)*

Indirect dark matter detection is a powerful search method that allows to explore a very wide range of the dark matter mass spectrum. I will present our recent results in which we exploit multi-messenger astrophysical observations to explore the allowed parameter space for various archetypal dark matter candidates with masses in the range 10 GeV to  $10^{16}$  GeV. Throughout my talk, I will explain our efforts to account for the systematic astrophysical uncertainties impacting our results.

**Primary author:** MACÍAS, Óscar (Virginia Tech)

**Presenter:** MACÍAS, Óscar (Virginia Tech)

**Session Classification:** Dark matter

Contribution ID: 86

Type: **Regular Talk (15'+5')**

## The KM3NeT Neutrino Observatory: opportunities for Latin American collaborators

*Tuesday, 1 December 2020 16:35 (20 minutes)*

The KM3NeT Neutrino Observatory is being built in two abyssal sites in the Mediterranean Sea: ARCA (IT), optimized for astrophysical neutrinos, and ORCA (FR), optimized for atmospheric neutrinos studies. The KM3NeT observatory will host the largest water Cherenkov detectors ever built in the northern hemisphere, providing real-time and high-bandwidth connection to shore, also available for Earth and Life Sciences. ARCA will be ready to survey the sky with unprecedented

angular resolution and sensitivity, providing an optimal view of the Southern sky including the Galactic Center. ORCA will provide a  $3\sigma$  sensitivity on NMH after three years of exposure and competitive precision in  $\Delta m^2_{32}$  and  $\sin^2 \theta_{23}$  using atmospheric neutrinos. The P2O proposal to shoot a long baseline neutrino beam from the Protvino accelerator to ORCA could also open new perspectives that will be briefly addressed in this contribution, particularly, regarding  $\delta CP$  estimation. Dark Matter, Non-Standard Interactions, Multi-Messenger Astronomy, and Neutrino Tomography complete the KM3NeT scope. Regarding Earth and Sea Sciences programs, the abyss of the Mediterranean as site of prime interest, is used for research and long-term monitoring of geohazards, marine life and ocean dynamics.

Exciting times for Latin America scientific community are coming in Space, Earth and Sea Sciences, with the arrival of KM3NeT to the region. The construction, operation and scaling taking place nowadays are the main target of this contribution. Nonetheless, main technical aspects and calibrations of the observatory are presented, together expectations of its ambitious science programs, and some first physics results. Opportunities for Colombian collaborators are particularly motivated, looking for triggering a fore coming discussion next in time.

**Primary author:** Prof. YEPES RAMIREZ, HAROLD (Yachay Tech / KM3NeT)

**Presenter:** Prof. YEPES RAMIREZ, HAROLD (Yachay Tech / KM3NeT)

**Session Classification:** Neutrinos

**Track Classification:** Neutrinos

Contribution ID: 87

Type: **not specified**

## The diffuse supernova neutrino background, a new window to the Universe

*Thursday, 3 December 2020 08:00 (30 minutes)*

Since the first generation of stars, core-collapse supernovae have produced a steady flux of neutrinos, which could be detectable in the next-generation of experiments. Measuring this continuous flux, known as the diffuse supernova neutrino background (DSNB), could put novel bounds on possible beyond-the-Standard Model scenarios, such as lifetimes and oscillations expected if neutrinos are pseudo-Dirac particles. Moreover, the DSNB could teach us about cosmology and astrophysics since these neutrinos have been propagating in an expanding Universe. We will explore these possibilities in this talk.

**Primary author:** PEREZ, Yuber

**Presenter:** PEREZ, Yuber

**Session Classification:** Neutrinos

Contribution ID: **88**

Type: **not specified**

## **BSM physics program at DUNE**

*Thursday, 3 December 2020 08:40 (30 minutes)*

**Presenter:** Prof. SOUSA, Alexandre (University of Cincinnati (US))

**Session Classification:** Neutrinos

Contribution ID: **89**

Type: **not specified**

## **Cosmic Rays Experiments in Latin America**

*Thursday, 3 December 2020 14:00 (30 minutes)*

**Presenter:** ASOREY, Hernan (Comision Nacional de Energia Atomica)

**Session Classification:** Cosmology / Astroparticles



Contribution ID: **90**

Type: **not specified**

## **A guidance for building dark energy models**

*Thursday, 3 December 2020 14:40 (30 minutes)*

**Presenter:** GOMEZ, gabriel

**Session Classification:** Cosmology / Astroparticles

Contribution ID: **91**

Type: **not specified**

## **Review on CPV and mixing in the charm system**

*Friday, 4 December 2020 08:00 (30 minutes)*

**Presenter:** CORREA DOS REIS, Alberto (CBPF - Centro Brasileiro de Pesquisas Físicas (BR))

**Session Classification:** Heavy flavour

Contribution ID: 92

Type: **not specified**

## Searching for flavored new physics at the LHC

*Friday, 4 December 2020 08:40 (30 minutes)*

**Presenter:** MARTIN CAMALICH, Jorge (Inst. Astrophys. of Canary Islands (ES))

**Session Classification:** Heavy flavour

Contribution ID: 93

Type: **not specified**

## Open Discussion

*Wednesday, 2 December 2020 18:15 (20 minutes)*

**Session Classification:** Dark matter

Contribution ID: **94**

Type: **not specified**

## Discussion

*Monday, 30 November 2020 16:20 (20 minutes)*

**Session Classification:** Formal theory

Contribution ID: 95

Type: **not specified**

## Discussion

*Tuesday, 1 December 2020 10:55 (20 minutes)*

**Session Classification:** Formal theory

Contribution ID: **96**

Type: **not specified**

## **Discussion**

*Tuesday, 1 December 2020 17:15 (20 minutes)*

**Session Classification:** Neutrinos

Contribution ID: 97

Type: **not specified**

## Discussion

*Wednesday, 2 December 2020 11:30 (20 minutes)*

**Session Classification:** LHC results



Contribution ID: **98**

Type: **not specified**

## Discussion

*Thursday, 3 December 2020 11:35 (20 minutes)*

**Session Classification:** Neutrinos

Contribution ID: **99**

Type: **not specified**

## Discussion

*Thursday, 3 December 2020 18:15 (20 minutes)*

**Session Classification:** Cosmology / Astroparticles

Contribution ID: 100

Type: **Regular Talk (15'+5')**

## QMexico - Latin America's role in the second quantum revolution

*Friday, 4 December 2020 14:00 (20 minutes)*

QMexico is an organization that brings quantum computing researchers and enthusiasts from Colombia and Mexico together. Its main goal is to popularize quantum technologies and software. Also, through education and skill development opportunities, QMexico is training the next generation of quantum scientists. For this talk, we will focus on recent efforts to promote this emerging technology in Latin America through a hands-on workshop on the basics of quantum computing and the organization of QTalks, where Hispanic researchers talk about their work on the subject. We will also highlight the importance of bridging these innovative tools with high energy physics to solve some of the computational challenges facing the HEP community in the upcoming years.

**Primary authors:** DELGADO, Andrea (Texas A & M University (US)); DELGADO, Andrea

**Presenters:** DELGADO, Andrea (Texas A & M University (US)); DELGADO, Andrea

**Session Classification:** Common Session

**Track Classification:** Formal theory

Contribution ID: **101**

Type: **Regular Talk (15'+5')**

## **CONHEP status**

*Friday, 4 December 2020 17:00 (20 minutes)*

**Presenter:** SANDOVAL USME, Carlos (Universidad Nacional de Colombia)

**Session Classification:** CONHEP

Contribution ID: **102**

Type: **Regular Talk (15'+5')**

## **LASF4RI status**

*Friday, 4 December 2020 17:20 (20 minutes)*

**Presenter:** Prof. RESTREPO, Diego (Universidad de Antioquia)

**Session Classification:** CONHEP

Contribution ID: **103**

Type: **Regular Talk (15'+5')**

## **Colaboración Arte y Ciencia**

**Session Classification:** CONHEP

Contribution ID: **104**

Type: **Regular Talk (15'+5')**

## **LA-CoNGA Physics**

*Friday, 4 December 2020 14:20 (20 minutes)*

**Presenter:** NÚÑEZ, Luis A. (University of Los Andes)

**Session Classification:** Common Session

Contribution ID: **105**

Type: **not specified**

## **Las fronteras inexistentes entre el arte y la ciencia**

*Friday, 4 December 2020 14:40 (10 minutes)*

**Presenter:** BRILL, Daniela (Suratómica)

**Session Classification:** Common Session



Contribution ID: **106**

Type: **not specified**

## **Proyectos Suratómica**

*Friday, 4 December 2020 14:50 (10 minutes)*

**Presenters:** RIVERA, Carlos; ARIAS, Margareth

**Session Classification:** Common Session