

# LHCP2021

The Ninth Annual Conference on Large Hadron Collider Physics


Online

7-12 June 2021

~~Paris (France), Sorbonne Université~~

~~(IN2P3/CNRS, IRFU/CEA)~~


## Heavy-flavour collectivity in heavy-ion collisions



ALICE



ATLAS  
EXPERIMENT




*Fabrizio Grosa* on behalf of the ALICE, ATLAS, and CMS Collaborations

CERN


- a colour-deconfined medium, called *quark-gluon plasma* (QGP) is created in ultrarelativistic heavy-ion collisions
- The initial geometrical anisotropy is transferred to a momentum anisotropy by the pressure gradients

$$E \frac{d^3N}{dp_T} = \frac{1}{2\pi} \frac{d^2N}{p_T dp_T dy} \left\{ 1 + \sum_{n=1}^{\infty} v_n \cos[n(\varphi - \Psi_n)] \right\}$$

$$v_n = \langle \cos[n(\varphi - \Psi_n)] \rangle \quad \text{Flow harmonic coefficients}$$



- Azimuthal anisotropy in HF hadron production
  - ➔ Participation of charm and beauty quarks in the collective motions and their possible thermalisation
  - ➔ Study path-length dependence of in-medium parton energy loss
  - ➔ Sensitivity to initial-state event-by-event fluctuations
  - ➔ Probe strong initial electromagnetic fields in the QGP



- Significant  $v_2$  for open and hidden charm mesons  $\rightarrow$  charm participation to collective motions
- $p_T < 3-4$  GeV/c: mass hierarchy
  - $\rightarrow v_2(J/\psi) < v_2(D, D^0) < v_2(\pi)$
- $3-4 < p_T < 6-8$  GeV/c: coalescence
  - $\rightarrow v_2(J/\psi) < v_2(D, D^0) = v_2(\pi)$
- $p_T > 8$  GeV/c: path-length dependence of in-medium energy loss
  - $\rightarrow v_2(J/\psi) = v_2(D, D^0) = v_2(\pi)$

ALI-PUB-483361


ALICE prompt D: PLB 813 (2021) 136054

ALICE  $J/\psi$  forward  $y$ : JHEP 10 (2020) 141


ALICE  $\pi^\pm$ : JHEP 09 (2018) 006

CMS prompt  $D^0$ : PRL 120 (2018) 202301

ALICE  $J/\psi$  mid  $y$ : JHEP 10 (2020) 141



- D-meson  $v_2\{4\}$  smaller than  $v_2\{2\}$ 
  - ➔ Expected in case of event-by-event fluctuations in the flow signal
- Similar effect for charm and light hadrons
- Described by theoretical models based on charm-quark transport in a hydrodynamically expanding QGP



- Significant  $v_3$  for open and hidden charm mesons → charm sensitive to initial-state event-by-event fluctuations
- $p_T < 3-4$  GeV/c: mass hierarchy  
→  $v_3(J/\psi) < v_3(D, D^0) < v_3(\pi)$
- $3-4 < p_T < 6-8$  GeV/c: coalescence  
→  $v_3(J/\psi) < v_3(D, D^0) = v_3(\pi)$
- $p_T > 8$  GeV/c: path-length dependence of in-medium energy loss  
→  $v_3(J/\psi) = v_3(D, D^0) = v_3(\pi)$

ALI-PUB-483366


ALICE prompt D: PLB 813 (2021) 136054

ALICE  $J/\psi$  forward  $y$ : JHEP 10 (2020) 141

CMS prompt  $D^0$ : PRL 120 (2018) 202301

ALICE  $\pi^\pm$ : JHEP 09 (2018) 006

ALICE prompt D: PLB 813 (2021) 136054



- Transport of charm quarks in an hydrodynamically expanding medium via Boltzmann or Langevin equations
  - ▶ diffusion coefficient  $D_s$  related to thermalisation time of charm quark


$1.5 < 2\pi D_s T_c < 7$  for models that describe data with  $\chi^2/\text{ndf} < 2$

$\tau_{\text{charm}} = (m_{\text{charm}}/T) \cdot D_s = 3 - 14 \text{ fm}/c \approx \tau_{\text{QGP}}$

TAMU: PRL 124 (2020) 042301	Catania: PLB 805 (2020) 135460
MC@sHQ+EPOS2: PRC 89 (2014) 014905	POWLANG: EPJC (2019) 79:494
LGR: arXiv:1912.08965	LBT: PRC 94 (2016) 014909
LIDO: PRC 98 (2018) 064901	BAMPS: JPG 42 (2015) 11, 115106
PHSD: PRC 93 (2016) 034906	DAB-MOD: PRC 102 (2020) 024906

- D-meson  $v_2$  studied as a function of average event flow with event-shape engineering technique

- ▶ Positive correlation between heavy and light hadron  $v_2$
- ▶ Sensitivity for charm to event-by-event fluctuations




20% smallest  $q_2^{\text{TPC}}$

$$\langle v_2 \rangle_{\text{small-}q_2} < \langle v_2 \rangle_{\text{unb}}$$

20% largest  $q_2^{\text{TPC}}$

$$\langle v_2 \rangle_{\text{large-}q_2} > \langle v_2 \rangle_{\text{unb}}$$

ALICE prompt D: PLB 813 (2021) 136054



ALI-PUB-490285

POWLANG: EPJC (2019) 79:494

Catania: PLB 805 (2020) 135460

DAB-MOD: PRC 102 (2020) 024906

LIDO: PRC 98 (2018) 064901


- Initial strong electric (magnetic) fields are expected to produce a charge dependence in the elliptic (directed) flow

→ Charm quarks ideal probes due to their early production


U. Gürsoy et al, PRC 98 (2018) 055201

S. K. Das et al, PLB 768 (2017) 260-264

A. Dubla et al, MPA Vol. 35, No. 39 (2020) 2050324





CMS prompt  $D^0$ : PLB 816 (2021) 136253



ALI-PUB-342594

ALICE  $D^0$ : PRL 125 (2020) 022301

- No indication of charge dependent  $v_2$  as a function of rapidity
- $2.7\sigma$  significant indication of  $y$ -dependent charge difference for  $D^0$  mesons





- Elliptic BW: PRC 101 (2020) 064905
- MC@sHQ+EPOS2: PRC 89 (2014) 014905
- LIDO: PRC 98 (2018) 064901
- PHSD: PRC 93 (2016) 034906
- DAB-MOD: PRC 102 (2020) 024906
- DREENA-B: PLB 791 (2019) 236-241





- Positive  $v_2$  of leptons from beauty-hadron decays both at low and high  $p_T$ 
  - ➔ Indication of participation in the collective motions of the system
  - ➔ Lower than charm (larger mass)
  - ➔ Expected contribution from hadronisation via coalescence with light quarks from the medium



ATLAS  $c, b \rightarrow \mu$ : PLB 807 (2020) 135595

DAB-MOD: PRC 102 (2020) 024906




- No indication of positive  $v_3$  of muons from beauty-hadron decays neither in central nor semicentral collisions
  - ➔ Compatible with very low  $v_3$  predicted by DAB-MOD transport model
  - ➔ Lower  $p_T$  range important to draw a firm conclusion

-  CMS  $\Upsilon(1S)$ : [arXiv:2006.07707](https://arxiv.org/abs/2006.07707)
-  CMS  $\Upsilon(2S)$ : [arXiv:2006.07707](https://arxiv.org/abs/2006.07707)
-  ALICE  $\Upsilon(1S)$ : PRL 123 (2019) 192301
-  ALICE  $J/\psi$ : JHEP 10 (2020) 141



- $v_2$  of  $\Upsilon(1S)$  and  $\Upsilon(2S)$  compatible with zero within large uncertainties in all the  $p_T$  and centrality intervals measured
  - ➔ No contribution from coalescence with light quarks from the QGP
  - ➔ Does beauty quark flow?
- For more information about quarkonia see [presentation by Minjung Kim on Mon June 7th at 17:15](#)



ALI-DER-486560

ALICE  $\pi^\pm$ : JHEP 09 (2018) 006

ALICE prompt D: PLB 813 (2021) 136054


ALICE  $J/\psi$ : JHEP 10 (2020) 141

ALICE  $b \rightarrow e$ : PRL 126 (2021) 162001

ALICE  $\Upsilon(1S)$ : PRL 123 (2019) 192301

TAMU: PRL 124 (2020) 042301, NPA 943 (2015) 147–158, PRC (2017) 96, 054901

- Precise measurements of different hadron species crucial for setting constraints to models



- Simultaneous description of different observables challenging

ATLAS  $\mu$ : ATLAS-CONF-2021-020

DAB-MOD: PRC 102 (2020) 024906

DREENA-B: PLB 791 (2019) 236-241

- Charm-quark flow: positive  $v_2$  and  $v_3$  of open and hidden charm mesons, correlated with light hadrons
  - ➔ Indicate significant degree of thermalisation of charm quarks in the QGP
  - ➔ Indicate sensitivity to initial-state event-by-event fluctuations
  - ➔ Contribution from charm-quark hadronisation via coalescence
- Search for strong early electromagnetic fields in heavy-ions
  - ➔ Electric field: no significant D-meson charge-dependent  $v_2$
  - ➔ Magnetic field: hint of D-meson charge-dependent  $v_1$
- Beauty-quark flow: positive  $v_2$  for open-beauty hadrons, no significant  $v_2$  for bottomonia
  - ➔ Does beauty quark flow?
  - ➔ Only contribution from beauty-quark hadronisation via coalescence?

D. Longieras IJClab 2021

# LHCP2021

The Ninth Annual Conference on Large Hadron Collider Physics


Online

7-12 June 2021


~~Paris (France), Sorbonne Université~~

~~(IN2P3/CNRS, IRFU/CEA)~~


ADDITIONAL SLIDES



- QCD calculations on lattice predict a phase transition from the ordinary nuclear matter to a colour-deconfined medium, called *quark-gluon plasma* (QGP)
  - ➔ created in ultrarelativistic heavy-ion collisions
  - ➔ very high energy density  $\varepsilon > 15 \text{ GeV/fm}^3$
  - ➔ after a pre-equilibrium phase expands hydrodynamically
- Heavy flavours: produced in shorter time scales than QGP formation time
  - ➔  $\tau_{\text{HF}} \approx \hbar/m \approx 0.05\text{-}0.1 \text{ fm}/c$  depending on  $p_{\text{T}}$
  - ➔  $\tau_{\text{QGP form}} (\text{LHC}) \approx 0.3 \text{ fm}/c$
 } HF experience the full system evolution



- HF propagate in the QGP with a **Brownian motion**
  - interact with medium constituents
  - loose energy via **elastic collisions** and **radiative processes**
  - heavy-quark **thermalisation** in the QGP?



→  $\tau_{\text{HF}} \approx \hbar/m \approx 0.05\text{-}0.1 \text{ fm}/c$  depending on  $p_T$


→  $\tau_{\text{QGP form}} (\text{LHC}) \approx 0.3 \text{ fm}/c$

PRC 89 (2014) 034906

→  $\tau_{\text{QGP lifetime}} \approx 10 \text{ fm}/c$

PLB 696 (2011) 328-315

- Competing mechanisms for the HF hadronisation in the QGP




## 1. Fragmentation $D_{q \rightarrow h}(z_q, Q^2)$

- energy-loss of partons while traversing the QGP modifies fraction of the parton momentum  $z_q$  taken by the hadron
  - equal for all hadron species

## 2. Coalescence

- partons close in phase space can recombine into hadrons
- quarks with different mass coalesce if have similar velocities





- 📖 PRL 90 (2003) 202302
- 📖 PRL 90 (2003) 202303
- 📖 PRC 67 (2003) 064902
- 📖 PLB 595 (2004) 202-208

- Statistical hadronisation model

- Hadrons emitted from the interaction region in statistical equilibrium at the QGP phase boundary

- 📖 PLB (2008) 659:149-155


S. K. Das et al, PLB 768 (2017) 260-264



- Originated by two competing effects:

- ➔ Faraday effect: electric field induced by decreasing  $\vec{B}$

- ➔ Hall effect: Lorentz force induced by moving charges  $\vec{F} = q \vec{v} \times \vec{B}$




ALICE prompt D: PLB 813 (2021) 136054

ALICE prompt  $D_s^+$ : preliminary


- No significant difference between  $D_s^+$  and non-strange D mesons within current uncertainties

# Centrality and rapidity dependence of charm meson elliptic flow





ALICE J/ψ: JHEP 10 (2020) 141  
ALICE  $\pi^\pm$ : JHEP 09 (2018) 006

CMS prompt  $D^0$ : PLB 816 (2021) 136253  
CMS charged particles: PLB 776 (2017) 195



PLB 813 (2021) 136054




- The event-shape engineering (ESE) technique relies on the classification of events at a certain centrality according to the magnitude of the second-harmonic reduced flow vector:

$$q_2 = |\vec{Q}_2| / \sqrt{M}$$


$$\vec{Q}_2 = \sum_{j=1}^M e^{i2\varphi_j}$$

ALI-PUB-348698




20% smallest  $q_2^{TPC}$

$$\langle v_2 \rangle_{\text{small-}q_2} < \langle v_2 \rangle_{\text{unb}}$$




20% largest  $q_2^{TPC}$

$$\langle v_2 \rangle_{\text{large-}q_2} > \langle v_2 \rangle_{\text{unb}}$$



ALICE prompt D: PLB 813 (2021) 136054

- Models based on charm-quark transport in an hydrodynamically expanding medium describe  $q_2$  dependence of D-meson  $v_2$  flow
- Predicted variation of D-meson  $v_2$  in ESE-selected samples with respect to unbiased sample similar for different transport parameters (e.g. POWLANG HTL vs. IQCD)




POWLANG: EPJC (2019) 79:494

Catania: PLB 805 (2020) 135460


DAB-MOD: PRC 102 (2020) 024906

LIDO: PRC 98 (2018) 064901




ALICE prompt D: PLB 813 (2021) 136054

- No significant modification of the  $p_T$ -differential production yields of D-mesons in different ESE-selected samples of events



POWLANG: EPJC (2019) 79:494  
LIDO: PRC 98 (2018) 064901




- Charm-quark production time comparable with maximum of magnetic field created in heavy-ion collisions


S. K. Das et al, PLB 768 (2017) 260-264  
 A. Beraudo et al, arXiv:2102.08064

➔ Charge-dependent directed flow ideal observable to study early magnetic fields

- Indication of  $y$ -dependent charge difference
- ➔ Different magnitude for charged particles and D<sup>0</sup> mesons
- ➔ 2.6 $\sigma$  significance for charged particles
- ➔ 2.7 $\sigma$  significance for D<sup>0</sup> mesons



ALI-PUB-325477



CMS  $\Upsilon(1S)$ : arXiv:2006.07707

ALICE  $\Upsilon(1S)$ : PRL 123 (2019) 192301


ALICE  $J/\psi$ : JHEP 10 (2020) 141

TAMU: PRC (2017) 96, 054901

BBJS: PRC 100 (2019) 051901

Yao: PRD 100 (2019) 014008

Hong, Lee: PLB 801 (2020) 135147



- $v_n$  of D mesons can be obtained as a weighted average of the one of charm and light quarks:

$$v_n^D(p_T^D) = v_n^q(p_T^q) + v_n^c(p_T^c)$$

- ➔ Quark  $p_T$  and  $v_n$  obtained as half of those of charged pions and J/ψ mesons
- ➔ Best agreement in case of  $p_T^D/p_T^q = 0.4$



ALI-PUB-483386

ALICE  $\pi^\pm$ : JHEP 09 (2018) 006

ALICE J/ψ forward  $y$ : JHEP 10 (2020) 141

CMS prompt  $D^0$ : PRL 120 (2018) 202301

ALICE J/ψ mid  $y$ : JHEP 10 (2020) 141




● No significant  $v_2$  of muons ( $D^0$ ) from beauty-hadron decays in pp (pPb) collisions

➔ Is it zero or just very small?

➔  $v_2$  mainly driven by beauty-quark coalescence in heavy-ion collisions?

➔ For more information about small systems see presentation by Georgios Krintiras on Wed June 9th at 16:00



ATLAS  $c \rightarrow \mu$ : PLB 807 (2020) 135595

CMS prompt D<sup>0</sup>: PRL 120 (2018) 202301