

The new Δa_μ measurement

Muon anomalous magnetic moment

$$\Delta a_\mu^{\text{exp}} = a_\mu^{\text{exp}} - a_\mu^{\text{SM}}$$

Experimental measurement

Standard Model prediction

Brookhaven National Laboratory (2006): $\Delta a_\mu^{\text{exp}} = (279 \pm 76) \times 10^{-11}$ (3.7σ)

Fermilab 2021 + BNL 2006 new combined result: $\Delta a_\mu^{\text{exp}} = (251 \pm 59) \times 10^{-11}$ (4.2σ)

NEW-PHYSICS EXPLANATIONS

Supersymmetric models
Left-right symmetric models
Scotogenic models
331 models

$L_\mu - L_\tau$ models
Seesaw models
Zee-Babu model

2HDMs

Impact of the new measurement in **type-X** and **type-II** models
2HDMs + vector-like leptons (VLLs) which do not mix with the muon


Δa_μ in 2HDMs + VLLs

2HDM extension with VLLs (NO mixing with the muon)

$$-\mathcal{L}_{\text{VLL}} = m_L \bar{\chi}_L \chi_R + m_E \bar{E}_L E_R + \lambda_L \bar{\chi}_R \Phi_1 E_L + \lambda_R \bar{\chi}_L \Phi_1 E_R + \text{H.c.}$$

	$\chi_{L,R}$	$E_{L,R}$
$SU(2)$	2	1
$U(1)$	-1/2	-1

- One neutral lepton N
- Two charged leptons L_a^-
- Left and right-handed charged-current interactions
- No new one-loop contributions




New Barr-Zee diagrams involving the new charged and neutral leptons

Charged VLLs masses


Charged VLLs mixing angles

Benchmark cases $\left\{ \begin{array}{l} B_1: M_2 = 2.5 M_1 = 250 \text{ GeV} \\ B_2: M_2 = 1.5 M_1 = 150 \text{ GeV} \end{array} \right.$ $\sin \theta_L = 0.5$ $\sin \theta_R = 0.4$

RESULTS




Above the solid (dashed) horizontal line, $\lambda_{\text{max}} \equiv \max\{|\lambda_L|, |\lambda_R|\} > 4\pi$ for B_1 (B_2).



- The ranges of m_A can be substantially enlarged with respect to the 2HDMs ones
- $\tan \beta$ can be shifted to lower values (drastically lowered in the μSpec model)
- This effect is more pronounced for B_1

The larger VLL coupling with the Higgs doublet Φ_1 is required to be either close or above the perturbative limit $\lambda_{\text{max}} = 4\pi$

New Δa_μ measurement in 2HDMs



Considering the one-loop and two-loop Barr-Zee contributions in a 2HDM without tree-level flavour changing neutral currents


Sizeable positive contribution from the two-loop pseudoscalar processes can account for the observed Δa_μ

leads to

A light pseudoscalar with coupling to the muon proportional to $\tan \beta$

Type-II, type-X and μSpec models

RESULTS



The impact of the new result is marginal

Type-X model can accommodate the discrepancy but requires large values of $\tan \beta$ and very light pseudoscalars

Type II requires light pseudoscalars in conflict with perturbativity, unitarity and electroweak precision constraints

The μSpec model requires extreme fine-tuning and values of $\tan \beta$ of $O(1000)$ in order to accommodate the discrepancy

Can the addition of VLLs relax these conclusions?

CONCLUSIONS

Fitting the Δa_μ discrepancy

Type-X and type-II models	Light pseudoscalars and large values of $\tan \beta$
2HDMs + VLLs	Parameter space is widened <ul style="list-style-type: none"> Much larger values of the pseudoscalar mass Lower values of $\tan \beta$ VLL Yukawa couplings \gtrsim perturbativity (can be alleviated considering more families of VLLs or VLL-muon mixing)

¹ Departamento de Física and CFTP, Instituto Superior Técnico, Universidade de Lisboa, Lisboa, Portugal

² Centro de Física Teórica e Computacional, Faculdade de Ciências, Universidade de Lisboa, Campo Grande, Edifício C8 1749-016 Lisboa, Portugal

³ Instituto Superior de Engenharia de Lisboa, Instituto Politécnico de Lisboa 1959-007 Lisboa, Portugal

⁴ High Energy Theory Group, William & Mary, Williamsburg, VA 23187, USA