

F3C-uR models further analysis

Disha Bhatia, Indian Association for the Cultivation of
Science

Inclusive k-factor:

1) **xx process**

generate p p > dm dm /excluded [QCD] → NLO

generate p p > dm dm /excluded → LO

Cuts: $p_{tj} > 100 \text{ GeV}$, $|\eta_{tj}| < 5$

Pdf: 260000

2) **xy process**

generate p p > dm yy /excluded [QCD] → NLO

generate p p > dm yy/excluded → LO Cuts: $p_{tj} > 30 \text{ GeV}$, $|\eta_{tj}| < 5$

3) yy QCD:

generate p p > yy yy /excluded [QCD] → NLO

generate p p > yy yy /excluded → LO

Cuts: $p_{tj} > 30 \text{ GeV}$


4) yy-t-channel

generate p p > yy yy /excluded DMT=2 QCD=0 QED=0 [QCD] → NLO

generate p p > yy yy/excluded DMT=2 QCD=0 QED=0 → LO


The mediator is not decayed for evaluating the inclusive k-factor

inclusive k-factor (xx)




K-factor
sensitive to
both mediator
and dark
matter
particle mass

inclusive k-factor (xy)




Sensitivity largely on the mediator

inclusive k-factor (yy-tchannel)



inclusive k-factor (yy-qcd)



Exclusive k-factor: Leading order commands

1) xx:

```
generate p p > dm dm /excluded ; add process p p > dm dm j /excluded $yy
```

2) xy:

```
generate p p > yy dm /excluded ; add process p p > yy dm j /excluded $yy
```


3) yy:

```
generate p p > yy yy /excluded [for t-channel: we use in addition DMT=2  
QCD=0 QED=0]
```



Merging plots (xx -mode):

$m_X = 200 \text{ GeV}$

$m_Y = 500 \text{ GeV}$



log10d01 []



mx= 550 GeV
My = 1000 GeV

Differential k-factor for $m_x=200$ GeV, $m_y=550$ GeV

