

Searching for long-lived particles at the LHC and beyond: Eighth workshop of the LHC LLP Community



Contribution ID: 15

Type: **not specified**

Extending the reach of FASER, MATHUSLA, and SHiP towards smaller lifetimes using secondary particle production (12'+3')

Thursday 19 November 2020 15:20 (15 minutes)

Many existing or proposed intensity-frontier search experiments look for decay signatures of light long-lived particles (LLPs), highly displaced from the interaction point. This approach is, however, limited to new particles with decay lengths similar to or larger than the baseline of those experiments. We will discuss how this basic constraint can be overcome in BSM models that go beyond the simplest scenarios due to additional secondary production of LLPs right in front of the detector. This way a new lifetime regime can be probed. We illustrate the prospects of such searches in the future experiments FASER, MATHUSLA, and SHiP. We also analyze additional advantages from employing dedicated neutrino detectors placed in front of the main decay volume.

The talk is based on <https://arxiv.org/pdf/1911.11346.pdf>

Primary authors: JODLOWSKI, Krzysztof (National Centre for Nuclear Research); ROSZKOWSKI, Leszek (National Centre for Nuclear Research (PL)); TROJANOWSKI, Sebastian; KLING, Felix (SLAC)

Presenter: JODLOWSKI, Krzysztof (National Centre for Nuclear Research)

Session Classification: New ideas