



Facultad de Ciencias
UNIVERSIDAD DE LA SERENA



Agencia
Nacional de
Investigación
y Desarrollo

Ministerio de Ciencia,
Tecnología, Conocimiento
e Innovación

Gobierno de Chile

Fondecyt

Fondo Nacional de Desarrollo
Científico y Tecnológico

LLP and multi-lepton signatures from neutrino mass models

Juan Carlos Helo


Searching for long-lived particles at the LHC and beyond: Eighth workshop of the LHC LLP Community.

November 2020

Long-lived charged particles and multilepton signatures from neutrino mass models Arbelaez, Cottin, Helo, Hirsch. *Phys.Rev.D* 101 (2020) 9, 095033

Type-I Seesaw

Minkowski; Gellman, Ramon, Slansky; Yanagida; Glashow; Mohapatra,
Senjanovic



Some displaced N studies in:

G. Cottin, J.C. Helo and M. Hirsch, Phys. Rev. D98 (2018).



E. Izaguirre and B. Shuve, Phys. Rev. D91 (2015)

S. Dube, D. Gadkari, and A. M. Thalapillil, Phys. Rev. D96 (2017)

J. C. Helo, M. Hirsch, and S. Kovalenko, Phys. Rev. D89 (2014)


... and others

$c\tau$ drops below mm for m_N above 100 GeV - No displaced vertex



1-loop models:

SCOTOGENIC



E. Ma, *Phys. Rev. D*73 (2006) 077301,


Additional symmetry to the SM $SU(2)_L \times U_Y \times Z_2$

$$(\nu_i, l_i) \sim (2, -1/2; +), \quad l_i^c \sim (1, 1; +), \quad N_i \sim (1, 0; -),$$

$$(\phi^+, \phi^0) \sim (2, 1/2; +), \quad (\eta^+, \eta^0) \sim (2, 1/2; -).$$

2 Dark Matter Candidates !

OUR MODEL



Arbelaez, Cottin, Helo Hirsch *Phys.Rev.D* 101 (2020) 9, 095033

No additional symmetries to the SM


$$S_{1,1,2} = S_1^{2+}$$

$$S_{1,3,3} = (S_3^{4+}, S_3^{3+}, S_3^{2+})$$

$$F_{1,2,5/2} = (F^{3+}, F^{2+})$$

No Dark Matter Candidates

OUR MODEL



All these particles will decay to SM particles with at least two same-sign leptons!






FIG. 2: Feynman diagrams for the decays $H^{4+} \rightarrow l_\gamma^+ l_\delta^+ l_\alpha^+ l_\beta^+$ and $H^{4+} \rightarrow W^+ W^+ l_\alpha^+ l_\beta^+$.

We define LNV multi-lepton events as final states with at least Four charged leptons!.



In our model multi-lepton events have larger rates than di-lepton events!



6l 2w LNV!!





Pair production of MCP is dominated by photon-photon fusion diagrams for large scalar masses.



Long-Lived Multi-Charged Particles:

Smallness of the observed neutrino masses, together with the high multiplicity of the final states, lead to Long-Lived Multi-Charged Particles!!!



$$m_\nu \simeq 0.05 \left(\frac{\lambda_5}{10^{-6}} \right) \left(\frac{h_F}{10^{-2}} \right) \left(\frac{h_{\bar{F}}}{10^{-2}} \right) \left(\frac{1 \text{ TeV}}{\Lambda} \right) \text{ eV}$$

Current limits: Non of existing searches covers exactly the multi-lepton signals!

$c\tau >$ detector size. MCP based on only 36 /fb

ATLAS collaboration, M. Aaboud et al., Phys. Rev. D99 (2019) 052003,

Based on the anomalously large ionization of MCP that are long-lived enough to reach the muon spectrometer $M_{H4+} > 980$ GeV. Limits only applies for $c\tau$ larger than 10m!

$c\tau = 1$ mm -1 m detector size. Displaced vertices - Disappearing tracks.

CMS collaboration, Phys. Rev. Lett. 114 (2015) 061801, [1409.4789].

CMS collaboration, Search for displaced leptons in the e-mu channel, CMS-PAS-EXO-16-022.

ATLAS collaboration, G. Aad et al. Phys. Rev. D92 (2015) 072004, [1504.05162].

CMS collaboration, A. M. Sirunyan et al., JHEP 08 (2018) 016, [1804.07321].

ATLAS collaboration, M. Aaboud et al., JHEP 06 (2018) 022, [1712.02118].

J. A. Evans and J. Shelton, JHEP 04 (2016) 056, [1601.01326].

R. Mahbubani, P. Schwaller and J. Zurita, JHEP 06 (2017) 119, [

ATLAS collaboration, M. Aaboud et al., Phys. Rev. D97 (2018) 052012, [1710.04901].

ATLAS collaboration, M. Aaboud et al., Phys. Rev. D99 (2019) 052003,



Displaced vertex and disappearing tracks searches can give limits to our model

$c\tau < 1$ mm. Prompt decays

CMS collaboration, A. M. Sirunyan et al., Search for physics beyond the standard model in multilepton final states in proton-proton collisions at $\sqrt{s} = 13$ TeV JHEP 03 (2020) 051

For the parameter region of our model where the scalars decay promptly, various “seesaw-searches” can give limits to our model.

Other models can be constructed that leads
To exotic multi-lepton signals!



$LNV!$
 $4e \ 2w \ 2j$

$$S_{33\frac{7}{3}} = (S^{4/3}, S^{2/3}, S^{10/3})$$

\swarrow
 all with color!!

Conclusions

We discussed 1-loop neutrino mass model which does not require additional symmetries to be the leading contribution to neutrino masses. We showed that in this approach multi-charged particles which are Long-lived naturally appears as a consequence of the smallness of neutrino masses and the high multiplicity of the final states. These particles have masses up to 10 TeV.

Different searches can be sensitive to our model: from searches of stable MCP, to displaced vertex signals, depending on the half-life of the new particles.

Multi-leptons signals can arise in our model. In particular we have discussed the possibility of having multi-lepton LNV final states at the LHC.

Other interesting models can be built that lead to multi-lepton signatures. We have showed another possibility that leads to Long-lived colored Multi-charge particles.

Thanks!!