Beyond the Standard Model



Lectures at the 5th Latin-American School of High-Energy Physics,

Recinto Quirama, Antioquia, Colombia, March 2009

John.Ellis@cern.ch

Plan of the Lectures

- The Standard Model and issues beyond it
- Origin of particle masses: Higgs boson(s)?
- Supersymmetry
- Searches for supersymmetry
- Neutrinos and Grand Unification
- Extra dimensions and string theory

Summary of the Standard Model

- Particles and SU(3) × SU(2) × U(1) quantum numbers:

L_L	$\begin{pmatrix} \nu_e \\ e^- \end{pmatrix}_L , \begin{pmatrix} \nu_\mu \\ \mu^- \end{pmatrix}_L , \begin{pmatrix} \nu_\tau \\ \tau^- \end{pmatrix}_L$	$(\mathbf{1},\mathbf{2},-1)$
E_R	$e_R^- , \mu_R^- , \tau_R^-$	$(\mathbf{1},\mathbf{1},-2)$
Q_L	$\begin{pmatrix} u \\ d \end{pmatrix}_L , \begin{pmatrix} c \\ s \end{pmatrix}_L , \begin{pmatrix} t \\ b \end{pmatrix}_L$	$(\mathbf{3},\mathbf{2},+1/3)$
U_R	u_R , c_R , t_R	$(\mathbf{3},\mathbf{1},+4/3)$
D_R	d_R , s_R , b_R	$(\mathbf{3},\mathbf{1},-2/3)$

- Lagrangian:

$$\begin{aligned} \mathcal{L} &= -\frac{1}{4} F^a_{\mu\nu} F^{a\,\mu\nu} \\ &+ i\bar{\psi}\, \not{D}\psi + h.c. \\ &+ \psi_i y_{ij} \psi_j \phi + h.c. \\ &+ |D_\mu \phi|^2 - V(\phi) \end{aligned}$$

gauge interactions

matter fermions

Yukawa interactions

Higgs potential

Gauge Interactions of the Standard Model

- Three separate gauge group factors:
  - SU(3) × SU(2) × U(1)
  - Strong × electroweak
- Three different gauge couplings:
  - g_3, g_2, g □
- Mixing between the SU(2) and U(1) factors:

$$\begin{pmatrix} Z^\mu \\ A^\mu \end{pmatrix} = \begin{pmatrix} \cos(\theta_W) & \sin(\theta_W) \\ -\sin(\theta_W) & \cos(\theta_W) \end{pmatrix} \begin{pmatrix} W_3^\mu \\ B^\mu \end{pmatrix} \qquad \sin^2(\theta_W) = \frac{g'^2}{g'^2 + g^2}$$

- Experimental value: $\sin^2\theta_W = 0.23120 \pm 0.00015$

Important clue for Grand Unification

Weak Interactions

- Interactions of lepton doublets: $L = \begin{pmatrix} \nu_e \\ e^- \end{pmatrix}_L$
- Charged-current interactions:

$$\mathcal{L}_{cc} = \frac{-g}{\sqrt{2}} \sum_{\alpha=e,\mu,\tau} \nu_{L_\alpha} \gamma_\mu l_{L_\alpha} W^\mu \; + \; h.c.$$

- Neutral-current interactions:

$$\mathcal{L}_{nc} = \frac{-g}{2\cos\theta_W} \sum_{\alpha=e,\mu,\tau} \nu_{L_\alpha} \gamma_\mu l_{L_\alpha} Z^\mu \; + \; h.c$$

- Mixing between quark flavours:

$$\begin{pmatrix} d' \\ s' \\ b' \end{pmatrix} = V_{CKM} \begin{pmatrix} d \\ s \\ b \end{pmatrix} = \begin{pmatrix} V_{ud} & V_{us} & V_{ub} \\ V_{cd} & V_{cs} & V_{cb} \\ V_{td} & V_{ts} & V_{tb} \end{pmatrix} \begin{pmatrix} d \\ s \\ b \end{pmatrix}$$

Status of the Standard Model

- Perfect agreement with all *confirmed* accelerator data
- Consistency with precision electroweak data (LEP et al) *only if there is a Higgs boson*
- Agreement seems to require *a relatively light Higgs boson* weighing < ~ 180 GeV
- Raises many unanswered questions:

mass? flavour? unification?

Precision Tests of the Standard Model

Lepton couplings



Pulls in global fit

	Measurement	Fit
$\Delta\alpha^{(5)}_{had}(m_Z)$	0.02758 ± 0.00035	0.02767
m_Z [GeV]	91.1875 ± 0.0021	91.1874
Γ_Z [GeV]	2.4952 ± 0.0023	2.4959
σ^0_{had} [nb]	41.540 ± 0.037	41.478
R_l	20.767 ± 0.025	20.742
$A^{0,l}_{fb}$	0.01714 ± 0.00095	0.01643
$A_l(P_\tau)$	0.1465 ± 0.0032	0.1480
R_b	0.21629 ± 0.00066	0.21579
R_c	0.1721 ± 0.0030	0.1723
$A^{0,b}_{fb}$	0.0992 ± 0.0016	0.1038
$A^{0,c}_{fb}$	0.0707 ± 0.0035	0.0742
A_b	0.923 ± 0.020	0.935
A_c	0.670 ± 0.027	0.668
A_l(SLD)	0.1513 ± 0.0021	0.1480
$\sin^2\theta^{lept}_{eff}(Q_{fb})$	0.2324 ± 0.0012	0.2314
m_W [GeV]	80.410 ± 0.032	80.377
Γ_W [GeV]	2.123 ± 0.067	2.092
m_t [GeV]	172.7 ± 2.9	173.3



Parameters of the Standard Model

- Gauge sector:
  - 3 gauge couplings: g_3, g_2, g □
  - 1 strong CP-violating phase
- Yukawa interactions:
  - 3 charge-lepton masses
  - 6 quark masses
  - 4 CKM angles and phase
- Higgs sector:
  - 2 parameters: μ, λ
- **Total: 19 parameters**

Unification?

Flavour?

Mass?

Open Questions beyond the Standard Model

- What is the origin of particle masses?

  due to a Higgs boson? + other physics?

  solution at energy < 1 TeV (1000 GeV)

  Susy
- Why so many types of matter particles?

  matter-antimatter difference?
- Unification of the fundamental forces?

  at very high energy ~ 10^{16} GeV?

  probe directly via neutrino physics, indirectly via masses, couplings

  Susy
- Quantum theory of gravity?

  (super)string theory: extra space-time dimensions?

  Susy

At what Energy is the New Physics?



Origin of mass

Dark matter

A lot accessible To the LHC, ILC

Some accessible only indirectly

Why do Things Weigh?

Newton:

Weight proportional to Mass

Einstein:

Energy related to Mass

Neither explained origin of Mass

Where do the masses come from?

Are masses due to Higgs boson? (the physicists' Holy Grail)



2008 Nobel Physics Prize: Nambu

Illustration of the Higgs Idea

The excitation crosses the room = the Higgs boson

Think of a Snowfield

Skier moves fast:
Like particle without mass
e.g., photon = particle of light

Snowshoer sinks into snow,
moves slower:
Like particle with mass
e.g., electron

The LHC will look for the snowflake: The Higgs Boson

Hiker sinks deep,
moves very slowly:
Particle with large mass

The Higgs Mechanism

- Postulated effective Higgs potential:

$$V[\phi] = -\mu^2 \phi^\dagger \phi + \lambda (\phi^\dagger \phi)^2$$

- Minimum energy at non-zero value:

$$\phi_0 = < 0|\phi|0 > = \frac{1}{\sqrt{2}} \begin{pmatrix} 0 \\ +v \end{pmatrix} \quad v = \sqrt{\frac{\mu^2}{\lambda}}$$

- Non-zero masses: $M_f = y_f \frac{v}{\sqrt{2}}$ $M_W = \frac{g\,v}{2}$
- Components of Higgs field: $\phi(x) = \frac{1}{\sqrt{2}}(v + \sigma(x))e^{i\pi(x)}$
- π massless, σ massive: $m_H^2 = 2\mu^2 = 2\lambda v$

Masses for Gauge Bosons

- Kinetic terms for SU(2) and U(1) gauge bosons:

$$\mathcal{L} = -\frac{1}{4}\, G^i_{\mu\nu} G^{i\mu\nu} - \frac{1}{4} F_{\mu\nu} F^{\mu\nu}$$

  where $G^i_{\mu\nu} \equiv \partial_\mu W^i_\nu - \partial_\nu W^i_\mu + i g \epsilon_{ijk} W^j_\mu W^k_\nu$ $F_{\mu\nu} \equiv \partial_\mu W^i_\nu - \partial_\nu W^i_\mu$

- Kinetic term for Higgs field:

$$\mathcal{L}_\phi = -|D_\mu \phi|^2 \quad D_\mu \equiv \partial_\mu - i\ g\ \sigma_i\ W^i_\mu - i\ g'\ Y\ B_\mu$$

- Expanding around vacuum: $\phi = < 0|\phi|0 > + \hat{\phi}$

$$\mathcal{L}_\phi \ni -\frac{g^2 v^2}{2}\ W^+_\mu\ W^{\mu -} - g'^2\ \frac{v^2}{2}\ B_\mu\ B^\mu + g\ g' v^2\ B_\mu\ W^{\mu 3} - g^2\ \frac{v^2}{2}\ W^3_\mu\ W^{\mu 3}$$

- Boson masses:

$$m_{W^\pm} = \frac{gv}{2} \qquad Z_\mu = \frac{g W^3_\mu - g' B_\mu}{\sqrt{g^2 + g'^2}} \ :\ m_Z = \frac{1}{2}\sqrt{g^2 + g'^2}\, v\ ; \quad A_\mu = \frac{g' W^3_\mu + g B_\mu}{\sqrt{g^2 + g'^2}} \ :\ m_A = 0$$

Constraints on Higgs Mass

- Electroweak observables sensitive via quantum loop corrections:

$$m_W^2 \sin^2\theta_W = m_Z^2 \cos^2\theta_W \sin^2\theta_W = \frac{\pi\alpha}{\sqrt{2}G_F}(1+\Delta r)$$

- Sensitivity to top, Higgs masses:

$$\frac{3G_F}{8\pi^2\sqrt{2}}m_t^2$$

$$\frac{\sqrt{2}G_F}{16\pi^2}m_W^2\left(\frac{11}{3}\ln\frac{M_H^2}{m_Z^2}+\ldots\right),\ M_H >> m_W$$

- Preferred Higgs mass: **$m_H \sim 91$ GeV**
- Compare with lower limit from direct searches:

$m_H > 114$ GeV

- No conflict!

Nambu and Higgs

Spontaneous breaking of symmetry





Higgs Decay Branching Ratios

- Couplings proportional to mass:
  - Decays into heavier particles favoured



- But: important couplings through loops:
  - gluon + gluon → Higgs → γγ

Summer 2008

Indications on the Higgs Mass

Sample observable:
W mass @ LEP & Tevatron



Combined information
on Higgs mass



The State of the Higgs: March 2009

- Direct search limit from LEP:

  $m_H > 114.4$ GeV
- Electroweak fit sensitive to m_t

  (Now $m_t = 173.1 \pm 1.3$ GeV)
- Best-fit value for Higgs mass:

  $m_H = 84^{+34}_{-26}$ GeV
- 95% confidence-level upper limit:

  $m_H < 154$ GeV, or 185 GeV including direct limit
- Tevatron exclusion:

  $m_H < 160$ GeV or > 170 GeV

Higgs Discovery @ Tevatron?



Tevatron excludes Higgs between 160 & 170 GeV

Combining the Higgs Information



$m_H = 116.4^{+\ 18.3}_{-1.3}$ GeV

Combining the Higgs Information



$m_H = 116.4^{+\,18.3}_{-1.3}$ GeV

Theoretical Constraints on Higgs Mass

- Large → large self-coupling → blow up at low energy scale Λ due to renormalization
- Small: renormalization due to t quark drives quartic coupling < 0
  → vacuum unstable
- Bounds on Higgs mass depend on Λ



How to Stabilize a Light Higgs Boson?

- Top quark destabilizes potential: introduce stop-like scalar:

$$\mathcal{L} \supset M^2|\phi|^2 + \frac{M_0}{v^2}|H|^2\,|\phi|^2$$

- Can delay collapse of potential:
- But new coupling must be fine-tuned to avoid blow-up:
- Stabilize with new fermions:
  - just like Higgsinos
- Very like **Supersymmetry!**








The Large Hadron Collider (LHC)
Proton- Proton Collider
7 TeV + 7 TeV
1,000,000,000 collisions/second
Primary targets:
•Origin of mass
•Nature of Dark Matter
•Primordial Plasma
•Matter vs Antimatter

Huge Statistics thanks to High Energy and Luminosity

Event rates in ATLAS or CMS at $L = 10^{33}$ cm^{-2} s^{-1}

Process	Events/s	Events per year	Total statistics collected at previous machines by 2007
$W \rightarrow e\nu$	15	10^8	10^4 LEP / 10^7 Tevatron
$Z \rightarrow ee$	1.5	10^7	10^7 LEP
$t\bar{t}$	1	10^7	10^4 Tevatron
$b\bar{b}$	10^6	10^{12} - 10^{13}	10^9 Belle/BaBar ?
H m=130 GeV	0.02	10^5	?
$\tilde{g}\tilde{g}$ m= 1 TeV	0.001	10^4	---
Black holes m > 3 TeV (M_D=3 TeV, n=4)	0.0001	10^3	---

LHC is a factory for anything: top, W/Z, Higgs, SUSY, etc….
mass reach for discovery of new particles up to m ~ 5 TeV

The LHC Physics Haystack(s)

Interesting cross sections



Susy

Higgs

- Cross sections for heavy particles
  ~ 1 /(1 TeV)^2
- Most have small couplings ~ α^2
- Compare with total cross section
  ~ 1/(100 MeV)^2
- Fraction ~ 1/1,000,000,000,000
- Need ~ 1,000 events for signal
- Compare needle
  ~ 1/100,000,000 m^3
- Haystack ~ 100 m^3
- Must look in ~ 100,000 haystacks

The LHC's Most Anticipated Discovery

6 December 2008

Evidence for a Higgs boson production in pp collisions at $\sqrt{s}$ = 14 TeV

CMS collaboration

Abstract

Experimental evidence for a Higgs boson production in pp collisions $\sqrt{s}$ = 14 TeV with an integrated luminosity 1 fb^{-1} [1] at the Large Hadron Collider at CERN is reported. The CMS experiment

...

Submitted to *European Journal of Physics*

A Simulated Higgs Event in CMS



A la recherche du Higgs perdu …

Higgs Production at the LHC







… not far away?





Combining direct, Indirect information

Some Sample Higgs Signals

YY

ZZ* -> llll

ttH

bbH







The Spin of the Higgs Boson @ LHC

Low mass: if H →γγ,
It cannot have spin 1

Higher mass: angular correlations in H → ZZ decays





Significance for exclusion of other J^{CP} states than 0^+

ATLAS + CMS, 2 x 300 fb^{-1}

m_H (GeV)	$J^{CP} = 1^+$	$J^{CP} = 1^-$	$J^{CP}=0^-$
200	6.5 σ	4.8 σ	40 σ
250	20 σ	19 σ	80 σ
300	23 σ	22 σ	70 σ

Measuring Higgs Self-Coupling

Light Higgs @ low-energy LC



Heavier Higgs possible @ SLHC



Higgs Detection at the LHC

The Higgs may be found quite quickly …



… in several different channels



International Linear Collider

- e^+e^- collisions up to E_{cm} = 1 TeV
- Preferred choice for next collider
- Now subject of Global Design Effort
- Hope for decision 2010 – 2012
- To be constructed by 2015 – 2020?

The GDE Plan and Schedule



Tasks for the TeV ILC

- Measure m_t to < ± 100 MeV
- If there is a light Higgs of any kind, pin it down:

  Does it have standard model couplings?

  What is its precise mass?
- If there are extra light particles:

  Measure mass and properties
- If LHC sees nothing new below ~ 500 GeV:

  Look for indirect signatures





Measuring Properties of Light Higgs

LC capabilities

Some new studies …

bb, ττ, gg, cc, WW, γγ



Measuring top-Higgs couplings



Higgs Measurements @ ILC & LHC



Measure Little Higgs Decays @ LC



Sensitivity to Strong WW scattering

@ LHC



@ 800 GeV LC



Measuring a WW Resonance

Form factor measurements @ 500 GeV LC



Resonance parameters @ LHC



Resonance parameters @ 500 GeV LC

After LHC @ CERN - CLIC?



Electron-Positron
collisions up to 3 TeV



The CLIC main parameters

Center of mass Energy (TeV)	0.5 TeV	3 TeV
Luminosity (10^{34} cm^{-1}s^{-1})	2.1	8.0
Mean energy loss (%)	4.4	21
Photons / electron	0.75	1.5
Coherent pairs per X	700	$6.8\ 10^{8}$
Rep. Rate (Hz)	200	100
10^{9} e± / bunch	4	4
Bunches / pulse	154	154
Bunch spacing (cm)	20	20
H/V ε_n (10^{-8} rad.m)	200/1	68/1
Beam size (H/V) (nm)	202/1.2	60/0.7
Bunch length (μm)	35	35
Accelerating gradient (MV/m)	150	150
Overall length (km)	10.2	33.2
Power / section (MW)	230	230
RF to beam efficciency (%)	23.1	23.1
AC to beam efficiency (%)	9.3	9.3
Total AC power for RF (MW)	105	319
Total site AC power (MW)	175	410

Large Cross Section @ CLIC

Can measure rare decay modes …





$\Delta g/g = 4\%$

$m_H = 120$ GeV



$\Delta g/g = 2\%$

$m_H = 180$ GeV

If the Higgs boson is heavier …

Can establish its existence beyond any doubt if < 1 TeV:
ee → H ee



Find resonance in strong WW scattering if > 1 TeV:
ee → H νν



Measuring Effective Higgs Potential @ Higher-Energy LC

Large cross section for HH pair production



Accuracy in measurement of HHH coupling



The Stakes in the Higgs Search

- How is gauge symmetry broken?
- Is there any elementary scalar field?
- Would have caused phase transition in the Universe when it was about 10^{-12} seconds old
- May have generated then the matter in the Universe: electroweak baryogenesis
- A related inflaton might have expanded the Universe when it was about 10^{-35} seconds old
- Contributes to today's dark energy: 10^{60} too much!