


NON-STATISTICAL EFFECTS IN BETA & GAMMA DECAYS AND BETA-DELAYED FISSION ANALYSIS



I. N. Izosimov, *Joint Institute for Nuclear Research, 141980 Dubna, Russia*



The probability $P_{\beta d}$ of β -delayed process is :

$$P_{\beta d} = \frac{Q_{\beta} \int_0^{Q_{\beta}} S_{\beta}(E) f(Q_{\beta} - E) \Gamma_d(E) / \Gamma_{\text{tot}}(E) dE}{\int_0^{Q_{\beta}} S_{\beta}(E) f(Q_{\beta} - E) dE}$$


where $\Gamma_d(E)$ – delayed process width, $\Gamma_{\text{tot}}(E)$ – total width.



The β -transition probability is proportional to the product of the lepton part described by the Fermi function $f(Q_{\beta} - E)$ and the nucleon part described by the β -decay strength function $S_{\beta}(E)$, $\Gamma_{\text{tot}} = \Gamma_d + \Gamma_{\gamma}$ and $\Gamma_d = \Gamma_f$ for delayed fission. The delayed fission probability P_{df} substantially depends on the resonance structure of the $S_{\beta}(E)$ both for β^- and β^+/EC decays. The analysis of the experimental data on delayed fission demonstrated that the P_{df} can be correctly described only by using the **non-statistical** $S_{\beta}(E)$, reflecting nuclear-structure effects (I.N. Izosimov, Yu.V. Naumov, *Bulletin of the Academy of Science USSR, Physical Series*, **42**, 25 (1978).

<https://www.researchgate.net/publication/322539669>)

In β -decay the simple (non-statistical) configurations are populated and as a consequence the non-statistical effects may be observed in γ -decay of such configurations. In delayed fission analysis the **γ -decay widths Γ_{γ} calculated using the statistical model**, which, in general, can only be an **approximation** (I.N. Izosimov, *Physics of Particles and Nuclei*, **30**, 131 (1999). DOI: 10.1134/1.953101)



Dependence of the multipole mixture δ on the incident-proton

- energy for nonanalog resonances with $I^{\pi} = \frac{3}{2}^-$ in ^{63}Cu . The excitation energies of resonances in ^{63}Cu ranged from 8040 to 9250 keV. The average value of δ is $\langle \delta \rangle = (0.6 \pm 0.1)$, while the statistical model gives $\langle \delta \rangle = 0$.

Non-statistical effects in (p, γ) nuclear reactions in the excitation and decay of the non-analog resonances, for which simple configurations play an important role, were analyzed. The strong non-statistical effects were observed both for $M1$ and $E2$ γ -transitions.

CONCLUSION

- For correct calculations of the beta-delayed processes probabilities $P_{\beta d}$ it is necessary to have experimental information and systematic both on $S_{\beta}(E)$ structure and Γ_{γ} values. Only after proper consideration of non-statistical effects both for β -decay and γ -decay it is possible to make a **quantitative** conclusion about fission barriers.
- For non-statistical effects study in delayed fission both for β -decay and γ -decay it is necessary to measure first of all the $S_{\beta}(E)$ structure for delayed fission nuclei. We are interesting in collaboration for such experiments.