



African School of Fundamental
Physics and Applications

Flavour Physics

Monica Pepe Altarelli (CERN)

ASP 20/10/2020

Outline of the lecture

- What is flavour physics and why it is interesting
- CP Violation and baryogenesis
- Some historical remarks
- The CKM Matrix
- The rise of b physics
- A brief mention of rare decays

Outline of the lecture

- What is flavour physics and why it is interesting
 - CP Violation and baryogenesis
 - Some historical remarks
 - The CKM Matrix
 - The rise of b physics
 - A brief mention of rare decays
- Some overlap with Sally Seidel's slides
 - Marco Gersabeck will go more into the experimental aspects of CPV measurements

A very vast subject..

- Flavour physics includes
 - neutrinos
 - charged leptons
 - kaon physics
 - charm & beauty physics
 - some aspects of top physics
- My focus here will be on some limited aspects of kaon, charm and beauty physics


What is flavour?



- In 1971, at a Baskin-Robbins ice-cream store in Pasadena, Murray Gell-Mann and his student Harald Fritzsch came up with the term “flavour” to describe **the different types of quarks**
- Just as ice cream has both color and flavour, so do quarks

Flavour physics refers to the study of the interactions that distinguish between the fermion generations

Who ordered that?



- ... asked I.Rabi after the discovery of the μ with a mass of $207 m_e$
- $m_t/m_u \sim \mathcal{O}(10^5)$!
- ν masses many orders of magnitude lighter than any other matter field!

The Higgs mechanism does not solve the problem of why each particle has a different mass (it does not allow us to predict/compute particle masses)

Many mysteries...

- ..even if the SM is, at the current level of experimental precision and at the energies reached so far, the most successful and best tested theory of nature at a fundamental level.

What determines the observed pattern of masses and mixing angles of quarks and leptons?


- In the SM, the only interaction distinguishing the three flavours is the Yukawa interaction (interaction of the matter fields with the Higgs boson). The complex phases present in the Yukawa couplings are also the only source of CP violation.

Are there other sources of flavour (and CP) symmetry breaking, beside the SM Yukawa couplings?

Why flavour is interesting

- To be able to answer these questions is likely to shed light on physics beyond the SM...
- Flavour physics might provide the first indications of new physics at energy scales that are beyond the reach of direct searches
- CP (Charge-Parity) violation is connected to the matter-antimatter asymmetry of the Universe

Where did the anti-matter go?



Where did the anti-matter go?

- What led to the disappearance of antimatter assuming an initial symmetric state (or that inflation washed out any possible prior asymmetry)?
 - There are anti-protons in cosmic rays, consistent with secondaries due to the interactions of cosmic-ray protons in the Interstellar Medium
 - We can produce and study anti-matter in accelerators
 - But apparently no anti-matter around us
 - This looks really strange, given that the properties of matter and antimatter are very similar.
 - **Where did it go? Why is the universe 100% matter-antimatter asymmetric ?**

Primordial Baryon Asymmetry

- We can define the **Baryon Asymmetry of the Universe (BAU)** just before antibaryons disappeared from the primordial plasma as

$$\Delta(t) = \frac{N_B - N_{\bar{B}}}{N_B + N_{\bar{B}}}$$

We already know that
 $\Delta(10^{10} \text{ years}) = 1$


- Since the end products of the annihilation processes are mostly photons and there are no antibaryons in the universe today, BAU can be estimated by the **baryon to photon ratio η**

$$\eta = \frac{N_B}{N_\gamma}$$

Primordial Baryon Asymmetry

- From observations:
 - $N_\gamma \simeq 410$ photons/cm³ (at T= 2.73⁰K)
 - $N_B \simeq 0.25$ nucleons/m³

$$\eta = \frac{N_B}{N_\gamma} \simeq 6 \times 10^{-10}$$



Small baryon-to-photon ratio in Universe today

- Conclusion is that Big Bang theory tells us that the baryon asymmetry of the early universe was a very small number , i.e., **today's huge matter-antimatter asymmetry was a tiny number in the past**

$$\Delta(t) = \frac{N_B - N_{\bar{B}}}{N_B + N_{\bar{B}}} \sim 10^{-10}$$

Beginning of Universe

10,000,000,000

matter

10,000,000,000

anti-matter

$\sim 10^{-6}$ seconds later

10,000,000,001

10,000,000,000

matter

anti-matter


Universe now



- Antimatter and matter particles annihilated massively in the early universe, but a tiny fraction of matter was left over: every 10 billion particles, a handful was not annihilated away
- We are very lucky!

Baryogenesis and Sakharov conditions

- A process called **baryogenesis** was hypothesized to generate this asymmetry dynamically from a matter-antimatter symmetric initial state
- In 1967 A.D. Sakharov enumerated three necessary conditions for baryogenesis (incidentally, his work went unnoticed for 11 years!)



Sakharov conditions

1. Baryon number violation

- Otherwise there's no way to produce an excess of baryons

2. C and CP violation

- If C and CP are exact symmetries, the total rate for any process which produces an excess of baryons is equal to the rate of the complementary process which produces an excess of antibaryons

3. Thermodynamic non equilibrium

- Otherwise any asymmetry would be washed away by simple thermodynamics

Can the SM explain baryogenesis?

- In principle SM carries all the ingredients to satisfy the Sakharov conditions
- Relevant measure is Jarlskog determinant J (I will come back to it!), an invariant that identifies CP violation in the SM and that depends on every physical quark mixing angle $J \sim \Pi(\delta m_q^2 / M_W^2) \Pi(\text{angles})$
- CP violation in the SM is proportional to J (a dimensionless quantity is constructed by dividing by the relevant temperature at which the BAU freezes out) $\sim \mathbf{10^{-20}}$
- **Many orders of magnitude below the observation!**


We need more CP violation!

- CP violation beyond the SM must exist!
- Where might we find it?
 - **quark sector** , as deviations from CKM predictions
 - **lepton sector**, e.g. as CP violation in neutrino oscillations
 - **other new physics**: almost all TEV-scale NP contains new sources of CP violation and precision measurements of flavour observables are generically sensitive to additions to the Standard Model

Some historical remarks

Cabibbo Theory

- First building block of what we now call “Flavour physics” was laid down by Nicola Cabibbo in 1963 well before many of the SM ingredients were clear
- The Cabibbo theory of semileptonic decays provided the first step towards a unified description of hadronic and leptonic weak interactions



The puzzling decays of strange particles

- $\Delta S = 1$ semileptonic weak decays (e.g. $K^+ \rightarrow \mu^+ \nu$) are suppressed relative to those with $\Delta S = 0$ (e.g. $\pi^+ \rightarrow \mu^+ \nu$)
- Cabibbo hypothesised that the weak interaction couples the up quark to an orthogonal combination of the down and strange quarks, determined by the “Cabibbo angle”

$$\begin{pmatrix} \nu_e \\ e \end{pmatrix}_L, \begin{pmatrix} \nu_\mu \\ \mu \end{pmatrix}_L, \begin{pmatrix} u \\ d' \end{pmatrix}_L = \begin{pmatrix} u \\ d \cos \theta_C + s \sin \theta_C \end{pmatrix}$$

To determine θ , let us compare the rates for $K^+ \rightarrow \mu^+ + \nu$ and $\pi^+ \rightarrow \mu^+ + \nu$; we find

$$\frac{\Gamma(K^+ \rightarrow \mu \nu)}{\Gamma(\pi^+ \rightarrow \mu \nu)} = \tan^2 \theta \frac{M_K (1 - M_\mu^2/M_K^2)^2/M_\pi (1 - M_\mu^2/M_\pi^2)^2}{M_\pi (1 - M_\mu^2/M_\pi^2)^2}. \quad (3)$$


From the experimental data, we then get^{5,6}

$$\theta = 0.257. \quad (4)$$

- The Cabibbo angle θ_C is the mixing angle expressing the weakly interacting down-quark d' in terms of fields with definite mass d, s
- Remarkable agreement of the theory with experiments, already at the time he wrote the article

GIM mechanism and charm

- However, Cabibbo's theory could not explain the suppression of strangeness-changing neutral current processes, e.g. $\frac{\Gamma(K_L \rightarrow \mu^+ \mu^-)}{\Gamma(K^+ \rightarrow \mu^+ \bar{\nu}_\mu)} \sim 10^{-8}$
- In 1970, Glashow, Iliopoulos and Maiani brought in a new, fourth, charge 2/3 quark: "charm" (small detail... not yet discovered!)
- This adds an additional decay amplitude almost identical to original one, but with opposite sign \Rightarrow (Almost) fully destructive interference
(Cancellation not perfect because u, c masses not quite the same, result proportional to $m_c^2 - m_u^2$)



- At the price of adding a second doublet, the unwanted $\Delta S = 1$ neutral currents were cancelled:



$$\begin{array}{ccc}
 & \text{weak} & \text{mass} \\
 & \text{eigenstates} & \text{eigenstates} \\
 \begin{pmatrix} d' \\ s' \end{pmatrix} & = & \begin{pmatrix} \cos \theta_C & \sin \theta_C \\ -\sin \theta_C & \cos \theta_C \end{pmatrix} \begin{pmatrix} d \\ s \end{pmatrix}
 \end{array}$$

$$\begin{pmatrix} \nu_e \\ e \end{pmatrix}_L, \begin{pmatrix} \nu_\mu \\ \mu \end{pmatrix}_L \\
 \begin{pmatrix} u \\ d' \end{pmatrix}_L, \begin{pmatrix} c \\ s' \end{pmatrix}_L$$

Remarkable symmetry between leptons and quarks!

Tremendous triumph of theory

- .. when on November 10, 1974 two groups, one at Brookhaven using a p beam on a fixed target and the other in e^+e^- at SLAC simultaneously announced the discovery of the J/Ψ resonance ($c\bar{c}$) with mass of 3.1 GeV



Ting&Richter, Nobel prize 1976

- The ADONE e^+e^- machine in Frascati was also pushed beyond its nominal limit of energy (2×1.5 GeV) and saw an overwhelming signal !

GIM-50

50 Years with the GIM Mechanism



Kobayashi and Maskawa

- With four quarks, matrix $V = \begin{pmatrix} \cos\theta & \sin\theta \\ -\sin\theta & \cos\theta \end{pmatrix}$, generally complex, can always be brought to a real form, thereby excluding CP violation from the weak interactions
- Three years later, in '73, Kobayashi and Maskawa showed that a complex phase does remain if the matrix is three by three (indicated as V_{CKM} , after Cabibbo, Kobayashi and Maskawa)
- **It is possible to incorporate the observed CP violation in a theory with six quark flavours** (remarkable conjecture when not even the second family was completed! b quark discovered in '77 by Lederman and t in '94)
- CP violation discovered in the **neutral kaon system** by Cronin and Fitch in 1964 (Nobel prize in 1980)

The neutral kaon system

- Neutral kaons $|K^0\rangle = |d\bar{s}\rangle$ and $|\bar{K}^0\rangle = |s\bar{d}\rangle$, generated in strong interactions and distinguished by their production mode, e.g., $\pi^- + p \rightarrow \Lambda + K^0$ or $p + \bar{p} \rightarrow K^+ + \bar{K}^0 + \pi^-$ (flavour eigenstates with definite quark content)
- They mix via the weak interactions \rightarrow physical states are superpositions of K^0, \bar{K}^0 (states with definite mass and lifetime)
- Weak interactions thought to be invariant under CP:


$$\text{CP eigenstates: } \begin{cases} |K_1\rangle = \frac{1}{\sqrt{2}}(|K^0\rangle - |\bar{K}^0\rangle), & CP|K_1\rangle = +|K_1\rangle \\ |K_2\rangle = \frac{1}{\sqrt{2}}(|K^0\rangle + |\bar{K}^0\rangle), & CP|K_2\rangle = -|K_2\rangle \end{cases}$$

distinguished by their mode of decay, with CP-even $K_1 \rightarrow \pi\pi$ and CP-odd $K_2 \rightarrow \pi\pi\pi$

- Large difference in lifetimes:

$$m_K - 2m_\pi \sim 220 \text{ MeV} \gg m_K - 3m_\pi \sim 80 \text{ MeV}$$




$$\Rightarrow \tau_1 \ll \tau_2$$





The Cronin & Fitch experiment

- Investigating some anomaly reported in the “regeneration” phenomenon with 2 magnetic spectrometers $\sim 20\text{m}$ away from K^0 production point ($\sim 300 K_1$ lifetimes), where only K_2 are left



- For “wrong” CP two-body decay $K_2 \rightarrow \pi\pi$, angle θ between vector sum of two momenta and beam direction should be $= 0$ and $\neq 0$ for three-body decays




The Cronin & Fitch experiment

- A clear peak of ~ 45 events in forward direction ($\cos \theta > 0.9999$) at $m^* \sim m_K$
 - Background from 3-body decays ($\pi^+\pi^-\pi^0, \pi^\pm\mu^\mp\nu_\mu, \pi^\pm e^\mp\nu_e$)
- These 45 events correspond to $K_L \rightarrow \pi^+\pi^-$ decays with BF $\sim 2 \cdot 10^{-3}$
- Observation of $K_L \rightarrow \pi^+\pi^-$ implies that K_L is not a pure CP-eigenstate
- The actual physical states are given by

$$|K_L\rangle = \frac{1}{\sqrt{1+|\epsilon|^2}} (|K_2\rangle + \epsilon|K_1\rangle) \sim |K_2\rangle$$

$$|K_S\rangle = \frac{1}{\sqrt{1+|\epsilon|^2}} (|K_1\rangle + \epsilon|K_2\rangle) \sim |K_1\rangle$$



A more modern notation (as a reference)

- $|K_s^0\rangle = p|K^0\rangle + q|\bar{K}^0\rangle$
- $|K_L^0\rangle = p|K^0\rangle - q|\bar{K}^0\rangle$ with
- $p = (1 + \epsilon)/\sqrt{2 + |\epsilon|^2}$
- $q = (1 - \epsilon)/\sqrt{2 + |\epsilon|^2}$ and
- $q/p = (1 - \epsilon)/(1 + \epsilon)$

Cabibbo-Kobayashi-Maskawa

- Generalization to 6 quarks by Kobayashi and Maskawa (1973, 10 years after Cabibbo's theory)
- CP violation introduced in a natural way if there are at least three families of quarks



Cabibbo-Kobayashi-Maskawa


- Generalization to 6 quarks by Kobayashi and Maskawa (1973, 10 years after Cabibbo's theory)
- CP violation introduced in a natural way if there are at least three families of quarks



- 2008 Nobel prize to K&M

CKM matrix

- V_{CKM} describes the rotation between flavour (d', s', b') and mass (d, s, b) eigenstates

$$\text{Flavour eigenstates} \begin{pmatrix} d' \\ s' \\ b' \end{pmatrix} = \begin{pmatrix} V_{ud} & V_{us} & V_{ub} \\ V_{cd} & V_{cs} & V_{cb} \\ V_{td} & V_{ts} & V_{tb} \end{pmatrix} \begin{pmatrix} d \\ s \\ b \end{pmatrix} \text{Mass eigenstates}$$


The diagram shows a horizontal line representing a quark transition. On the left, an arrow points right towards a vertex labeled 'b'. From this vertex, a wavy line representing a W^- boson extends to the right. At the second vertex, another arrow points right towards a vertex labeled 'u'. Below the wavy line, the label V_{ub} is written, indicating the CKM matrix element for this transition.

- V_{ij} proportional to transition amplitude from quark i to quark $j \rightarrow$
 V_{CKM} quark mixing matrix

- V_{CKM} induces flavour-changing transitions inside and between generations in the charged sector at tree level (W^\pm interaction). (By contrast, there are no flavour-changing transitions in the neutral sector at tree level. **No FCNC**)

How many independent parameters are needed to determine V_{CKM} ?

- $N \times N$ complex matrix (with $N = 3$)
- N^2 complex entries with N^2 unitarity constraints ($V^\dagger V = 1$) $\rightarrow N^2$ real parameters
- $2N - 1$ phases not physically meaningful \rightarrow
 V_{CKM} depends on $N^2 - 2N + 1 = (N - 1)^2$ real physical parameters
- An orthogonal matrix has $N(N - 1)/2$ independent parameters (mixing angles, e.g., for $N = 3$, 3 Euler angles)
- V_{CKM} has $N(N - 1)/2$ mixing angles and
 $(N - 1)^2 - N(N - 1)/2 = (N - 1)(N - 2)/2$ phases
- For $N = 2$, one mixing angle θ_c and no phases
- For $N = 3$, three angles $\theta_{12}, \theta_{13}, \theta_{23}$ and one complex phase δ

Important consequences

- If we want to see large CP-violating effects coming from the CKM matrix, we must look for processes which involve, even in leading approximation, quarks from all three generations.
- **Large CP violating asymmetries are expected in b decays!**
- CP violation in K decays is small, regardless of the value of the complex phase, because the dominant diagrams involve only quarks from the first two families

V_{CKM} parametrizations

- It can be written as product of three independent 2×2 block matrices

$$V_{CKM} = \begin{pmatrix} 1 & 0 & 0 \\ 0 & c_{23} & s_{23} \\ 0 & -s_{23} & c_{23} \end{pmatrix} \begin{pmatrix} c_{13} & 0 & s_{13}e^{-i\delta} \\ 0 & 1 & 0 \\ -s_{13}e^{i\delta} & 0 & c_{13} \end{pmatrix} \begin{pmatrix} c_{12} & s_{12} & 0 \\ -s_{12} & c_{12} & 0 \\ 0 & 0 & 1 \end{pmatrix} \quad s_{ij} = \sin \theta_{ij}, c_{ij} = \cos \theta_{ij}$$

Maiani,
Chau&Keung


- Advantage of this parametrization is that mixing angles are of different orders of magnitude. From experiment we know that

$$s_{12} \equiv \lambda, \quad s_{23} \sim \mathcal{O}(\lambda^2), \quad s_{13} \sim \mathcal{O}(\lambda^3) \quad \text{with } \lambda = \sin \theta_c \approx 0.22$$

- It is convenient to make this hierarchy more explicit, following **Wolfenstein**:
 $s_{12} = \lambda, \quad s_{23} = A\lambda^2, \quad s_{13}e^{i\delta} = A\lambda^3(\rho + i\eta)$ so that V_{CKM} can be expanded as

$$V_{CKM} = \begin{pmatrix} V_{ud} & V_{us} & V_{ub} \\ V_{cd} & V_{cs} & V_{cb} \\ V_{td} & V_{ts} & V_{tb} \end{pmatrix} \sim \begin{pmatrix} 1 - \frac{\lambda^2}{2} & \lambda & A\lambda^3(\rho - i\eta) \\ -\lambda & 1 - \frac{\lambda^2}{2} & A\lambda^2 \\ A\lambda^3(1 - \rho - i\eta) & -A\lambda^2 & 1 \end{pmatrix} + \mathcal{O}(\lambda^4)$$


Hierarchy in quark mixing



- Each quark has a preference to transform into a quark of its own generation.
- Very suggestive pattern
- No known reasons
- Completely different in neutrino sector

$$V_{CKM} = \begin{pmatrix} V_{ud} & V_{us} & V_{ub} \\ V_{cd} & V_{cs} & V_{cb} \\ V_{td} & V_{ts} & V_{tb} \end{pmatrix} \sim \begin{pmatrix} 1 - \frac{\lambda^2}{2} & \lambda & A\lambda^3(\rho - i\eta) \\ -\lambda & 1 - \frac{\lambda^2}{2} & A\lambda^2 \\ A\lambda^3(1 - \rho - i\eta) & -A\lambda^2 & 1 \end{pmatrix} + \mathcal{O}(\lambda^4)$$

Hierarchy in quark mixing



- Each quark has a preference to transform into a quark of its own generation.
- Very suggestive pattern
- No known reasons
- Completely different in neutrino sector

$$V_{CKM} = \begin{pmatrix} V_{ud} & V_{us} & V_{ub} \\ V_{cd} & V_{cs} & V_{cb} \\ V_{td} & V_{ts} & V_{tb} \end{pmatrix} \sim \begin{pmatrix} 0.97446 & 0.22452 & 0.00365 \\ 0.22438 & 0.97359 & 0.04214 \\ 0.00896 & 0.04133 & 0.999105 \end{pmatrix}$$

Unitarity conditions

- Unitarity of CKM matrix implies relations of the form

$$\sum_i V_{ij} V_{ik}^* = \delta_{j,k}, \text{ with } j \neq k$$







- Each of these 6 unitarity constraints can be seen as the sum of 3 complex numbers closing a triangle in the complex plane
- All triangles have the same area a , half of the Jarlskog invariant (independent of parametrization):

$$J = 2a = c_{12}c_{23}c_{13}^2 s_{12}s_{23}s_{13} \sin \delta \simeq \lambda^6 A^2 \eta \simeq 10^{-5}$$

- J is a measure of CPV in the SM (we introduced J in the context of baryogenesis); J equal to zero if any one of the mixing angles or phase is zero

Unitarity conditions

- Only db and ut triangles have sides of the same order (λ^3), i.e. are not squashed
- db triangle used to define angles α, β, γ (**Unitarity Triangle**)
- ut triangle of special relevance for physics of B_s mesons



Unitarity Triangle (UT)

$$V_{ud}V_{ub}^* + V_{cd}V_{cb}^* + V_{td}V_{tb}^* = 0$$

$$\mathcal{O}(\lambda^3) \quad \mathcal{O}(\lambda^3) \quad \mathcal{O}(\lambda^3)$$

$$\frac{V_{ud}V_{ub}^*}{V_{cd}V_{cb}^*} + 1 + \frac{V_{td}V_{tb}^*}{V_{cd}V_{cb}^*} = 0$$

$$\gamma = \arg\left(-\frac{V_{ud}V_{ub}^*}{V_{cd}V_{cb}^*}\right)$$

$$R_u \equiv \left| \frac{V_{ud}V_{ub}^*}{V_{cd}V_{cb}^*} \right|$$

$(\bar{\rho}, \bar{\eta})$

$\alpha = \phi_2$

$$\alpha = \arg\left(-\frac{V_{td}V_{tb}^*}{V_{ud}V_{ub}^*}\right)$$

$$R_t \equiv \left| \frac{V_{td}V_{tb}^*}{V_{cd}V_{cb}^*} \right|$$

$$\beta = \arg\left(-\frac{V_{cd}V_{cb}^*}{V_{td}V_{tb}^*}\right)$$

$\gamma = \phi_3$


$\beta = \phi_1$

$(0,0)$

$(1,0)$

- The triangle has vertices at $(0,0)$, $(1,0)$, $(\bar{\rho}, \bar{\eta})$ with $\bar{\rho} \equiv \rho(1 - \lambda^2/2)$, $\bar{\eta} \equiv \eta(1 - \lambda^2/2)$
- CP violation in the quark sector ($\bar{\eta} \neq 0$) is translated into a non flat UT
- Huge improvement in the knowledge of the CKM elements in the last decades!

15 years of $(\bar{\rho}, \bar{\eta})$ predictions



PDG2019

$$\bar{\rho} = 0.122^{+0.018}_{-0.017}$$

$$\bar{\eta} = 0.355^{+0.012}_{-0.011}$$


$$\lambda = 0.22453 \pm 0.00044$$

$$A = 0.836 \pm 0.015$$


- Black curves give contours at 68% and 95% probability (from <http://www.utfit.org>)

Consistency of CKM fits

www.utfit.org



ckmfitter.in2p3.fr



- Impressive effort from community and tremendous success of CKM paradigm!
- Constraints from many different quark transitions. Extensive measurements on K , D and B mesons performed at different experiments. These constraints depend also on theory input.
- At the current level of precision, all measurements are consistent and intersect in the apex of the UT
- New Physics effects (if there) are small!

A large experimental effort...

- Constraints coming from K mesons from. e.g., NA48 at CERN, KLOE at LNF, KTeV at FNAL
- Measurements of CKM parameters from D and B mesons pioneered by ARGUS at DESY, CLEO, and CLEO-c at CESR, Cornell, followed by the so-called B-factory experiments BaBar at SLAC and Belle at KEK
- Significant contributions from CDF and D0 at FNAL, especially on B_s^0 mesons
- All the above experiments have been terminated while Belle has been upgraded (Belle II)
- LHCb at the LHC is now dominating physics with b and c hadrons while the general purpose detectors ATLAS and CMS contribute in selected areas and Belle II is ramping up
- BESIII in China provides many results on c hadrons, NA62 at CERN and KOTO at J-Parc measure very rare Kaon decays


The rise of b physics

- An accurate test of the CKM paradigm requires extending the physics programme to heavy-flavoured hadrons, in particular to B -meson decays
- In the late 80s, studies indicated that the best source for such a physics programme was an e^+e^- collider operating at the $\Upsilon(4S)$ but in an asymmetric mode, i.e. with beams of unequal energy [Oddone 1987].
 - The $\Upsilon(4S)$ has a mass of 10.58 GeV and decays essentially into $B\bar{B}$ pairs (roughly equally to B^+B^- and $B^0\bar{B}^0$)
- The collider must also have unprecedented luminosity ($\mathcal{O}(\text{few } 10^{33})/\text{cm}^2/\text{sec}$), to provide enough B -mesons
- Two such asymmetric, high-luminosity e^+e^- colliders operating at the $\Upsilon(4S)$, so-called B -factories, were eventually built in the 1990s:
 - PEP-II at SLAC in the United States
 - KEK-B at KEK in Japan

Why asymmetric?

- At a symmetric B -factory, the small Q-value of the $\Upsilon(4S) \rightarrow B\bar{B}$ results in B -mesons almost at rest in the CM: $p \simeq 330 \text{ MeV} \rightarrow \beta\gamma \simeq 0.06$
 - $\tau_b \sim 1.5 \cdot 10^{-12} \text{ sec} \rightarrow d = \beta\gamma c\tau \sim 30 \mu\text{m}$ $\tau \sim 1/(m^5 |V_{cb}|^2)$
This is a decay length too small to be resolved by vertex detectors !
 - With an asymmetric B -factory, boost increases the decay length:
 - PEP-II collided 3.1 GeV e^+ and 9 GeV e^- head on $\rightarrow \beta\gamma = 0.56$ boost of the $\Upsilon(4S)$ and an average separation between the two B vertices of $260 \mu\text{s}$
 - KEK-B collided 3.5 GeV e^+ and 8 GeV e^- at ± 11 mrad crossing angle $\rightarrow \beta\gamma = 0.43$ boost of the $\Upsilon(4S)$ and an average separation between the two B vertices of $200 \mu\text{s}$
- You are in business!!
- This idea [Oddone] was a radical break with tradition, as it required two separate beam pipes, each with their own magnet system and a complex interaction region

KEKB and PEP-II




3.5 GeV e^+ 8 GeV e^-


3.1 GeV e^+ 9 GeV e^-

- Exceptional performance! The two machines broke any existing record of instantaneous and integrated luminosity of previous particle colliders and recorded over 10^9 $B\bar{B}$ pairs at the $\Upsilon(4S)$!

World record luminosities




Belle and BaBar




- BaBar at PEP-II and Belle at KEKB took data from 1999 to 2008 and 2010, respectively
- Each of the two experiments did an excellent job of reconstructing charged tracks and decay vertices, detecting photons even down to low energy (~ 30 MeV) and performing particle identification to reconstruct electrons, muons, pions, kaons and protons.

The golden mode: $B^0 \rightarrow J/\psi K_s^0$



- It allowed the first observation of CP violation in B decays at the B factories




- Final state f_{CP} common to both B^0 and \bar{B}^0 decays: $CP|f_{CP}\rangle = \eta_{CP}|f_{CP}\rangle$ with $\eta_{CP} = \pm 1$
- **Interference** between the amplitudes for the direct decay and that after $B^0 - \bar{B}^0$ oscillation results in a decay-time dependent CP asymmetry :

$$A_{CP}(\Delta t) = \frac{\Gamma(\bar{B}^0 \rightarrow J/\psi K_s) - \Gamma(B^0 \rightarrow J/\psi K_s)}{\Gamma(\bar{B}^0 \rightarrow J/\psi K_s) + \Gamma(B^0 \rightarrow J/\psi K_s)} \cong \sin(2\beta) \sin(\Delta m \Delta t), \text{ where}$$


$\Delta t \equiv t_{\text{rec}} - t_{\text{tag}}$ is the time difference between the two decays and Δm the mass difference between the heavy and light mass eigenstates of the $B^0 - \bar{B}^0$ system

$B^0 \rightarrow (c\bar{c})K_{S/L}^0$ at BaBar and Belle




$$A_{CP}(\Delta t) = \sin(2\beta) \sin(\Delta m_d \Delta t)$$

$B^0 \rightarrow (c\bar{c})K_{S/L}^0$ at BaBar and Belle




Legacy B-factories result
 $\sin 2\beta = 0.677 \pm 0.020$





The main actors today

ATLAS and CMS @ LHC are “General Purpose Detectors”, but can measure a few flavour observables, mainly with muons in final state





NA62 @ CERN is an experiment to measure the very rare decay $K^+ \rightarrow \pi^+ \nu \bar{\nu}$ (BF $\sim 10^{-10}$)



LHCb @ LHC and Belle II @ KEK are dedicated detectors for flavour physics performing a wide range of measurements

..plus BESIII, KOTO, Mu2e, MEG II, ..





LHCb

The LHC

The LHCb collaboration

- ~1400 members from 87 institutes in 18 countries
- ~530 publications, some with very high impact
- Main focus on heavy quark flavour...but plenty of other physics in the forward direction



The LHCb collaboration

- ~1400 members from 87 institutes in 18 countries
- ~530 publications, some with very high impact
- Main focus on heavy quark flavour...but plenty of other physics in the forward direction

CKM & CPV

EW and QCD

Rare decays

Spectroscopy



Semileptonic
decays

Ions and
fixed target


Exotica searches

Rare decays, in a nutshell

- In the SM, processes involving flavour changes between two up-type quarks (u,c,t) or between two down-type quarks (d,s,b) are forbidden at tree level and can only occur at loop level (penguin and box) → **Rare FCNCs**



- A new particle, too heavy to be produced at the LHC, can give sizeable effects when exchanged in a loop



- Strategy: use well-predicted observables to look for deviations
- Indirect approach to New Physics searches, complementary to that of ATLAS/CMS

One of the milestones of flavour programme


$$B_{(s)} \rightarrow \mu^+ \mu^-$$

- Very suppressed in the SM

- Theoretically “clean” → precisely predicted:

$$\mathcal{B}(B_s^0 \rightarrow \mu^+ \mu^-) = (3.65 \pm 0.23) \times 10^{-9} \quad (\sim 6\%)$$

$$\mathcal{B}(B^0 \rightarrow \mu^+ \mu^-) = (1.06 \pm 0.09) \times 10^{-10}$$



Bobeth et al.
PRL 112 (2014) 101801

- Sensitive to NP

- A large class of NP theories, such as SUSY, predict significantly higher values for the $B_{(s)}$ decay probability

- Very clean experimental signature

- Studied by all high-energy hadron collider experiments

$$B_{(s)} \rightarrow \mu^+ \mu^-$$

- First observation of $B_s \rightarrow \mu^+ \mu^-$ by LHCb!

PRL 118 (2017) 191801



- ATLAS, CMS, LHCb

$$B(B_s^0 \rightarrow \mu^+ \mu^-) = (2.69^{+0.37}_{-0.35}) \times 10^{-9}$$

$$B(B^0 \rightarrow \mu^+ \mu^-) < 1.9 \times 10^{-10} @ 95\% \text{ CL}$$

LHCb-CONF-2020-002
 CMS PAS BPH-20-003
 ATLAS-CONF-2020-049

$B_s \rightarrow \mu^+ \mu^-$ compatibility
 with SM of at 2.4σ level



Finding a needle in a haystack!




Very important constraints to many New Physics models

The SM stands its ground

- Sizeable effects expected in many MSSM models (cancellation of helicity suppression)


Straub, arXiv:1107.0266



The SM stands its ground

- Sizeable effects expected in many MSSM models (cancellation of helicity suppression)


Straub, arXiv:1107.0266



Now

Evidence for the decay $K^+ \rightarrow \pi^+ \nu \bar{\nu}$ from the NA62 experiment at CERN

- Another very rare decay that proceeds through electroweak box and penguin diagrams in the SM and could be sensitive to new physics



Rare FCNCs

- Precisely predicted in SM:

Buras et al, JHEP11 (2015) 033

$$BR(K^+ \rightarrow \pi^+ \nu \bar{\nu}) = (8.4 \pm 1.0) \times 10^{-11}$$


- Preliminary result shown at ICHEP2020 [R.Marchevski]:

- Observed 20 events with an expected background of 7

$$BR(K^+ \rightarrow \pi^+ \nu \bar{\nu}) = (11.0_{-3.5}^{+4.0}(\text{stat.}) \pm 0.3(\text{syst.})) \times 10^{-11}$$

- 3.5σ significance, compatible with SM within 1σ

Energy reach of various indirect precision tests of physics beyond the SM compared to direct searches



Matt Reece, DOE Basic Research Needs Study on HEP Detector R&D

A window with a view of a blue sky and white clouds. The window frame is light-colored wood or metal, and the view is a bright blue sky filled with large, fluffy white cumulus clouds. The text is centered in the middle of the window.

A window on
NP at high
scales

Take home message

- Flavour physics is very rich and is connected to many fundamental questions
 - What determines the observed pattern of masses and mixing angles of quarks and leptons?
 - Explaining the observed imbalance between matter and antimatter in the Universe requires CP violation. CP violation beyond the SM must exist! Keep on looking for deviations to the CKM theory
- A lesson from history is that new physics can show up at precision frontier before energy frontier
 - GIM mechanism before discovery of charm
 - CP violation and CKM before discovery of beauty and top
- A data-driven approach, in which we test precise SM predictions looking for discrepancies, is particularly relevant in the absence of direct collider production of new particles.
- Precise measurements of flavour observables provide a powerful way to probe for NP effects beyond the SM, complementing direct searches for NP

Supplementary material


Are there more than three generations?

- LEP operated at CERN from 1989 to 2000, and delivered e^+e^- collisions to four experiments at $\sqrt{s} \simeq M_Z$ and above
- Hadronic cross-section at the Z peak used to derive the number of light neutrino species N_ν

[LEP EW WG:
Phys. Rept. 427 (2006)]

$$\sigma_f^0(s = M_Z^2) = 12\pi \frac{\Gamma_e \Gamma_f}{M_Z^2 \Gamma_Z^2}$$

- Dependence on N_ν through:
 $\Gamma_Z = 3\Gamma_\ell + \Gamma_{\text{had}} + N_\nu \Gamma_\nu$



$$N_\nu = 2.9840 \pm 0.0082$$


Based on 17 million Z decays

Neutral meson oscillations

$$K^0 \leftrightarrow \bar{K}^0, \quad D^0 \leftrightarrow \bar{D}^0, \quad B^0 \leftrightarrow \bar{B}^0$$

- $\Delta S = 2, \quad \Delta C = 2, \quad \Delta B = 2$


Strangeness, charm and beauty are not conserved



- Formalism is the same even if difference in mass and CKM elements results in dramatically different phenomenology

- Flavour eigenstates M^0, \bar{M}^0 can mix into each other

- via short-distance (box diagrams) or long-distance processes



- Time evolution described by two-component Schrödinger equation

$$i \frac{d}{dt} \begin{pmatrix} M^0 \\ \bar{M}^0 \end{pmatrix} = H \begin{pmatrix} M^0 \\ \bar{M}^0 \end{pmatrix} = \left(M - \frac{i}{2} \Gamma \right) \begin{pmatrix} M^0 \\ \bar{M}^0 \end{pmatrix} = \begin{pmatrix} M - \frac{i}{2} \Gamma & M_{12} - \frac{i}{2} \Gamma_{12} \\ M_{12}^* - \frac{i}{2} \Gamma_{12}^* & M - \frac{i}{2} \Gamma \end{pmatrix} \begin{pmatrix} M^0 \\ \bar{M}^0 \end{pmatrix}$$

- H effective hamiltonian, M mass matrix, Γ decay matrix

Solving the Schrödinger equation

- Physical states: eigenstates of effective Hamiltonian $|M_{L,H}\rangle = p|M^0\rangle \pm q|\bar{M}^0\rangle$ with eigenvalues $\lambda_{L,H} = m_{L,H} - \frac{i}{2}\Gamma_{L,H}$
Labelled as either S,L (short, long-lived) or L,H (light, heavy) depending on values of Δm , $\Delta\Gamma$ (labels 1,2 usually reserved for CP eigenstates)

- They evolve as $|M_{H,L}(t)\rangle = e^{-im_{H,L}t} e^{-\Gamma_{H,L}t/2} |M_{H,L}(0)\rangle$

- By inverting, starting from a pure flavour eigenstate at $t = 0$, this will evolve into a superposition of $|M^0\rangle$ and $|\bar{M}^0\rangle$ (flavour oscillation):

$$|M(t)\rangle = g_+(t)|M^0\rangle + \frac{q}{p}g_-(t)|\bar{M}^0\rangle \quad g_+(t) = \frac{1}{2}e^{-iMt} \left(e^{-i\frac{1}{2}\Delta mt - \frac{1}{2}\Gamma_H t} + e^{+i\frac{1}{2}\Delta mt - \frac{1}{2}\Gamma_L t} \right)$$

$$|\bar{M}(t)\rangle = \frac{p}{q}g_-(t)|M^0\rangle + g_+(t)|\bar{M}^0\rangle \quad g_-(t) = \frac{1}{2}e^{-iMt} \left(e^{-i\frac{1}{2}\Delta mt - \frac{1}{2}\Gamma_H t} - e^{+i\frac{1}{2}\Delta mt - \frac{1}{2}\Gamma_L t} \right)$$





$$M = (M_H + M_L)/2, \Delta m = m_H - m_L$$

$$\Gamma = (\Gamma_L + \Gamma_H)/2, \Delta\Gamma = \Gamma_L - \Gamma_H$$

- Probability of measuring a state $|\bar{M}^0\rangle$ at time t starting from a pure sample of $|M^0\rangle$ particles:

$$|\langle\bar{M}^0|M^0(t)\rangle|^2 = |g_-(t)|^2 \left| \frac{p}{q} \right|^2 \quad \text{with} \quad |g_{\pm}(t)|^2 = \frac{e^{-\Gamma t}}{2} \left(\cosh \frac{\Delta\Gamma t}{2} \pm \cos \Delta m t \right)$$

Compare the mesons



Probability to observe an M^0 or \bar{M}^0 at time t starting from a pure M^0 meson



- Δm depends on rate of mixing diagram, $\Delta\Gamma$ depends on widths of decays into common final states ($K^0 \rightarrow \pi^+\pi^- \rightarrow \bar{K}^0$) (large for K^0 , small for D^0, B_d^0)
- $x = \Delta m/\Gamma$ gives the average number of oscillations before decay

	Δm → ($x = \Delta m/\Gamma$)	$\Delta\Gamma$ ($y = \Delta\Gamma/(2\Gamma)$)
K^0	large ~ 500	~ maximal ~ 1
D^0	small (0.63 ± 0.19)%	small (0.75 ± 0.12)%
B^0	medium 0.770 ± 0.008	small 0.008 ± 0.009
B_s^0	large 26.49 ± 0.29	medium 0.075 ± 0.010

- B^0 mixing, first observed by Argus in 1987, then measured precise by B factories, ..
- B_s^0 mixing first measured by CDF in 2006 and then by LHCb

$B^0 \leftrightarrow \bar{B}^0$ mixing

Phys. Lett. B719 (2013) 318




$$\Delta m_d = 0.5065 \pm 0.0019 \text{ ps}^{-1}$$

One period of B^0 oscillations
 $\Delta T \simeq 12 \text{ ps} \rightarrow$ oscillation
 frequency $\Delta m_d \simeq 0.5 \text{ ps}^{-1}$

$B_s^0 \leftrightarrow \bar{B}_s^0$ mixing

- $\Delta m_s = 17.757 \pm 0.021 \text{ ps}^{-1}$ CDF,LHCb



New J.Phys.15 (2013) 053021

One period of B_s^0 oscillations
 $\Delta T \simeq 350 \text{ fs} \rightarrow$ oscillation
frequency $\Delta m_s \simeq 17.8 \text{ ps}^{-1}$