Three-Loop Calculation of the Higgs Boson Mass in Supersymmetry

Monday 5 September 2011 14:00 (25 minutes)

A Key feature of the minimal supersymmetric extension of the Standard Model (MSSM) is the existence of a light Higgs boson, the mass of which is not a free parameter but an observable that can be predicted from the theory. Given that the LHC is able to measure the mass of a light Higgs with very good accuracy, a lot of effort has been put into a precise theoretical prediction.

We present a calculation of the SUSY-QCD corrections to this observable to three-loop order. We perform multiple asymptotic expansions in order to deal with the multi-scale three-loop diagrams, making heavy use of computer algebra and keeping a keen eye on the numerical error introduced.

We provide a computer code in the form of a Mathematica package that combines our three-loop SUSY-QCD calculation with the literature of one- and two-loop corrections to the Higgs mass, providing a state-of-the-art prediction for this important observable.

Primary author: Dr KANT, Philipp (Humboldt-Universität zu Berlin)

Co-authors: Dr MIHAILA, Luminita (KIT); Prof. STEINHAUSER, Matthias (KIT); Prof. HARLANDER, Robert V. (Bergische Universität Wuppertal)

Presenter: Dr KANT, Philipp (Humboldt-Universität zu Berlin)

Session Classification: Monday 05th - Computations in Theoretical Physics

Track Classification: Track 3: Computations in Theoretical Physics - Techniques and Methods