

# ATLAS results on charmonium production and $B_c$ production and decays



Semen Turchikhin  
*on behalf of ATLAS Collaboration*

Joint Institute for Nuclear Research






Light Cone 2021: Physics of Hadrons on the Light Front  
29 November – 4 December 2021



## ▶ Charmonium production

- ▶  $J/\psi$  and  $\psi(2S)$  production measurement at  $\sqrt{s} = 13$  TeV  
[ATLAS-CONF-2019-047](#) 
- ▶ Associated prompt  $J/\psi + W$  production measurement at  $\sqrt{s} = 8$  TeV  
[JHEP 01 \(2020\) 095](#) 

## ▶ Physics of $B_c$ meson






- ▶ Measurement of relative  $B_c^+/B^+$  production at  $\sqrt{s} = 8$  TeV  
[PRD 104 \(2021\) 012010](#) 
- ▶ Study of  $B_c^+ \rightarrow J/\psi D_s^{(*)+}$  decays with  $\sqrt{s} = 13$  TeV data  
[ATLAS-CONF-2021-046](#) 

- ▶ Heavy quarkonium is a unique probe for our understanding of strong interactions
- ▶ Two production mechanisms:
  - ▶ *Prompt* in  $pp$  interaction or feed-down from heavier states
  - ▶ *Non-prompt* from  $b$  hadron decays
  - ▶ Distinguished by 2D fit of dimuon mass and pseudo-proper lifetime
- ▶ ATLAS Run 1 results ([EPJC 76 \(2016\) 283](#) ) were well described by the conventionally used approaches: **NRQCD** (prompt) and **FONLL** (non-prompt, although slightly worse at highest  $p_T$ )
  - ▶ Used di-muon triggers, reaching only  $p_T \sim 100$  GeV
- ▶ New analysis uses full Run 2 dataset, aims at high- $p_T$  region
  - ▶ Use **single-muon trigger** with  $p_T(\mu) > 50$  GeV





# Prompt production results

- ▶  $p_T$  range goes up to 360 GeV – much beyond what was achieved before
- ▶ Non-prompt fraction stable above 60 GeV
  - ▶ cf. growth from  $\sim 0.3$  to  $\sim 0.7$  between 8 and 40 GeV



# Non-prompt production results

- ▶ FONLL agrees with data at low  $p_T$  within theory uncertainty
- ▶ Overestimates them above 150 GeV for  $J/\psi$
- ▶ not enough statistics for  $\psi(2S)$



- ▶  $W + \text{prompt } J/\psi$  production is often assumed to be a signature of colour-octet (CO)  $J/\psi$  production
  - ▶ although theorists argue that higher-order colour-singlet (CS) processes may dominate

▶ ATLAS analysis uses Run 1  $\sqrt{s} = 8$  TeV data

▶ Ratio of  $J/\psi + W$  to inclusive  $W$  production cross-section

▶  $8.5 \text{ GeV} < p_T(J/\psi) < 150 \text{ GeV}, |y(J/\psi)| < 2.1$


▶  $R_{J/\psi}^{\text{incl}} = \frac{\sigma_{\text{incl}}(pp \rightarrow J/\psi + W^\pm)}{\sigma(pp \rightarrow W^\pm)} \cdot \mathcal{B}(J/\psi \rightarrow \mu^+ \mu^-) = (5.3 \pm 0.7(\text{stat.}) \pm 0.8(\text{syst.})_{-0.7}^{+1.5} (\text{spin})) \cdot 10^{-6}$

▶ Subtraction of the DPS contribution done under two assumptions on  $\sigma_{\text{eff}}$

▶  $R_{J/\psi}^{\text{DPSsub}} = (3.6 \pm 0.7_{-1.0}^{+1.1+1.5}) \cdot 10^{-6}, \sigma_{\text{eff}} = 15.0_{-4.2}^{+5.8} \text{ mb}$



▶  $R_{J/\psi}^{\text{DPSsub}} = (1.3 \pm 0.7 \pm 1.5_{-0.7}^{+1.5}) \cdot 10^{-6}, \sigma_{\text{eff}} = 6.3 \pm 1.9 \text{ mb}$

▶ Theory predictions using CO LDME (from fits of Tevatron data) gives  $(0.428 \pm 0.017) \cdot 10^{-6}$







# $J/\psi + W$ production measurement (2)

- ▶ Excess of SPS production is more visible in differential  $J/\psi + W$  production rate at high  $p_T(J/\psi)$



- ▶ May be an indication of significant contribution of CS mechanism not accounted for in the predictions

- ▶  $B_c^+$  production involves producing  $b\bar{b}$  and  $c\bar{c}$  simultaneously
  - ▶ x-section predicted at  $\sim 0.2\%$  of inclusive  $b\bar{b}$  production
  - ▶ dominated by  $gg \rightarrow B_c^+ + b + \bar{c}$
- ▶ Measure the ratio:  $\frac{\sigma(B_c^+) \cdot \mathcal{B}(B_c^+ \rightarrow J/\psi \pi^+)}{\sigma(B^+) \cdot \mathcal{B}(B^+ \rightarrow J/\psi K^+)}$ 
  - ▶ common systematic uncertainties largely cancel
- ▶ Use Run 1 data at  $\sqrt{s} = 8 \text{ TeV}$
- ▶ Total x-section in fiducial range
  - ▶  $p_T(B) > 13 \text{ GeV}$ ,  $|y(B)| < 2.3$
- ▶ and differentially, in two bins of  $p_T(B)$  and  $|y(B)|$







# $B_c^+ / B^+$ production measurement results

- ▶ Total x-section ratio:

$$\frac{\sigma(B_c^+) \cdot \mathcal{B}(B_c^+ \rightarrow J/\psi \pi^+)}{\sigma(B^+) \cdot \mathcal{B}(B^+ \rightarrow J/\psi K^+)} = (0.34 \pm 0.04(\text{stat.})_{-0.02}^{+0.06}(\text{syst.}) \pm 0.01(\text{lifetime}))\%$$



- ▶ With  $p_T(B)$ , the  $B_c^+$  production decreases faster than  $B^+$
- ▶ No significant dependence on  $|y(B)|$  found






*Shaded areas show the inclusive measurement result, the points correspond to individual  $p_T(B)$  or  $|y(B)|$  bins*




# $B_c^+ / B^+$ production measurement results

Measurement	$p_T(B)$	$ y(B) $	Result [%]
ATLAS 8 TeV	$> 13$ GeV	$< 2.3$	$(0.34 \pm 0.04^{+0.06}_{-0.02} \pm 0.01)$
bin 1	13 – 22 GeV	$< 2.3$	$(0.44 \pm 0.07 \pm^{+0.09}_{-0.04} \pm 0.01)$
bin 2	$> 22$ GeV	$< 2.3$	$(0.24 \pm 0.04 \pm^{+0.05}_{-0.01} \pm 0.01)$
LHCb 8 TeV	$< 20$ GeV	2.0 – 4.5	$(0.683 \pm 0.018 \pm 0.009)$
CMS 7 TeV	$> 15$ GeV	$< 1.6$	$(0.48 \pm 0.05 \pm 0.03 \pm 0.05)$

- ▶ ATLAS result is lower than LHCb measurement  in forward and softer kinematics
  - ▶ the lower- $p_T$  bin is more consistent though
- ▶ Fairly consistent with CMS result at  $\sqrt{s} = 7$  TeV  in a similar kinematics



- ▶ Observed earlier by LHCb (PRD 87 (2013) 112012 ) and ATLAS (EPJC 76 (2016) 4 ) in Run 1.
- ▶ Aiming at more precise measurement of  $\mathcal{B}$ 's and polarization with full Run 2 dataset
  - ▶ Test various approaches used for their predictions, e.g. pQCD calculation , relativistic potential models , sum rules calculations  etc.
- ▶ Signal channel uses  $D_s^+ \rightarrow \phi(K^+K^-)\pi^+$  reconstruction, and  $D_s^{*+} \rightarrow D_s^+\pi^0/\gamma$  with incomplete reconstruction
- ▶ Use  $B_c^+ \rightarrow J/\psi\pi^+$  *reference channel* for  $\mathcal{B}$  measurement
- ▶ Fiducial range:  $p_T(B_c^+) > 15 \text{ GeV}$ ,  $|\eta(B_c^+)| < 2.0$



$$N_{B_c^+ \rightarrow J/\psi\pi^+} = 8440^{+550}_{-470}$$


# $B_c^+ \rightarrow J/\psi D_s^{(*)+}$ fit

- ▶ 2D fit to extract the signal parameters
  - ▶  $m(J/\psi D_s^+)$  and the  $J/\psi$  helicity angle
- ▶ Both sensitive to polarization
  - $B_c^+ \rightarrow J/\psi D_s^{*+}$
- ▶ Total yields
  - ▶  $N_{B_c^+ \rightarrow J/\psi D_s^+} = 241 \pm 28$
  - ▶  $N_{B_c^+ \rightarrow J/\psi D_s^{*+}} = 424 \pm 46$



# $B_c^+ \rightarrow J/\psi D_s^{(*)+}$ results

- ▶  $R_{D_s^+/\pi^+} = 2.76 \pm 0.33(\text{stat.}) \pm 0.29(\text{syst.}) \pm 0.16(\text{BF})$
- ▶  $R_{D_s^{*+}/\pi^+} = 5.33 \pm 0.61(\text{stat.}) \pm 0.67(\text{syst.}) \pm 0.32(\text{BF})$
- ▶  $R_{D_s^{*+}/D_s^+} = 1.93 \pm 0.24(\text{stat.}) \pm 0.10(\text{syst.})$
- ▶  $\Gamma_{\pm\pm}/\Gamma = 0.70 \pm 0.10(\text{stat.}) \pm 0.04(\text{syst.})$



- ▶ Generally agree with the earlier measurements
- ▶  $R_{D_s^{*+}/\pi^+}$  described well by the predictions
- ▶  $R_{D_s^+/\pi^+}$  and  $R_{D_s^{*+}/D_s^+}$  predictions consistently deviate from data
  - ▶ except QCD PM (PRD 61 (2000) 034012) perfectly agreeing
- ▶  $\Gamma_{\pm\pm}/\Gamma$  agrees with naive spin-counting estimate of 2/3 and larger than the dedicated predictions





- ▶ The most precise measurement of these decay parameters to date

- ▶ Charmonium precise measurements provide important input for both theory and experimental studies
- ▶ ATLAS provides wide opportunities here, still room for many other measurements with Run 2
- ▶ For  $B_c^+$  physics, Run 2 provides more than order-of-magnitude higher yields
- ▶ Lots of further possible measurements on production, decays, and spectroscopy of  $B_c$  mesons

**Stay tuned for further results!**

Backup slides





# $J/\psi$ mass–lifetime fits







# $B_c^+ \rightarrow J/\psi D_s^{(*)+}$ trigger datasets

- ▶ Two datasets defined based on triggers
  - ▶ Dedicated  $J/\psi\phi$  triggers inefficient for the reference channel but contribute much to signal
  - ▶ Used only for  $R_{D_s^{*+}/D_s^+}$  and  $\Gamma_{\pm\pm}/\Gamma$  measurements



# $J/\psi D_s^+$ angular fits in the signal mass region



# ATLAS detector and trigger for B-physics

- ▶ Inner Detector in solenoid field for reconstructing tracks and vertices ( $|\eta| < 2.5$ )
- ▶ Muon Spectrometer in toroid field for muon identification ( $|\eta| < 2.7$ )
- ▶ Trigger selection primarily bases on di-muon signature

