



Contribution ID: 3

Type: **not specified**

Flavored axions and the flavor problem

Wednesday 7 October 2020 14:25 (25 minutes)

A Peccei-Quinn (PQ) symmetry is proposed, in order to generate in the Standard Model (SM) quark sector a realistic mass matrix ansatz with five texture-zeros. Limiting our analysis to Hermitian mass matrices we show that this requires a minimum of 4 Higgs doublets. This model allows assigning values close to 1 for several Yukawa couplings, giving insight into the origin of the mass scales in the SM. Since the PQ charges are non-universal the model features Flavor-Changing Neutral Currents (FCNC) at the tree level. We calculate the FCNC couplings of the most general low-energy effective Lagrangian for the axion in a procedure valid for an arbitrary number of Higgs doublets. Finally, we report the allowed region in the parameter space obtained from the measurements of branching ratios of semileptonic meson decays. By choosing conveniently the parameters of our model, it is possible to adjust the recently reported XENON1T anomaly. Dark matter interpretations for the axion will be also discussed.

Primary authors: ROJAS, Eduardo (Universidad de Nariño); MARTINEZ, Roberto (Universidad Nacional de Colombia); Prof. GIRALDO, Yithsbey (Universidad de Nariño); Prof. SALAZAR, Juan Carlos (Universidad de Nariño)

Presenter: ROJAS, Eduardo (Universidad de Nariño)

Session Classification: MOCa