

Connecting fluctuation measurements and grand canonical susceptibilities: global conservation effects

Roman Poberezhnyuk (BITP, FIAS)



NA61/SHINE Open Seminar (virtual)

August 20, 2020


with Volodymyr Vovchenko, Oleh Savchuk, Volker Koch, Mark Gorenstein, Kirill Taradiy, Viktor Begun, Leonid Satarov, Jan Steinheimer, Horst Stoecker



FIAS Frankfurt Institute
for Advanced Studies



Fluctuations in strongly interacting matter





Fluctuations probe finer details of the (QCD) equation of state

Fluctuation measures --- Cumulants (susceptibilities) of distributions --- are sensitive to fine details of interactions, e.g., phase structure

in GCE:


$$\chi_{l_1 \dots l_N}^{Q_1 \dots Q_N} = \frac{\partial^{l_1 + \dots + l_N} (p/T^4)}{\partial (\mu_{Q_1}/T)^{l_1} \dots \partial (\mu_{Q_N}/T)^{l_N}} = \frac{{}^{\text{gce}} \kappa_{l_1 \dots l_N}^{Q_1 \dots Q_N}}{VT^3}$$

Lattice QCD (fluctuations of conserved charges)





Bazavov et al. (HotQCD), 1701.04325


Statistical models (HRG, effective QCD,...)



V.Vovchenko et al, PRC 2015




R.P. et al, PRC 2019



A.Motornenko. et al, PRC 2020

Grand canonical ensemble



coordinate space


Total system, $V=V_1+V_2$, is in the canonical ensemble (CE)

Subsystem (**observed**), $V_1 = \alpha V$

Reservoir (thermostat), V_2

$$\alpha = V_1/V$$

Grand canonical ensemble




Requirements for GCE:

- 1) $V_1 \gg \xi$ --- (Thermodynamic limit) --- ratios of extensive quantities become V -independent
- 2) $V \gg V_1$ --- (Subsystem is a small part of the system $\Leftrightarrow \alpha \ll 1$)

$$V \gg V_1 \gg \xi$$

Equivalence of ensembles



Begun, Gorenstein, Gazdzicki, Zozulya, PRC 2004

Subensemble

$$V \gg V_1 \gg \xi$$
A diagram showing the relationship between three variables: V, V1, and xi. The variables are arranged in a descending sequence: V >> V1 >> xi. Each variable is crossed out with a diagonal line. A blue arrow points from the text 'Global conservation effects' below to the V >> V1 part of the sequence. Another blue arrow points from the text 'Finite size effects' below to the V1 >> xi part of the sequence.

Global conservation effects

Finite size effects


Finite subsystem in finite reservoir corresponds to Subensemble ---
the generalization of grand canonical and canonical statistical ensembles

V.Vovchenko, O.Savchuk, **R.P.**, M.Gorenstein, V.Koch, 2003.13905 [hep-ph], 2020

R.P., O.Savchuk,, M.Gorenstein, V.Vovchenko, K.Taradiy, V.Begun, L.Satarov, J.Steinheimer, H.Stoecker
PRC, 2020

Relevant for heavy-ion collision experiments

Relativistic heavy-ion collisions (Event-by-event fluctuations)




Facilities: CERN-LHC, BNL-RHIC, CERN-SPS, FAIR-GSI, ...

$$k_2 \sim \xi^2, \quad \xi \rightarrow \infty$$

NA61/SHINE program

STAR-BES program


CP \Rightarrow "FLUCTUATION HILL"



M. Gazdzicki, CPOD2016

Event-by-event fluctuations

Cumulants are connected with moments of particle multiplicity/conserved charge distribution:

$$\begin{aligned}\kappa_1 &= \langle B \rangle \\ \kappa_2 &= \langle B^2 \rangle - \langle B \rangle^2 \\ \kappa_3 &= \langle B^3 \rangle - 3\langle B^2 \rangle \langle B \rangle + 2\langle B \rangle^3 \\ \kappa_4 &= \langle B^4 \rangle - 4\langle B^3 \rangle \langle B \rangle + 12\langle B^2 \rangle \langle B \rangle^2 - 3\langle B^2 \rangle^2 - 6\langle B \rangle^4 \\ &\dots\end{aligned}$$


(2)

Intensive (volume independent in GCE) ratios of cumulants:

$$\langle B \rangle = \kappa_1, \quad \omega = \frac{\kappa_2}{\kappa_1}, \quad S\sigma = \frac{\kappa_3}{\kappa_2}, \quad \kappa\sigma^2 = \frac{\kappa_4}{\kappa_2}.$$


also strongly intensive quantities - expressed through cumulant ratios

Grand-canonical ensemble:

$$\kappa_n = \frac{1}{VT^3} \chi_B^n(T, \mu), \quad \chi_B^n(T, \mu) = \frac{\partial^n(p/T^4)}{\partial(\mu_B/T)^n}$$

Applicability of the GCE in heavy-ion collisions

Experiments measure fluctuations in a finite momentum acceptance



V. Koch, 0810.2520


GCE applies if $\Delta Y_{total} \gg \Delta Y_{accept} \gg \Delta Y_{kick}, \Delta Y_{corr}$ and momentum-space correlation is strong (e.g. Bjorken flow)

In practice, difficult to satisfy all conditions simultaneously...

If $\Delta Y_{total} \gg \Delta Y_{accept}$ does not hold, corrections from global conservation appear

Applicability of the GCE in heavy-ion collisions

Experiments measure fluctuations in a finite momentum acceptance



V. Koch, 0810.2520

GCE applies if $\Delta Y_{\text{total}} \gg \Delta Y_{\text{accept}} \gg \Delta Y_{\text{kick}}, \Delta Y_{\text{corr}}$ and momentum-space correlation is strong (e.g. Bjorken flow)

In practice, difficult to satisfy all conditions simultaneously...

Subensemble acceptance method allows to account for global conservation effects and provides fluctuation measures insensible to global conservation!

Theory vs experiment: Caveats

- proxy observables in experiment (net-proton, net-kaon) vs actual conserved charges in QCD (net-baryon, net-strangeness)
Asakawa, Kitazawa, PRC '12; V.V., Jiang, Gorenstein, Stoecker, PRC '18
- volume fluctuations
Gorenstein, Gazdzicki, PRC '11; Skokov, Friman, Redlich, PRC '13;
Braun-Munzinger, Rustamov, Stachel, NPA '17
- non-equilibrium (memory) effects
Mukherjee, Venugopalan, Yin, PRC '15
- final-state interactions in the hadronic phase
Steinheimer, V.V., Aichelin, Bleicher, Stoecker, PLB '18
- accuracy of the grand-canonical ensemble (global conservation laws)
Jeon, Koch, PRL '00; Bzdak, Skokov, Koch, PRC '13;
Braun-Munzinger, Rustamov, Stachel, NPA '17

Theory vs experiment: Caveats

- proxy observables in experiment (net-proton, net-kaon) vs actual conserved charges in QCD (net-baryon, net-strangeness)
Asakawa, Kitazawa, PRC '12; V.V., Jiang, Gorenstein, Stoecker, PRC '18
- volume fluctuations
Gorenstein, Gazdzicki, PRC '11; Skokov, Friman, Redlich, PRC '13;
Braun-Munzinger, Rustamov, Stachel, NPA '17
- non-equilibrium (memory) effects
Mukherjee, Venugopalan, Yin, PRC '15
- final-state interactions in the hadronic phase
Steinheimer, V.V., Aichelin, Bleicher, Stoecker, PLB '18
- **accuracy of the grand-canonical ensemble (global conservation laws)**
Jeon, Koch, PRL '00; Bzdak, Skokov, Koch, PRC '13;
Braun-Munzinger, Rustamov, Stachel, NPA '17

Subensemble acceptance method


Partition a thermal system with a globally conserved charge B (*canonical ensemble*) into two subsystems which can exchange the charge

Neglect surface effects:

$$\hat{H} = \hat{H}_1 + \hat{H}_2 + \hat{V}_{1,2} \approx \hat{H}_1 + \hat{H}_2$$

The canonical partition function then reads:

$$Z^{\text{ce}}(T, V, B) = \text{Tr} e^{-\beta \hat{H}} \approx \sum_{B_1} Z^{\text{ce}}(T, V_1, B_1) Z^{\text{ce}}(T, V - V_1, B - B_1)$$



$$V_1 + V_2 = V$$

The probability to have charge B_1 is:

$$P(B_1) \propto Z^{\text{ce}}(T, \alpha V, B_1) Z^{\text{ce}}(T, (1 - \alpha)V, B - B_1), \quad \alpha \equiv V_1/V$$

If the canonical partition function known, B_1 -cumulants can be calculated explicitly

Subensemble acceptance: Thermodynamic limit

In the thermodynamic limit, $V \rightarrow \infty$, Z^{ce} expressed through free energy density

$$Z^{ce}(T, V, B) \stackrel{V \rightarrow \infty}{\simeq} \exp \left[-\frac{V}{T} f(T, \rho_B) \right]$$

B_1 cumulant generating function:

$$G_{B_1}(t) = \ln \left\{ \sum_{B_1} e^{t B_1} \exp \left[-\frac{\alpha V}{T} f(T, B_1) \right] \times \exp \left[-\frac{\beta V}{T} f(T, B_2) \right] \right\} + \tilde{C}.$$

B_1 cumulants:


$$\kappa_n[B_1] = \left. \frac{\partial^n G_{B_1}(t)}{\partial t^n} \right|_{t=0} \equiv \tilde{\kappa}_n[B_1(t)]|_{t=0} \quad \text{or} \quad \kappa_n[B_1] = \left. \frac{\partial^{n-1} \tilde{\kappa}_1[B_1(t)]}{\partial t^{n-1}} \right|_{t=0}$$

All κ_n can be calculated by determining the t -dependent first cumulant $\tilde{\kappa}_1[B_1(t)]$

Subensemble acceptance: Thermodynamic limit

$$\tilde{\kappa}_1[B_1(t)] = \frac{\sum_{B_1} B_1 \tilde{P}(B_1; t)}{\sum_{B_1} \tilde{P}(B_1; t)} \equiv \langle B_1(t) \rangle \quad \text{with} \quad \tilde{P}(B_1; t) = \exp \left\{ tB_1 - V \frac{\alpha f(T, \rho_{B_1}) + \beta f(T, \rho_{B_2})}{T} \right\}.$$

Thermodynamic limit: $\tilde{P}(B_1; t)$ highly peaked at $\langle B_1(t) \rangle$



$\langle B_1(t) \rangle$ is a solution to equation $d\tilde{P}/dB_1 = 0$:

$$t = \hat{\mu}_B[T, \rho_{B_1}(t)] - \hat{\mu}_B[T, \rho_{B_2}(t)]$$

where $\hat{\mu}_B \equiv \mu_B/T$, $\mu_B(T, \rho_B) = \partial f(T, \rho_B)/\partial \rho_B$

$t = \mathbf{0}$: $\rho_{B_1} = \rho_{B_2} = B/V$, $B_1 = \alpha B$, i.e. conserved charge uniformly distributed between the two subsystems

Subensemble acceptance: $\kappa_2[B_1]$

$$t = \hat{\mu}_B[T, \rho_{B_1}(t)] - \hat{\mu}_B[T, \rho_{B_2}(t)] \quad (*)$$

$$\rho_{B_1}(t) = \langle B_1(t) \rangle / (\alpha V) \quad \rho_{B_2}(t) = [B - \langle B_1(t) \rangle] / [(1 - \alpha)V]$$

$$\frac{\partial(*)}{\partial t} : \quad 1 = \left(\frac{\partial \hat{\mu}_B}{\partial \rho_{B_1}} \right)_T \left(\frac{\partial \rho_{B_1}}{\partial \langle B_1 \rangle} \right)_V \frac{\partial \langle B_1 \rangle}{\partial t} - \left(\frac{\partial \hat{\mu}_B}{\partial \rho_{B_2}} \right)_T \left(\frac{\partial \rho_{B_2}}{\partial \langle B_2 \rangle} \right)_V \frac{\partial \langle B_2 \rangle}{\partial \langle B_1 \rangle} \frac{\partial \langle B_1 \rangle}{\partial t}$$

$$\text{GCE susceptibilities: } [\partial \hat{\mu}_B(T, \rho_{B_{1,2}}) / \partial \rho_{B_{1,2}}]_T = [T^3 \chi_2^B(T, \rho_{B_{1,2}})]^{-1}$$

Solve the equation for $\tilde{\kappa}_2$:

$$\tilde{\kappa}_2[B_1(t)] = \frac{V T^3}{[\alpha \chi_2^B(T, \rho_{B_1})]^{-1} + [(1 - \alpha) \chi_2^B(T, \rho_{B_2})]^{-1}}$$

$t = 0$:

$$\kappa_2[B_1] = \alpha (1 - \alpha) V T^3 \chi_2^B$$

Connection between cumulant measured in the subsystem and GCE susceptibility

Higher-order cumulants: iteratively differentiate $\tilde{\kappa}_2$ w.r.t. t

Subensemble acceptance: Full result up to κ_6

$$\kappa_1[B_1] = \alpha VT^3 \chi_1^B$$

$$\kappa_2[B_1] = \alpha VT^3 \beta \chi_2^B$$

$$\kappa_3[B_1] = \alpha VT^3 \beta (1 - 2\alpha) \chi_3^B$$

$$\kappa_4[B_1] = \alpha VT^3 \beta \left[\chi_4^B - 3\alpha\beta \frac{(\chi_3^B)^2 + \chi_2^B \chi_4^B}{\chi_2^B} \right]$$

$$\kappa_5[B_1] = \alpha VT^3 \beta (1 - 2\alpha) \left\{ [1 - 2\beta\alpha] \chi_5^B - 10\alpha\beta \frac{\chi_3^B \chi_4^B}{\chi_2^B} \right\}$$

$$\kappa_6[B_1] = \alpha VT^3 \beta [1 - 5\alpha\beta(1 - \alpha\beta)] \chi_6^B + 5 VT^3 \alpha^2 \beta^2 \left\{ 9\alpha\beta \frac{(\chi_3^B)^2 \chi_4^B}{(\chi_2^B)^2} - 3\alpha\beta \frac{(\chi_3^B)^4}{(\chi_2^B)^3} - 2(1 - 2\alpha)^2 \frac{(\chi_4^B)^2}{\chi_2^B} - 3[1 - 3\beta\alpha] \frac{\chi_3^B \chi_5^B}{\chi_2^B} \right\}$$

$$\beta = 1 - \alpha$$

$$\chi_n^B = \frac{\partial^n(p/T^4)}{\partial(\mu_B/T)^n} \quad - \text{grand-canonical susceptibilities}$$

Model-independent!

Subensemble acceptance: Cumulant ratios

Some common cumulant ratios:

scaled variance $\frac{\kappa_2[B_1]}{\kappa_1[B_1]} = (1 - \alpha) \frac{\chi_2^B}{\chi_1^B},$

skewness $\frac{\kappa_3[B_1]}{\kappa_2[B_1]} = (1 - 2\alpha) \frac{\chi_3^B}{\chi_2^B},$

kurtosis $\frac{\kappa_4[B_1]}{\kappa_2[B_1]} = (1 - 3\alpha\beta) \frac{\chi_4^B}{\chi_2^B} - 3\alpha\beta \left(\frac{\chi_3^B}{\chi_2^B} \right)^2.$

Subensemble acceptance: Cumulant ratios

Some common cumulant ratios:


scaled variance $\frac{\kappa_2[B_1]}{\kappa_1[B_1]} = (1 - \alpha) \frac{\chi_2^B}{\chi_1^B},$

skewness $\frac{\kappa_3[B_1]}{\kappa_2[B_1]} = (1 - 2\alpha) \frac{\chi_3^B}{\chi_2^B},$

kurtosis $\frac{\kappa_4[B_1]}{\kappa_2[B_1]} = (1 - 3\alpha\beta) \frac{\chi_4^B}{\chi_2^B} - 3\alpha\beta \left(\frac{\chi_3^B}{\chi_2^B} \right)^2.$

- Global conservation (α) and equation of state (χ_n^B) effects factorize in cumulants up to the 3rd order, starting from κ_4 not anymore
- $\alpha \rightarrow 0$ – GCE limit
- $\alpha \rightarrow 1$ – CE limit
- Charge conservation suppresses the magnitude of scaled variance and skewness
- For ideal uncorrelated gas all cumulant ratios equal to 1 and subensemble acceptance is reduced to binomial acceptance


Binomial acceptance vs actual acceptance



Binomial acceptance: accept each particle (charge) with a probability α independently from all other particles

The binomial acceptance does not account for correlations between particles

Subensemble:





Accounts for correlations, connects measurements with GCE susceptibilities

Subensemble acceptance: van der Waals fluid

van der Waals equation of state: first-order phase transition and a **critical point**

$$Z_{\text{vdW}}^{\text{ce}}(T, V, N) = Z_{\text{id}}^{\text{ce}}(T, V - bN, N) \exp\left(\frac{aN^2}{VT}\right) \theta(V - bN)$$

Rich structures in cumulant ratios close to the CP





Vovchenko, R.P., Anchishkin, Gorenstein, 1507.06537

Subensemble acceptance: van der Waals fluid

Calculate cumulants $\kappa_n [N]$ in a subvolume directly from the partition function

$$P(N) \propto Z_{\text{vdW}}^{\text{ce}}(T, xV_0, N) Z_{\text{vdW}}^{\text{ce}}(T, (1-x)V_0, N_0 - N)$$

and compare with the subensemble acceptance results




Results agree with subensemble acceptance in thermodynamic limit ($N_0 \rightarrow \infty$)


Finite size effects are strong near the critical point: a consequence of large correlation length ξ

Global conservation effects at $\alpha = 0.4$

$$\frac{\kappa_4[B_1]}{\kappa_2[B_1]} = (1 - 3\alpha\beta) \frac{\chi_4^B}{\chi_2^B} - 3\alpha\beta \left(\frac{\chi_3^B}{\chi_2^B} \right)^2.$$





GCE ($\alpha \rightarrow 0$)



$\alpha = 0.4$

Net baryon fluctuations at LHC and top RHIC

$$\left(\frac{\kappa_4}{\kappa_2}\right)_{LHC} = (1 - 3\alpha\beta) \frac{\chi_4^B}{\chi_2^B} \quad \left(\frac{\kappa_6}{\kappa_2}\right)_{LHC} = [1 - 5\alpha\beta(1 - \alpha\beta)] \frac{\chi_6^B}{\chi_2^B} - 10\alpha(1 - 2\alpha)^2\beta \left(\frac{\chi_4^B}{\chi_2^B}\right)^2$$



Lattice data for χ_4^B/χ_2^B and χ_6^B/χ_2^B from [Borsanyi et al., 1805.04445](#)

For $\alpha > 0.2$ difficult to distinguish effects of the EoS and baryon conservation in χ_6^B/χ_2^B , $\alpha \leq 0.1$ is a sweet spot where measurements are mainly sensitive to the EoS

Estimates: $\alpha \approx 0.1$ corresponds to $\Delta Y_{acc} \approx 2(1)$ at LHC (RHIC)

Multiple conserved charges

Let us have a vector $Q = (Q_1, \dots, Q_N)$ of N independent conserved charges in the system.

NOTATIONS

Diagonal and off-diagonal GCE susceptibilities:

$$\chi_{l_1 \dots l_N}^{Q_1 \dots Q_N} = \frac{\partial^{l_1 + \dots + l_N} (p/T^4)}{\partial(\mu_{Q_1}/T)^{l_1} \dots \partial(\mu_{Q_N}/T)^{l_N}}, \quad l_1 + \dots + l_N = M.$$

Other notation:

$$\hat{\chi}_{i_1 \dots i_M} = \frac{\partial^M (p/T^4)}{\partial(\mu_{i_1}/T) \dots \partial(\mu_{i_M}/T)} = \frac{\hat{\kappa}_{i_1 \dots i_M}^{\text{gce}}}{VT^3}, \quad i_j \in 1 \dots N.$$

$$\chi_4^B \equiv \hat{\chi}_{1111},$$

$$\chi_{211}^{BQS} \equiv \hat{\chi}_{1123} = \hat{\chi}_{1132} = \dots = \hat{\chi}_{3211}.$$

<https://github.com/vlvovch/SAM>

$$Z(T, V, \hat{Q}) = \sum_{\hat{Q}^1} Z(T, \alpha V, \hat{Q}^1) Z(T, \beta V, \hat{Q} - \hat{Q}^1)$$

Special cases

$$\beta \equiv 1 - \alpha$$

Two conserved charges: B,Q

$$\kappa_4[B^1] = \alpha VT^3 \beta \left[(1 - 3\alpha\beta) \chi_4^B - 3\alpha\beta \frac{(\chi_3^B)^2 \chi_2^Q - 2\chi_{21}^{BQ} \chi_{11}^{BQ} \chi_3^B + (\chi_{21}^{BQ})^2 \chi_2^B}{\chi_2^B \chi_2^Q - (\chi_{11}^{BQ})^2} \right].$$

Three conserved charges: B,Q,S

$$\begin{aligned} \kappa_4[B^1] = \alpha VT^3 \beta \left[(1 - 3\alpha\beta) \chi_4^B - \frac{3\alpha\beta}{D[\hat{\chi}_2]} \times \right. \\ \left. \left\{ (\chi_3^B)^2 [\chi_2^Q \chi_2^S - (\chi_{11}^{QS})^2] + (\chi_{21}^{BQ})^2 [\chi_2^B \chi_2^S - (\chi_{11}^{BS})^2] + (\chi_{21}^{BS})^2 [\chi_2^B \chi_2^Q - (\chi_{11}^{BQ})^2] \right. \right. \\ \left. \left. - 2\chi_3^B \chi_{21}^{BQ} (\chi_2^S \chi_{11}^{BQ} - \chi_{11}^{BS} \chi_{11}^{QS}) - 2\chi_3^B \chi_{21}^{BS} (\chi_2^Q \chi_{11}^{BS} - \chi_{11}^{BQ} \chi_{11}^{QS}) \right\} \right]. \end{aligned}$$

The fact that electric (strange) charge is fixed has an effect on observables which do not involve explicitly the electric charge.

But only for fluctuations of non-conserved quantities and for higher order fluctuations of conserved charges

General formulas

$$\hat{\kappa}_{i_1}[\hat{Q}^1] = \alpha VT^3 \hat{\chi}_{i_1}$$

$$\hat{\kappa}_{i_1 i_2}[\hat{Q}^1] = \alpha VT^3 \beta \hat{\chi}_{i_1 i_2}$$

$$\hat{\kappa}_{i_1 i_2 i_3}[\hat{Q}^1] = \alpha VT^3 \beta (1 - 2\alpha) \hat{\chi}_{i_1 i_2 i_3}$$

$$\hat{\kappa}_{i_1 i_2 i_3 i_4}[\hat{Q}^1] = \alpha VT^3 \beta \left[(1 - 3\alpha\beta) \hat{\chi}_{i_1 i_2 i_3 i_4} - \frac{\alpha\beta}{2! 2! 2!} \sum_{\sigma \in S_4} \hat{\chi}_{b_1 b_2}^{-1} \hat{\chi}_{i_{\sigma_1} i_{\sigma_2} b_1} \hat{\chi}_{i_{\sigma_3} i_{\sigma_4} b_2} \right]$$

For results up to the 6-th order see [arXiv:2007.03850](https://arxiv.org/abs/2007.03850)

General formulas

$$\hat{\kappa}_{i_1}[\hat{Q}^1] = \alpha VT^3 \hat{\chi}_{i_1}$$

$$\hat{\kappa}_{i_1 i_2}[\hat{Q}^1] = \alpha VT^3 \beta \hat{\chi}_{i_1 i_2}$$

$$\hat{\kappa}_{i_1 i_2 i_3}[\hat{Q}^1] = \alpha VT^3 \beta (1 - 2\alpha) \hat{\chi}_{i_1 i_2 i_3}$$

$$\hat{\kappa}_{i_1 i_2 i_3 i_4}[\hat{Q}^1] = \alpha VT^3 \beta \left[(1 - 3\alpha\beta) \hat{\chi}_{i_1 i_2 i_3 i_4} - \frac{\alpha\beta}{2! 2! 2!} \sum_{\sigma \in S_4} \hat{\chi}_{b_1 b_2}^{-1} \hat{\chi}^{i_{\sigma_1} i_{\sigma_2} b_1} \hat{\chi}^{i_{\sigma_3} i_{\sigma_4} b_2} \right]$$

Cumulants up to 3-d order have the same simple α -dependence as in the case of single conserved charge

For results up to the 6-th order see [arXiv:2007.03850](https://arxiv.org/abs/2007.03850)

General formulas

$$\begin{aligned}\hat{\kappa}_{i_1}[\hat{Q}^1] &= \alpha VT^3 \hat{\chi}_{i_1} \\ \hat{\kappa}_{i_1 i_2}[\hat{Q}^1] &= \alpha VT^3 \beta \hat{\chi}_{i_1 i_2} \\ \hat{\kappa}_{i_1 i_2 i_3}[\hat{Q}^1] &= \alpha VT^3 \beta (1 - 2\alpha) \hat{\chi}_{i_1 i_2 i_3}\end{aligned}$$

Cumulants up to third order have the same simple α -dependence as in the case of single conserved charge

$$\hat{\kappa}_{i_1 i_2 i_3 i_4}[\hat{Q}^1] = \alpha VT^3 \beta \left[(1 - 3\alpha\beta) \hat{\chi}_{i_1 i_2 i_3 i_4} - \frac{\alpha\beta}{2! 2! 2!} \sum_{\sigma \in S_4} \hat{\chi}_{b_1 b_2}^{-1} \hat{\chi}^{i_{\sigma_1} i_{\sigma_2} b_1} \hat{\chi}^{i_{\sigma_3} i_{\sigma_4} b_2} \right]$$


For results up to the 6-th order see [arXiv:2007.03850](https://arxiv.org/abs/2007.03850)

The global conservation effects cancel out in any ratio of second order cumulants and in any ratio of third order cumulants

$$\frac{\kappa_2^Q}{\kappa_2^B} = \frac{\chi_2^Q}{\chi_2^B}, \quad \frac{\kappa_3^Q}{\kappa_3^B} = \frac{\chi_3^Q}{\chi_3^B}, \quad \frac{\kappa_2^{BQ}}{\kappa_2^S} = \frac{\chi_2^{BQ}}{\chi_2^S}, \dots$$



Ensemble-independent fluctuation measures, not sensitive to global conservation

Check in Hadron Resonance Gas




Two sources of correlations:

- 1) Global conservation: different hadron species do carry different values of B, Q, S charges
- 2) Resonance decays





Check in Hadron Resonance Gas (third order)




Two sources of correlations:

- 1) Global conservation: different hadron species do carry different values of B, Q, S charges
- 2) Resonance decays



Check in Hadron Resonance Gas (higher order)



B and S charge conservation do influence the higher order fluctuations of electric charge.

The effect is stronger for non-conserved quantities, even higher cumulants of conserved charges

LHC energies

All odd-order GCE susceptibilities vanish

$$\kappa_4[B^1]|_{\hat{\mu}=0} = \alpha VT^3 \beta (1 - 3\alpha\beta) \chi_4^B$$

Now cumulants up to fifth order have the same simple α -dependence as for single conserved charge.

$$\begin{aligned} \kappa_6[B^1]|_{\hat{\mu}=0} = \alpha VT^3 \beta & \left[(1 - 5\alpha\beta(1 - \alpha\beta))\chi_6^B - \frac{10\alpha\beta(1 - 2\alpha)^2}{D[\hat{\chi}_2]} \times \right. \\ & \left\{ (\chi_{31}^{BS})^2 [\chi_2^B \chi_2^Q - (\chi_{11}^{BQ})^2] + (\chi_{31}^{BQ})^2 [\chi_2^B \chi_2^S - (\chi_{11}^{BS})^2] \right. \\ & + (\chi_4^B)^2 [\chi_2^Q \chi_2^S - (\chi_{11}^{QS})^2] + 2\chi_{31}^{BS} \chi_{31}^{BQ} (\chi_{11}^{BS} \chi_{11}^{BQ} - \chi_2^B \chi_{11}^{QS}) \\ & \left. \left. + 2\chi_{31}^{BS} \chi_4^B (\chi_{11}^{QS} \chi_{11}^{BQ} - \chi_2^Q \chi_{11}^{BS}) + 2\chi_{31}^{BQ} \chi_4^B (\chi_{11}^{QS} \chi_{11}^{BS} - \chi_2^B \chi_{11}^{BQ}) \right\} \right]. \end{aligned}$$

Strongly intensive fluctuation measures

$$\Delta[Q_a, Q_b] = C_{\Delta}^{-1} \left\{ \kappa_1[Q_b] \frac{\kappa_2[Q_a]}{\kappa_1[Q_a]} - \kappa_1[Q_a] \frac{\kappa_2[Q_b]}{\kappa_1[Q_b]} \right\},$$

$$\Sigma[Q_a, Q_b] = C_{\Sigma}^{-1} \left\{ \kappa_1[Q_b] \frac{\kappa_2[Q_a]}{\kappa_1[Q_a]} + \kappa_1[Q_a] \frac{\kappa_2[Q_b]}{\kappa_1[Q_b]} - 2 \kappa_{1,1}[Q_a, Q_b] \right\}.$$

M.Gorenstein, M.Gazdzicki, PRC, 2011

$$\Delta[Q_a, Q_b] = C_{\Delta}^{-1} VT^3 \alpha(1 - \alpha) \left\{ \chi_1^{Q_b} \frac{\chi_2^{Q_a}}{\chi_1^{Q_a}} - \chi_1^{Q_a} \frac{\chi_2^{Q_b}}{\chi_1^{Q_b}} \right\},$$

$$\Sigma[Q_a, Q_b] = C_{\Sigma}^{-1} VT^3 \alpha(1 - \alpha) \left\{ \chi_1^{Q_b} \frac{\chi_2^{Q_a}}{\chi_1^{Q_a}} + \chi_1^{Q_a} \frac{\chi_2^{Q_b}}{\chi_1^{Q_b}} - 2 \chi_{1,1}^{Q_a Q_b} \right\}.$$

$$\frac{\Sigma[Q_a, Q_b]}{\Delta[Q_a, Q_b]} = \frac{\chi_1^{\Delta}}{\chi_1^{\Sigma}} \frac{\chi_1^{Q_b} \frac{\chi_2^{Q_a}}{\chi_1^{Q_a}} + \chi_1^{Q_a} \frac{\chi_2^{Q_b}}{\chi_1^{Q_b}} - 2 \chi_{1,1}^{Q_a Q_b}}{\chi_1^{Q_b} \frac{\chi_2^{Q_a}}{\chi_1^{Q_a}} - \chi_1^{Q_a} \frac{\chi_2^{Q_b}}{\chi_1^{Q_b}}}.$$

Both strongly intensive and ensemble independent (insensitive to global conservation)

Non-conserved quantities (net-proton, pion, kaon,...)

Example: net proton number

$$Z(T, V, \hat{Q}) = \sum_{N_p} W(T, V, \hat{Q}; N_p) .$$


Variance of non-conserved quantity:

$$\check{\chi} = \begin{pmatrix} \chi_{\hat{Q}_i \hat{Q}_j} & \chi_{\hat{Q}_i p} \\ \chi_{p \hat{Q}_j} & \chi_{pp} \end{pmatrix}, \quad i, j = 1, \dots, N.$$

$$\kappa_{pp} = \alpha VT^3 \left[(1 - \alpha) \chi_{pp} + \alpha \frac{\det \check{\chi}}{\det \chi} \right]$$

GCE susceptibility


CE susceptibility





Off-diagonal cumulants involving non-conserved quantity

$$\kappa_{p\hat{Q}_j} = \alpha VT^3 \beta \chi_{p\hat{Q}_j} .$$

$$\frac{\kappa_{p\hat{Q}_j}}{\kappa_{\hat{Q}_i\hat{Q}_j}} = \frac{\chi_{p\hat{Q}_j}}{\chi_{\hat{Q}_i\hat{Q}_j}} \quad \frac{\kappa_{p\hat{Q}_j}}{\kappa_{k\hat{Q}_i}} = \frac{\chi_{p\hat{Q}_j}}{\chi_{k\hat{Q}_i}} .$$



Net-proton and net- Λ fluctuations



$$\alpha_B^p = \frac{\langle N_p^{\text{acc}} \rangle + \langle N_{\bar{p}}^{\text{acc}} \rangle}{\langle N_B^{4\pi} \rangle + \langle N_{\bar{B}}^{4\pi} \rangle}$$

Summary

- Subensemble acceptance method (SAM) provides general formulas to correct cumulants of distributions in heavy-ion collisions for global conservation of all QCD charges
- Formulas connect cumulants measured in the subsystem of the thermal system with grand canonical susceptibilities. The method works for an arbitrary equation of state and sufficiently large systems, such as created in central collisions of heavy ions.
- Some fluctuation measures are insensitive to global conservation and, thus, can be especially convenient for experimental measurement:

$$\frac{\kappa_2^Q}{\kappa_2^B}, \quad \frac{\kappa_3^Q}{\kappa_3^B}, \quad \frac{\kappa_2^{BQ}}{\kappa_2^S}, \quad \frac{\kappa_{p\hat{Q}_j}}{\kappa_{\hat{Q}_i\hat{Q}_j}}, \quad \frac{\kappa_{p\hat{Q}_j}}{\kappa_{p\hat{Q}_i}}, \quad \frac{\kappa_{p\hat{Q}_j}}{\kappa_{k\hat{Q}_i}}, \quad \frac{\Sigma[Q_a, Q_b]}{\Delta[Q_a, Q_b]}$$

Summary

- Subensemble acceptance method (SAM) provides general formulas to correct cumulants of distributions in heavy-ion collisions for global conservation of all QCD charges
- Formulas connect cumulants measured in the subsystem of the thermal system with grand canonical susceptibilities. The method works for an arbitrary equation of state and sufficiently large systems, such as created in central collisions of heavy ions.
- Some fluctuation measures are insensitive to global conservation and, thus, are can be especially convenient for experimental measurement:

$$\frac{\kappa_2^Q}{\kappa_2^B}, \quad \frac{\kappa_3^Q}{\kappa_3^B}, \quad \frac{\kappa_2^{BQ}}{\kappa_2^S}, \quad \frac{\kappa_{p\hat{Q}_j}}{\kappa_{\hat{Q}_i\hat{Q}_j}}, \quad \frac{\kappa_{p\hat{Q}_j}}{\kappa_{p\hat{Q}_i}}, \quad \frac{\kappa_{p\hat{Q}_j}}{\kappa_{k\hat{Q}_i}}, \quad \frac{\Sigma[Q_a, Q_b]}{\Delta[Q_a, Q_b]}$$

Outlook: Application of method to intermediate collision energies (NA61/SHINE, RHIC-BES,...). Effects of thermal smearing and “imperfect” space-momentum correlations, net proton higher order fluctuations.

Summary

- Subensemble acceptance method (SAM) provides general formulas to correct cumulants of distributions in heavy-ion collisions for global conservation of all QCD charges
- Formulas connect cumulants measured in the subsystem of the thermal system with grand canonical susceptibilities. The method works for an arbitrary equation of state and sufficiently large systems, such as created in central collisions of heavy ions.
- Some fluctuation measures are insensitive to global conservation and, thus, are can be especially convenient for experimental measurement:

$$\frac{\kappa_2^Q}{\kappa_2^B}, \quad \frac{\kappa_3^Q}{\kappa_3^B}, \quad \frac{\kappa_2^{BQ}}{\kappa_2^S}, \quad \frac{\kappa_{p\hat{Q}_j}}{\kappa_{\hat{Q}_i\hat{Q}_j}}, \quad \frac{\kappa_{p\hat{Q}_j}}{\kappa_{p\hat{Q}_i}}, \quad \frac{\kappa_{p\hat{Q}_j}}{\kappa_{k\hat{Q}_i}}, \quad \frac{\Sigma[Q_a, Q_b]}{\Delta[Q_a, Q_b]}$$

Outlook: Application of method to intermediate collision energies (NA61/SHINE, RHIC-BES,...). Effects of thermal smearing and “imperfect” space-momentum correlations, net proton higher order fluctuations.

Thank you for attention!