

## **Transonic accretion flow with time-dependent boundary conditions in HARM.**

*Thursday, 24 September 2020 16:55 (20 minutes)*

Accretion disks in High mass X-ray binaries (HMXB's) are mostly fed by the stellar wind from their companion star. These winds also affect the observed X-ray spectra arising from the hot coronal flow.

Cygnus X-1 and its companion star, HDE-226868 is one of such HMXBs. It is one of the brightest X-ray sources observed and shows the X-ray intensity variations in both the soft and hard X-rays. I will present my recent work on 2D numerical modeling using GRMHD code - HARM, replicating such focused, clumpy wind from the binary companion fed for accretion onto the black hole. We model an inviscid, non-magnetized, transonic accretion flow with a low angular momentum profile. I will discuss my prescribed time-dependent boundary conditions in this code and how it affects the hydrodynamics of the flow in the relativistic framework.

**Primary authors:** PALIT, Ishika (Center for Theoretical Physics, PAS); Prof. JANIUK, Agnieszka (Center for Theoretical Physics PAS); Prof. CZERNY, Bozena (Center for Theoretical Physics PAS)

**Presenter:** PALIT, Ishika (Center for Theoretical Physics, PAS)

**Session Classification:** Relativistic Astrophysics