Contribution ID: 24 Type: selected talk

## Mutual interaction of binary black hole and misaligned circumbinary disk

Thursday, 24 September 2020 16:15 (20 minutes)

The talk will cover the recent numerical investigation of a system composed of a Supermassive Black Hole Binary (SMBHB) and a non-self-gravitating, thin, locally isothermal, viscous disk.

The evolution of such a configuration is relevant not only for the expected gravitational-wave signal, but also for electromagnetic searches for SMBHB candidates. In 2-dimensional, Newtonian, numerical simulations, we analyze the influence of the two model parameters: q — the *mass ratio* of the binary and  $\iota$  — the *inclination angle* between the binary and the disk. We found that configurations with relatively low mass ratio, composed of central mass and satellite mass, always settle down in a *quasi steady state*. On the other hand, configurations characterized by equal or comparable masses may manifest an inability to reach quasi steady state for inclinations  $\iota \in (20^\circ, 55^\circ)$ . This problem does not exist for moderately inclined or highly inclined configurations, i.e., inclinations  $\iota \leq 20^\circ$  or  $\iota \geq 55^\circ$ . We try to understand the nature of these phenomena by investigating the binary and viscous torque densities which determine the disk's final density distribution and, in particular, the size of the *central gap*.

**Primary author:** PIRÓG, Michał (West Virginia University)

Co-authors: Dr MCWILLIAMS, Sean (West Virginia University); Mr MAHESH, Siddharth (West Virginia

University)

**Presenter:** PIRÓG, Michał (West Virginia University) **Session Classification:** Relativistic Astrophysics