



Contribution ID: 24

Type: **Parallel session talk**

Studying electron-positron annihilation into $K_S K \pi$ and $K_S K \pi \pi^0$ with the CMD-3 detector

Tuesday, January 11, 2022 3:20 PM (20 minutes)

Studying electron-positron annihilation into $K_S K \pi$ and $K_S K \pi \pi^0$ with the CMD-3 detector

A. A. Uskov.

Budker Institute of Nuclear Physics, Siberian Branch of the Russian Academy of Sciences.

We studied the process $e^+e^- \rightarrow K K \pi$ with the CMD-3 detector at the electron-positron collider VEPP-2000. The statistics collected by the CMD-3 detector in the energy range of 1.2 – 2 GeV during the 2011, 2012, 2017, 2019 runs, with a total luminosity integral of $\sim 120 pb^{-1}$, was used for the analysis.

The measured cross-section is crucial for the physics of light hadrons from u, d, s quarks, clarifying the hadronic contribution to the anomalous magnetic moment of the muon $(g - 2)_\mu$, and independently measuring the resonance parameters of both $\phi(1680)$, $\rho(1450)$. The intermediate dynamics of this process is also of interest, allowing us to check the isotopic relations and prove the dominance of the neutral $K^*(892)$ channel.

We developed a novel methodology for selecting signal events. Including multi-staged kinematic reconstruction. By adding the second stage, we reduced the background ~ 5 more times. This stage depends on a rigorous study of the background. Since we identified - the main physical background to be the process $e^+e^- \rightarrow 4\pi$.

The outline of the designed steps:

1. Kinematic reconstruction with 4 charged tracks.
2. Track combination, to distinguish the K_S meson by invariant mass and decay vertex.
3. Restriction on the energy of γ not bound to tracks.
4. Final selection of signal events based on analysis of 2D-distribution of four-track events by energy imbalance ΔE and momentum vector sum modulus of all four particles Δp .
5. Simulation to find efficiency ε and estimate systematic errors.

So, as a result of this work we not only designed a new selection algorithm, but also studied the theoretical aspect of the cross-section in the Vector Meson Dominance Framework. Therefore, we improved the world precision of $\phi(1680)$, $\rho(1450)$ and measured the cross-section of $e^+e^- \rightarrow K K \pi$ with the best accuracy.

Primary author: USKOV, Artem (Budker Institute of Nuclear Physics)

Presenter: USKOV, Artem (Budker Institute of Nuclear Physics)

Session Classification: Precision SM Measurements

Track Classification: Standard Model