

Probing dark matter at the ILC

Wojciech Kotlarski¹ and Aleksander Filip Żarnecki^{2*}

on behalf of the ILC International Development Team Physics and Detector Working Group

¹*National Centre for Nuclear Research, Warsaw, Poland*

²*Faculty of Physics, University of Warsaw, Poland*

**filip.zarnecki@fuw.edu.pl*

April 15, 2022

Abstract

The International Linear Collider (ILC) offers a number of unique opportunities for searching dark sector particles in e^+e^- collisions. The collider program itself will offer important capabilities, including precision measurements in the Higgs sector and direct searches based on the mono-photon signature. With extreme intensities of the high-energy electron and positron beams, the ILC will also enable new fixed-target measurements, both beam dump experiments and dedicated experiments using extracted beams. Described in this contribution are the expectations for these programs, which address various possible dark sector portals.

Presented at the 30th International Symposium
on Lepton Photon Interactions at High Energies,
hosted by the University of Manchester,
10-14 January 2022.

1 Introduction

While there are many hints for existence of Dark Matter (DM), its nature is not known. Many possible scenarios need to be considered with different possible connections between Standard Model (SM) and DM sectors, a wide range of DM particle masses, coupling values and other model parameters. The ILC, with its high energy reach, clean environment and kinematic constraints of e^+e^- collisions, is a unique machine offering many options for DM searches. Additional capabilities may become available with the use of single high-intensity electron or positron beams in a fixed target mode.

The ILC will make use of the superconducting accelerating cavities. In the assumed staged construction scheme, it will start as a 250 GeV Higgs factory, followed by 500 GeV collisions, with a possible upgrade to 1 TeV. The unique feature of the ILC is the possibility of having both electron and positron beams polarised. This is crucial for many precision measurements as well as for beyond SM searches. By combining data collected with four different polarisation combinations accuracy of precision measurements is significantly increased, more constraints can be imposed in global fits and analyses, ambiguity between different scenarios can be removed in many BSM studies and the sensitivity to systematic effects can be significantly reduced.

This is an updated version of the contribution previously presented at [1, 2].

2 Higgs portal

Invisible decays of the Higgs boson are considered as a possible evidence for existence of DM particles. In Higgs-portal models, new scalar fields coupled to DM particles mix with the SM Higgs field. For small mixing angle θ , one of the mass eigenstates is SM-like (the observed 125 GeV scalar) and its couplings can show no deviations from SM prediction. If the new scalar is too heavy to be directly produced, invisible decays of the SM-like Higgs boson can be the only signature for DM production.

The first ILC running stage will clearly focus on the Higgs boson measurements. At 250 GeV its production is dominated by the Higgs-strahlung process, $e^+e^- \rightarrow ZH$. In this channel the so called “Z-tagging” method can be used for unbiased selection of Higgs production events by reconstructing the recoil mass. This allows for fully model independent analysis avoiding any dependence on the Higgs decay channel. The recoil mass technique results also in high sensitivity to invisible Higgs boson decays [3, 4]. For 2 ab^{-1} of data collected at 250 GeV ILC the expected 95% C.L. limit on the invisible Higgs boson decays is 0.23% [5], an order of magnitude better than the estimated HL-LHC sensitivity. The expected recoil mass distribution for the ILC running at 250 GeV, assuming an invisible Higgs boson branching ratio of 10%, is shown in Fig. 1 (left). If the second scalar mass eigenstate is also light, it can be produced in e^+e^- collisions in the same way as the SM-like Higgs boson and should be visible as an additional resonance in the recoil mass distribution. Such scenarios can be probed at the ILC down to $\sin^2 \theta \sim 10^{-2}$, see Fig. 1 (right) [6].

3 Mono-photon events

DM particles can be pair produced in the e^+e^- collisions via exchange of a new mediator particle, Y , which couples to both electrons and DM particles, χ . This process can be detected,




Figure 1: Left: recoil mass distribution after event selection for ILC running at 250 GeV and invisible Higgs branching ratio of 10% [5]. Right: expected sensitivity of the International Large Detector (ILD) experiment at the ILC, for ILC running at 500 GeV to the scalar sector mixing parameter $\sin^2 \theta$ [6].




Figure 2: The expected sensitivity of the ILD experiment, for ILC running at 500 GeV, to DM pair production. Left: EFT limits on mediator mass scale Λ as a function of the DM particle mass M_χ , for different mediator types [7]. Right: mediator coupling limits as a function of the vector mediator mass M_Y for DM particle mass $M_\chi=50$ GeV [8].

if additional hard photon radiated from the initial state, $e^+e^- \rightarrow \chi\chi + \gamma$, is observed in the detector. The mono-photon signature is considered to be the most general approach to search for DM particle production at e^+e^- colliders, as it is largely independent of the BSM scenario. Signal events have a striking signature: single high-energy photon and no other activity in the detector. “Irreducible” background in this channel comes from the radiative neutrino pair-production, $e^+e^- \rightarrow \nu\bar{\nu} + \gamma$, while huge background from the radiative Bhabha scattering, $e^+e^- \rightarrow e^+e^- + \gamma$, can be efficiently suppressed by the selection cuts.

Scenarios with DM pair-production via heavy mediator exchange were considered in a full simulation study [7], using an EFT approach and assuming coupling values $\mathcal{O}(1)$. For light DM particles mediator masses up to about 3 TeV can be excluded at the ILC, as shown in Fig. 2 (left). DM production via light mediator exchange is still not excluded if very small mediator couplings to SM particles (electrons) are considered. Most general, “experimental-like” approach to such scenarios was presented in [8], where limits on the DM pair-production




Figure 3: An experimental setup proposed for new light particle searches at the ILC [10].




Figure 4: Expected sensitivity of main beam dump experiments at the 250 GeV ILC after one year (red lines) and 20 years (black lines) of running to processes with light axion-like particle production (left) and light scalar production (right). Results taken from [10].

were studied as a function of mediator mass and width. Dedicated simulation procedure was used [9], with all “detectable” photons generated on a Matrix Element level, matched with soft Initial State Radiation (ISR). With 2-dim distribution of $(p_T^\gamma, \eta^\gamma)$ used to constrain DM production, ILC running at 500 GeV is sensitive to mediator couplings down to $\mathcal{O}(10^{-3})$, as shown in Fig. 2 (right).

4 Fixed target experiments

Extreme intensities of electron and positron beams at the ILC, $\mathcal{O}(10^{22})$ particles per year, open unique options for fixed-target experiments focused on rare processes. A general concept of main beam dump experiments searching for axion-like particles or new scalars is presented in Fig. 3 [10]. Light axion-like particles (a) could be copiously produced in the beam dump and, if sufficiently long-lived, decay only after passing the massive muon shield. Expected sensitivity at ILC running at 250 GeV to $a\gamma\gamma$ coupling as a function of the axion-like particle mass, resulting from the search for two-photon decay, $a \rightarrow \gamma\gamma$, is presented in Fig. 4 (left).




Figure 5: Expected sensitivity of main beam dump experiments at the 250 GeV ILC after one year (red lines) and 20 years (black lines) of running to processes with dark photon production with visible dark photon decays (left) [11] and invisible decays to DM particles (right) [13].

Thanks to the large beam intensities, the sensitivity of the ILC beam dump experiments is an order of magnitude better than those expected at other projects. Similar increase in sensitivity is expected for models with new light scalars coupling to charged leptons [10, 11]. Expected sensitivity at the ILC to the scenario with scalar coupling proportional to the lepton mass is shown in Fig. 4 (right).

Dark photons (A') could be produced in the ILC beam dump by pair-annihilation and bremsstrahlung processes. For visible decay channels, expected sensitivity of the ILC beam dump experiment to A' production is comparable with dedicated future projects [11]. If DM particles are light, invisible A' decays to DM particles (χ), $A' \rightarrow \chi\chi$, can dominate. Dark photon production in the beam dump would then result in a collimated stream of DM particles from A' decay, which can be probed by looking for elastic DM interactions in the dedicated detector. This approach, previously used in SLAC Beam Dump Experiment E137 [12], will offer huge improvement in sensitivity for $m_{A'} \lesssim 1$ GeV thanks to much higher statistics at the ILC [13].

Experiments with extracted positron beams can search for dark photon production independent on its decays [14]. Associated production, $e^+e^- \rightarrow A' \gamma$ can be searched for with thick active target reconstructing the missing energy (idea based on LDMX experiment for SLAC [15]) or in thin target detectors reconstructing the missing mass (as in the PADME experiment at Frascati [16]). In both approaches, sensitivity of the fixed-target experiments at the ILC can probe the dark photon couplings down to the minimum couplings allowed by relic density measurements.

5 Conclusions

ILC will offer many complementary options for DM searches. Different scenarios can be constrained via precision Higgs studies. Direct searches for DM pair-production are possible in the mono-photon channel with sensitivity to heavy mediator exchange reaching TeV mass scales, one order of magnitude higher than the collision energy, and sensitivity to couplings of $\mathcal{O}(10^{-3})$

in the case of light mediator exchange. The ILC will also offer highest energy electron and positron beams, with unprecedented intensities, for beam dump and extracted beam experiments. Fixed-target experiments offer many interesting opportunities for dark sector searches in the low mass domain.

Acknowledgements

This study was supported by the National Science Centre, Poland, the OPUS project under contract UMO-2017/25/B/ST2/00496 (2018-2021) and the HARMONIA project under contract UMO-2015/18/M/ST2/00518 (2016-2021), and by the German Research Foundation (DFG) under grant number STO 876/4-1 and STO 876/2-2.

References

- [1] Wojciech Kotlarski and Aleksander Filip Zarnecki. “Probing dark matter with ILC”. *Proceedings of The European Physical Society Conference on High Energy Physics — PoS(EPS-HEP2021)*. Vol. 398. 2022, p. 135. DOI: [10.22323/1.398.0135](https://doi.org/10.22323/1.398.0135).
- [2] Wojciech Kotlarski and Aleksander Filip Zarnecki. “Probing dark matter at the ILC”. *Proceedings of Particles and Nuclei International Conference 2021 — PoS(PANIC2021)*. Vol. 380. 2022, p. 124. DOI: [10.22323/1.380.0124](https://doi.org/10.22323/1.380.0124).
- [3] Yu Kato. “Probing the dark sector via searches for invisible decays of the Higgs boson at the ILC”. *Proceedings of European Physical Society Conference on High Energy Physics — PoS(EPS-HEP2019)*. Vol. 364. 2020, p. 358. DOI: [10.22323/1.364.0358](https://doi.org/10.22323/1.364.0358). arXiv: [2002.12048](https://arxiv.org/abs/2002.12048) [[hep-ex](#)].
- [4] Amanda Steinhebel, Jim Brau, and Chris Potter. “H→invisible at the ILC with SiD”. *International Workshop on Future Linear Colliders*. Apr. 2021. arXiv: [2105.00128](https://arxiv.org/abs/2105.00128) [[hep-ex](#)].
- [5] Yu Kato. “Probing the dark sector via searches for invisible decays of the Higgs boson at the ILC”. *PoS EPS-HEP2019* (2020), p. 358. DOI: [10.22323/1.364.0358](https://doi.org/10.22323/1.364.0358). arXiv: [2002.12048](https://arxiv.org/abs/2002.12048) [[hep-ex](#)].
- [6] Yan Wang, Mikael Berggren, and Jenny List. “ILD Benchmark: Search for Extra Scalars Produced in Association with a Z boson at $\sqrt{s} = 500$ GeV” (May 2020). arXiv: [2005.06265](https://arxiv.org/abs/2005.06265) [[hep-ex](#)].
- [7] Moritz Habermehl, Mikael Berggren, and Jenny List. “WIMP Dark Matter at the International Linear Collider”. *Phys. Rev. D* 101.7 (2020), p. 075053. DOI: [10.1103/PhysRevD.101.075053](https://doi.org/10.1103/PhysRevD.101.075053). arXiv: [2001.03011](https://arxiv.org/abs/2001.03011) [[hep-ex](#)].
- [8] Jan Kalinowski et al. “Sensitivity of future linear e^+e^- colliders to processes of dark matter production with light mediator exchange”. *Eur. Phys. J. C* 81.10 (2021), p. 955. DOI: [10.1140/epjc/s10052-021-09758-6](https://doi.org/10.1140/epjc/s10052-021-09758-6). arXiv: [2107.11194](https://arxiv.org/abs/2107.11194) [[hep-ph](#)].
- [9] J. Kalinowski et al. “Simulating hard photon production with WHIZARD”. *Eur. Phys. J. C* 80.7 (2020), p. 634. DOI: [10.1140/epjc/s10052-020-8149-6](https://doi.org/10.1140/epjc/s10052-020-8149-6). arXiv: [2004.14486](https://arxiv.org/abs/2004.14486) [[hep-ph](#)].

- [10] Yasuhito Sakaki and Daiki Ueda. “Searching for new light particles at the international linear collider main beam dump”. *Phys. Rev. D* 103.3 (2021), p. 035024. DOI: [10.1103/PhysRevD.103.035024](https://doi.org/10.1103/PhysRevD.103.035024). arXiv: [2009.13790](https://arxiv.org/abs/2009.13790) [hep-ph].
- [11] Kento Asai et al. “New physics searches at the ILC positron and electron beam dumps”. *JHEP* 09 (2021), p. 183. DOI: [10.1007/JHEP09\(2021\)183](https://doi.org/10.1007/JHEP09(2021)183). arXiv: [2105.13768](https://arxiv.org/abs/2105.13768) [hep-ph].
- [12] Brian Batell, Rouven Essig, and Ze’ev Surujon. “Strong Constraints on Sub-GeV Dark Sectors from SLAC Beam Dump E137”. *Phys. Rev. Lett.* 113.17 (2014), p. 171802. DOI: [10.1103/PhysRevLett.113.171802](https://doi.org/10.1103/PhysRevLett.113.171802). arXiv: [1406.2698](https://arxiv.org/abs/1406.2698) [hep-ph].
- [13] D. Ueda, *Fixed-target experiment using ILC main beam dump*, presented at the ILC Workshop on Potential Experiments (ILCX2021), [link to contribution](#).
- [14] M. Perelstein, *First estimates of the dark sector capabilities at ILC*, presented at the 2021 International Workshop on Future Linear Colliders (LCWS2021), [link to contribution](#).
- [15] L. Marsicano et al. “Novel Way to Search for Light Dark Matter in Lepton Beam-Dump Experiments”. *Phys. Rev. Lett.* 121.4 (2018), p. 041802. DOI: [10.1103/PhysRevLett.121.041802](https://doi.org/10.1103/PhysRevLett.121.041802). arXiv: [1807.05884](https://arxiv.org/abs/1807.05884) [hep-ex].
- [16] Andre Frankenthal. “Searching for Dark Photons with PADME”. *Meeting of the Division of Particles and Fields of the American Physical Society*. Oct. 2019. arXiv: [1910.00764](https://arxiv.org/abs/1910.00764) [physics.ins-det].