

hep-ph/0507244, 0603030, 0606169, 0611204

A Global Fit to Scattering Data with NLL BFKL Resummation


Chris White, NIKHEF

DIS2007 - April 18th 2006

Overview

- ▶ Solution of the [BFKL](#) equation at NLL order with running coupling.
- ▶ Comparison of gluon splitting function with other resummation approaches ([ABF](#), [CCSS](#)).
- ▶ Treatment of Heavy Flavours.
- ▶ Global parton fit using NLL resummed coefficient and splitting functions!

The High Energy Problem



- Coefficient and splitting functions for the proton structure functions unstable at low x due to terms

$$\sim x^{-1} \bar{\alpha}_S^n \log^m(1/x),$$

$$m \leq n - 1.$$

- Divergence due to t -channel gluon exchange at LL order, with some quark mixing at NLL order.
- Must resum the gluon 4-point function by solving the [BFKL](#) equation.
- Relate gluon to structure functions using the k_T factorisation formula ([Collins & Ellis](#); [Catani, Ciafaloni & Hautmann](#)).

Running coupling solution of BFKL equation

- Mellin moments:

$$f(\gamma, N) = \int_0^\infty (k^2)^{-\gamma-1} \int_0^1 dx x^N f(x, k^2)$$

- Substitute LO running coupling into BFKL equation (Collins & Kwiecinski):

$$\begin{aligned} \frac{d^2 f(\gamma, N)}{d\gamma^2} &= \frac{d^2 f_l(\gamma, Q_0^2)}{d\gamma^2} - \frac{1}{\bar{\beta}_0 N} \frac{d(\chi_0(\gamma) f(\gamma, N))}{d\gamma} \\ &\quad + \frac{\pi}{3\bar{\beta}_0^2 N} \chi_1(\gamma) f(\gamma, N), \end{aligned}$$

with $\bar{\beta}_0 = 3/(\pi\beta_0)$.

- Solve with ansatz:

$$f(N, \gamma) = \exp\left(-\frac{X_1(\gamma)}{\bar{\beta}_0 N}\right) \int_\gamma^\infty A(\tilde{\gamma}) \exp\left(\frac{X_1(\tilde{\gamma})}{\bar{\beta}_0 N}\right) d\tilde{\gamma}$$

(Ciafaloni & Colferai).

- ▶ Can shift lower limit $\gamma \rightarrow 0$ up to power-suppressed corrections (Thorne).
- ▶ Gluon factorises:

$$\mathcal{G}(N, t) = \mathcal{G}_E(N, t) \mathcal{G}_I(Q_0^2, N)$$

$$(t = \log Q^2/\Lambda^2).$$

- ▶ Perturbative piece:

$$\mathcal{G}_E^1(N, t) = \frac{1}{2\pi i} \int_{1/2-i\infty}^{1/2+i\infty} \frac{f^{\beta_0}}{\gamma} \exp [\gamma t - X_1(\gamma, N)/(\bar{\beta}_0 N)] d\gamma$$

with:

$$X_1(\gamma, N) = \int_{\frac{1}{2}}^{\gamma} \left[\chi_0(\tilde{\gamma}) + N \frac{\chi_1(\tilde{\gamma})}{\chi_0(\tilde{\gamma})} \right] d\tilde{\gamma}.$$


- ▶ Similarly, get structure functions:

$$\mathcal{F}_E^1(N, t) = \frac{1}{2\pi i} \int_{1/2-i\infty}^{1/2+i\infty} \frac{h(\gamma, N) f^{\beta_0}}{\gamma} \exp [\gamma t - X_1(\gamma, N)/(\bar{\beta}_0 N)] d\gamma$$

- ▶ If impact factors known, can disentangle all resummed coefficient and splitting functions (within a particular factorisation scheme).
- ▶ However, NLL impact factors $h(\gamma, N)$ not known. Work in progress ([Bartels, Colferai, Gieseke & Kyrieleis](#)).
- ▶ Instead LL factors with exact gluon kinematics have been calculated ([Bialas, Navelet & Peschanski](#); [White, Peschanski & Thorne](#)).
- ▶ These provide a very good estimate to the full NLL impact factors ([White & Thorne](#)).
- ▶ Can use these to calculate all the NLL resummed coefficient and splitting functions in the DIS scheme.
- ▶ Finally, combine resummed results with NLO DGLAP:


$$P^{tot.} = P^{NLL} + P^{NLO} - \left[P^{NLL(0)} + P^{NLL(1)} \right]$$

Results for Splitting Functions




- ▶ Results shown at $n_f = 4$, $t = 6$.
- ▶ Running coupling suppresses low x divergence.
- ▶ NLL kernel and impact factor effects lead to even more suppression.
- ▶ Main feature is a dip below the NLO DGLAP result.

Comparison with Alternative Approaches



- ▶ The **ABF** and **CCSS** groups calculate $P_+ (= P_{gg} + \frac{4}{9}P_{qg})$ with $n_f = 0$.
- ▶ Results agree closely...
- ▶ Now need to consider heavy flavour coefficients...

Variable Flavour Number Schemes





- ▶ In DIS, can produce final state heavy quarks by boson gluon fusion ([Witten](#)).
- ▶ Diagrams diverge as $Q^2 \rightarrow \infty$ due to terms $\sim \alpha_S^n \log^n M^2/Q^2$.
- ▶ Get round this by defining parton distributions for the heavy species above a suitable matching scale e.g. $Q^2 = M^2$.
- ▶ Matching conditions exist at this scale ([Buza](#), [Matiounine](#), [Smith & van Neerven](#)) between the high Q^2 and low Q^2 partons.


The DIS(χ) Scheme

- ▶ Both sets of partons are ambiguous - one must fix a particular variable flavour scheme according to two types of choice.
- ▶ Have developed a scheme that allows one to disentangle the meaning of the impact factors in terms of heavy flavour coefficient and matching functions.
- ▶ Called the DIS(χ) scheme by analogy with the DIS scheme for massless quarks.
- ▶ Allows the consistent implementation of small x resummations in the heavy flavour sector.
- ▶ Approximate results obtained for massive resummed quantities at NLL+NLO order.
- ▶ Thus have everything necessary for a global parton fit!

Quality of Fit



- ▶ Neutral and charged current data for F_2^P (including F_2^c) from [H1](#), [ZEUS](#).
- ▶ Data for F_2^P , F_2^n from [BCDMS](#), [NMC](#), [SLAC](#) and [E665](#).
- ▶ $F_{2,3}^{\nu(\bar{\nu})N}$ from [CCFR](#). F_2^D/F_2^P from [NMC](#).
- ▶ Non-DIS: DY data from [E866](#) / [NuSea](#); DY asymmetry from [NA51](#); $\sigma_{DY}^{PD}/\sigma_{DY}^{pp}$ from [E866](#); W asymmetry from [CDF](#).
- ▶ NLL resummed fit gives an overall fit quality $\chi^2 = 2249$ for 2181 data points.
- ▶ Compare NLO DIS scheme $\chi^2 = 2352$ and $\overline{\text{MS}}$ scheme $\chi^2 = 2307$.
- ▶ A previous LL resummed fit gave $\chi^2 = 2336$, with significant momentum conservation violation.
- ▶ Main improvement in the HERA data, as expected.
- ▶ Description of F_2^c benefits from DIS(χ) scheme.
- ▶ Resummation seems to decrease tension between data sets.

Results - F_2 


Results - F_2^c 

- ▶ Resummed fit performs better for small x data - note slope as Q^2 increases.
- ▶ Fit is also improved over the whole range of x .
- ▶ Resummed F_2^c at lower end of range allowed by data.

Comparison with Tevatron jet data




Gluon Distribution




- ▶ Gluons differ for $x \lesssim 10^{-2}$.
- ▶ NLL resummed gluon positive and growing at small x !
- ▶ Not true at fixed order.
- ▶ Positive gluon avoids negative structure functions.
- ▶ See this in e.g. F_L ...

Longitudinal Structure Function




- ▶ NLO result goes negative.
- ▶ LL prediction more stable but turns over due to negative gluon. Also inconsistent with NLO at high x .
- ▶ NLL result is much more sensible!

Longitudinal Structure Function



- ▶ Clearly see perturbative instability in fixed order results.
- ▶ This is cured by the resummation.

Reduced Cross-Section



- Turnover required by data at low x (high y) - NLO fails.
- Resummation helps!
Interesting to compare with NNLO.

Conclusions

- ▶ Have combined NLL resummations with a NLO fixed order QCD expansion in massive and massless sectors, including running coupling effects.
- ▶ Running coupling and NLL kernel corrections each lead to large suppressions in the small x divergence. NLL impact factors give further suppression.
- ▶ Global fit with improved splitting and coefficient functions gives an excellent description of data.
- ▶ Gluon from resummed fit positive at low x and Q^2 .
- ▶ Prediction for F_L is stable in this regime.

⇒ Very compelling evidence for BFKL effects!