



Contribution ID: 53

Type: **Session: Dark Matter and Beyond Standard Model Searches**

CMB spectral μ -distortion during phase transition in the Bound Dark Matter (BDM) model

Thursday, 26 November 2020 13:30 (30 minutes)

Spectral distortions of the CMB provide independent and complementary probes to study energy injection processes in the early universe. In this work was obtained the spectral μ -distortion associated with phase transition of the dark particles in the BDM model. Several scenarios were simulated numerically with parameters: $a_c = [4.9 \times 10^{-7}, 3.3 \times 10^{-6}]$, $f_{eff} = [0, 1]$, $v_c = [0, 0.71]$, $f_x = 10^{-2}, 10^{-3}, 10^{-4}, 10^{-5}$ and 10^{-6} . Some constrictions were obtained for FIRAS sensitivity for $a_c < 1.68 \times 10^{-6}$ ($f_x = 10^{-2}$, $f_{eff} = 0.5$) and for PIXIE sensitivity $a_c > 1 \times 10^{-6}$ ($f_x = 10^{-5}$, $f_{eff} = 0.5$).

Primary authors: HENRÍQUEZ, Raúl (Universidad de El Salvador); MASTACHE, Jorge (Mesoamerican Centre for Theoretical Physics); CABALLERO, Karen (Universidad Autónoma de Chiapas)

Presenter: HENRÍQUEZ, Raúl (Universidad de El Salvador)

Session Classification: Dark Matter and Beyond Standard Model Searches