



Detecting the Δ^+ 1232 Baryon from an Electron-Proton Inelastic Scattering

Nations' Flying Foxes Proposal presentation


About us

- International school of Geneva- Campus des Nations
- Student body of 1000, with 113 nationalities (8 of which are represented in our team!)
- Born in 2002-2004, currently in our senior year of high school
- University plans ranging from physics and engineering to law and economics




Rationale

1. Detecting short-lived particle by studying less massive particles that come out of high-energy collisions.
2. “Peak” into a proton. What are its properties, otherwise unobservable ?




Background Theory




(Fig.1) // Source - W. BARTEL, B. DUDELZAKI, H. KREHBIEL, J. MCELROY, U. MEYER-BERKHOUT, W. SCHMIDT, V. WALTHER ^{ttt} and G. WEBER
*, ELECTROPRODUCTION OF PIONS NEAR THE A(1236) ISOBAR AND THE FORM FACTOR C*M(q²) OF THE (gammaNA)-VERTEX - Deutsches Elektronen-Synchrotron DESY, Hamburg, Germany and II. Institut für Experimentalphysik der Universität Hamburg, Germany*

(Fig.3) Synthesis of a Neutron and a Pion Plus from a decaying Delta plus



(Fig.2) Inelastic scattering of a Proton with an Electron



Equation 1 : $m_0(N^+) + KE(e^- \text{ initial}) = m_0(\Delta^+) + KE(e^- \text{ final}) = m_0(N^0) + m_0(\pi^+) + KE(\pi^+) + KE(e^- \text{ final})$

$\Rightarrow m_0(\Delta^+) = m_0(N^+) + KE(e^- \text{ initial}) - KE(e^- \text{ final}),$

$\Rightarrow m_0(\Delta^+) = m_0(N^0) + m_0(\pi^+) + KE(\pi^+)$, where $m_0(\pi^+) + KE(\pi^+)$ gives the total energy of π^+ .

Equation 2 : $p(e^- \text{ initial}) = p(\pi^+) + p(e^- \text{ final})$

Rectangular

Hypothesis



Latest Update:

Energy gained by p^+ during inelastic scattering = $x - y$ GeV

, where x = initial energy of electron in GeV and y = final energy of electron in GeV

As a result, p^+ gains $m(?) - 0.938$ GeV c^{-2} of mass

, where $m(?)$ = rest mass of particle created in GeV c^{-2} ,
and 0.938 = rest mass of proton in GeV c^{-2} .

KE gained by ? = $x - y - m(?) + 0.938$ GeV

As relativistic KE of ? = $m(?)C^2(\gamma - 1)$, where $\gamma = \frac{1}{\sqrt{1 - \frac{v^2}{c^2}}}$ and $C = 1$,

$$v(?) = \sqrt{1 - \frac{m^2(?)}{(x - y + 0.938)^2}} \text{ in ms}^{-1}$$

Now,

$$p(?) = \frac{KE + m(?)c^2}{c^2} \cdot v(?)$$

$$p(?) = (x - y + 0.938) \sqrt{1 - \frac{m^2(?)}{(x - y + 0.938)^2}}$$

, where $p(?)$ = momentum of particle created in $\text{GeV}c^{-1}$.

Solving for $m(?)$ gives:

$$m(?) = \sqrt{(x - y + 0.938)^2 - p^2(?)}$$

- Using the law of conservation of momentum,

$$p(?) = \frac{-\beta \sin \theta}{\sin \phi}, \quad p(?) = \frac{\alpha - \beta \cos \theta}{\cos \phi}$$

, where β = final momentum of the electron in $\text{GeV}c^{-1}$, θ = angle of deflected electron, ϕ = angle of particle created, and α = initial momentum of the electron in $\text{GeV}c^{-1}$.

Hence equating the expressions for $p(?)$ gives

$$\phi = \arctan\left(\frac{-\beta \sin \theta}{\alpha - \beta \cos \theta}\right)$$

And hence

$$p(?) = \frac{-\beta \sin \theta}{\sin \left[\arctan\left(\frac{-\beta \sin \theta}{\alpha - \beta \cos \theta}\right) \right]}$$


Finally,

$$m(?) = \sqrt{(x - y + 0.938)^2 - \left[\frac{-\beta \sin \theta}{\sin \left[\arctan \left(\frac{-\beta \sin \theta}{\alpha - \beta \cos \theta} \right) \right]} \right]^2}$$

So, measuring x , y , α , β , and θ of the electron gives us the rest mass of the particle created.


If $m(?) = 1.232 \text{ GeV}c^{-2}$, we have found Δ^+ 1232 baryon!

B: \otimes



- 1: Electron beam shaft
- 2: Collimator
- 3: Electron beam trajectory
- 4: Lead panel (proton source)
- 5: Pi-plus projected trajectory

- 6: Scintillator (trigger mechanism)
- 7: Two pairs of MicroMegas detectors
- 8: PCMAG
- 9: Lead crystal calorimeter



source: Cristovao Beirão Da Cruz E Silva