

Introduction and some first FCC-ee flavour studies

Donal Hill

23/9/20

FCC & LHCb software discussion





A short introduction

- Thank you all for coming!
- I approached Clement and the FCC software team a short while ago to voice an interest in flavour studies at FCC-ee
 - Have had lots of help getting up and running with FCCSW to simulate $e^+e^- \rightarrow Z^0 \rightarrow b\bar{b}$ with Pythia + Delphes
- Will show some studies using *awkward array*
- However, detailed analysis will likely require:
 - Robust candidate selection framework for **full decay chains**
 - **Tupling** of full decay chain with a **range of quantities calculated**
 - **MC-matching** of full chain to understand signal efficiencies
- Begin discussions today about how LHCb Run 3 software developments could be of use

Basic features of $Z^0 \rightarrow b\bar{b}$ events

b -quark truth-level kinematics

- At the Z^0 -pole, expect Z^0 production basically at rest so b -quarks should be produced back-to-back in lab frame
- Plot lab frame η and ϕ distribution of the two b -quarks
 - η equal and opposite, ϕ differs by π - consistent with back-to-back production



B-hadron production fractions

- Look at the *genParticles* container and count fraction of each true particle type produced according to *pdgId*
- Production of B_c^+ mesons is a nice feature at FCC-ee - no B_c^+ produced at Belle II

<i>B</i> -hadron	Production fraction (%)
B^0	43.0
B^\pm	43.0
B_s^0	9.6
B_c^\pm	0.04
Λ_b^0	3.7


Simple selection example

Particle combinations

- Access to stable final-state hadrons via the *pfcharged* container
- Perform particle combinations per-event using awkward array with ROOT file loading using uproot
- These packages provide *numpy*-like access to jagged data from ROOT files
 - Nice for dealing with events containing different numbers of particles in each event
 - Solid support within the HSF and PyHEP communities
- Particle combinatorics using *ak.combinations* (same particle type) and *ak.cartesian* (different particle types)

Example decay: $B^- \rightarrow (D^0 \rightarrow K^- \pi^+) \pi^-$

- High-statistics mode at LHCb which forms part of the CKM angle γ measurements
- Related decays $B^- \rightarrow (D^0 \rightarrow K^- \pi^+) \pi^-$ and $B^- \rightarrow (D^0 \rightarrow K^- \pi^+) K^-$ (note K charge w.r.t. B) sensitive to γ



2011 – 2016 LHCb data [LHCb-PAPER-2017-021]

Expected yield at FCC-ee


Factor	Value
Z^0 produced at FCC-ee	5×10^{12}
$\mathcal{B}(Z^0 \rightarrow b\bar{b})$	0.15
B from either b or \bar{b}	2
B^\pm production fraction	0.43
$\mathcal{B}(B^- \rightarrow D^0\pi^-)$	4.7×10^{-3}
$\mathcal{B}(D^0 \rightarrow K^-\pi^+)$	0.04
Total yield	121.3 million

- The yield above is for 7.5×10^{11} $b\bar{b}$ pairs
- My sample contains 13 million $Z^0 \rightarrow b\bar{b}$, so expect **2100** $B^- \rightarrow D^0\pi^-$ decays in the generated sample¹

¹Charge conjugation implied


D^0 candidates in inclusive $Z^0 \rightarrow b\bar{b}$ sample

- Invariant mass distribution of $D^0 \rightarrow K^\mp \pi^\pm$ candidates built from kaons and pions in the *pfcharged* container
- Select K and π based on their *pdgId* values (assumes perfect PID performance, in reality will have some mis-ID)
- Apply $p > 4$ GeV cut to all candidate tracks - no vertex fits e.t.c.
 - Some background present, but pretty clean given no dedicated selection is in place



B^\pm candidates in inclusive $Z^0 \rightarrow b\bar{b}$ sample

- Take $K\pi$ combinations passing a ± 25 MeV mass window around PDG D^0 mass, and combine them with pion candidates
- Encouraging to see a well-resolved B peak and low background level with minimal selection applied
- Peaks at low mass are due to partially reconstructed $B \rightarrow D^*\pi$ decays (studied in detail at LHCb, see previous slide)



Rough yield estimate (no mass fit)

- $B^- \rightarrow D^0\pi^-$ peak is 330 events high (background-subtracted) and 5 bins wide
- Assume a triangle to give $N = 330 \times 5 \times 0.5 = 825$
- This is **40%** of the total expected $B^- \rightarrow D^0\pi^-$ yield in the 13M $Z^0 \rightarrow b\bar{b}$ generated sample
- Includes hadron efficiency estimates in the IDEA Delphes card [here](#) and the $p > 4$ GeV track cuts
- This would give a total $B^- \rightarrow D^0\pi^-$ yield of ~ 50 million in the FCC-ee data
 - Yield in 9 fb^{-1} LHCb Run 1 + 2 sample is ~ 2 million after all selection, **so FCC-ee about an order higher**²

²LHCb Upgrade I + II targets 300 fb^{-1} , which would surpass this

Software moving forward

Software requirements for flavour studies


- Ability to generate samples of **exclusive decays using EvtGen**
 - Work required in FCCSW to configure EvtGen appropriately
- General candidate building tools at the FCCSW level (rather than at user-level as I have done with *awkward*)
 - Use not only kinematic information but track fit and vertex fit information
 - Include MC-matching of the full decay chain
 - **Can LHCb Run 3 software help here?**
- Ability to study how often the entire event can be reconstructed
 - If one side of the $Z^0 \rightarrow b\bar{b}$ event can be fully reconstructed, this **gives the 4-vector of the other side** (given well-known \sqrt{s})
 - Valuable for modes with missing momentum - same ideas adopted at Belle II

Summary


- Many thanks to Clement and co. for the help in getting started with FCCSW and simulating events
- Early studies indicate:
 - Sensible rates of B -hadron production
 - High selection efficiencies for hadronic modes
 - Low background levels with minimal selections applied
- Ability to study more decay modes, and in more detail, will help guide detector design requirements
- Porting code and ideas from the LHCb selection and nTupling frameworks could be profitable for these studies
 - Hence todays discussion!
 - **Should also keep in mind FCC-hh, where rates will be much higher**

Backup


$m(\pi\pi)$ spectrum - ρ visible




$m(KK)$ spectrum - ϕ and D^0





$m(\mu\mu)$ spectrum - J/ψ



$m(ee)$ spectrum - J/ψ and γ conversion



True particle multiplicities in 1 million events



Generating inclusive $Z^0 \rightarrow b\bar{b}$

- Setup FCCSW from [here](#)
- Used *ee_Z_ddbar.cmd* to create a *ee_Z_bbbar.cmd* Pythia config file, and run Pythia with Delphes using *PythiaDelphes_config.py*
- Generation is inclusive, in that the b -quarks produced hadronise to all sorts
 - *EvtGen* config options have been added to FCCSW, but early tests indicate that it isn't generating purely the decays specified by the user in a *.dec* file
 - Under discussion [here](#)
- Have generated 13 million events for the work shown today

Python analysis software for performance studies

- Initial studies using *uproot* for jagged ROOT file loading and *awkward* for manipulation
- Both packages are well-supported (Jim Pivarski main author) and many tutorials
 - Analysed 13 million events in under 5 mins to produce the D^0 and B^\pm candidates, making combinations across all tracks in every event
 - Interface seamlessly with other industry-standard packages like *numpy* for calculations and *matplotlib* for plotting
 - The *vector* package (Henry Schreiner) is being worked on - will provide support for Lorentz vectors e.t.c. in the *awkward* format
- FCC studies can make use of these packages, with perhaps a common set of helper functions
 - No need to reinvent the wheel with these tools in existence